

# KENTUCKY OPIOID REPLACEMENT STUDY

2022 ANNUAL OUTCOME REPORT



# PROJECT ACKNOWLEDGMENTS

## PRESENTED BY:

Kentucky Department for Behavioral  
Health, Developmental and Intellectual  
Disabilities  
Division of Behavioral Health  
275 East Main St.  
Frankfort, KY 40621-0001  
(502) 564-4527

### **Wendy Morris**

*Commissioner*  
Department for Behavioral Health, Developmental  
and Intellectual Disabilities

### **Koleen Slusher**

*Director*  
Division of Behavioral Health

### **Maggie Schroeder**

*Program Manager*  
Substance Abuse Treatment and Recovery Services

### **Karla Thompson**

*Program Administrator*  
State Opioid Treatment Authority

## PREPARED BY:

University of Kentucky, Center on Drug  
and Alcohol Research  
333 Waller Ave, Suite 480  
Lexington, KY 40504  
<http://cdar.uky.edu/kortos>

Suggested citation: Logan, TK, Scrivner,  
A., & Cole, J. (2022). *Kentucky's Opioid  
Replacement Treatment Program  
Outcome Study 2022 Annual Report*.  
Lexington, KY: University of Kentucky,  
Center on Drug & Alcohol Research.

The 2022 KORTOS report includes data  
from 34 clients at Kentucky opioid  
treatment programs (OTPs) who  
completed both an intake interview  
between January 1, 2020 and December  
31, 2020 and a six-month follow-up  
interview targeted between July 2020 and  
June 2021.

# TABLE OF CONTENTS

<b>Project Acknowledgments</b> .....	<b>2</b>
<b>Executive Summary</b> .....	<b>5</b>
<b>Introduction and Overview</b> .....	<b>11</b>
<b>Section 1. KORTOS Client Characteristics</b> .....	<b>15</b>
Description of Kortos Clients at Treatment Intake.....	15
KORTOS Follow-up Sample .....	29
<b>Section 2. Substance Use</b> .....	<b>32</b>
Alcohol And/or Drug Use .....	33
Any Illegal Drugs.....	35
Prescription Opioid Misuse .....	37
Non-prescribed Use of Methadone.....	38
Non-prescribed Use of Bup-nx.....	39
Heroin .....	40
Non-opioid Drug Use .....	41
Injection Drug Use.....	43
Alcohol Use .....	43
Problems Experienced with Substance Use in the Past 30 Days.....	45
Readiness for Substance Abuse Treatment.....	45
Substances That Client Had the Most Trouble with .....	46
Self-reported Severity of Alcohol and Drug Use.....	46
Substance Abuse Treatment History.....	50
Tobacco Use.....	51
<b>Section 3: Mental and Physical Health</b> .....	<b>56</b>
Depression Symptoms.....	56
Anxiety Symptoms.....	57
Comorbid Depression and Anxiety Symptoms.....	59
Suicidal Ideation And/or Attempts.....	59
Self-medication for Mental Health Symptoms.....	60
Post-traumatic Stress Disorder.....	61
General Health Status.....	61
Chronic Pain.....	65
Health Insurance.....	67
Quality of Life Ratings .....	68
<b>Section 4. Criminal Justice System Involvement and Interpersonal Victimization</b>	<b>70</b>
Arrests.....	70
Incarceration .....	71
Criminal Justice System Supervision .....	71
Interpersonal Victimization Experiences .....	72
Worry About Personal Safety .....	72

<b>Section 5. Education, Economic Status, and Living Circumstances.....</b>	<b>74</b>
Education .....	74
Employment.....	74
Living Circumstances .....	79
Economic Hardship .....	81
<b>Section 6. Recovery Supports .....</b>	<b>84</b>
Mutual Help Recovery Group Meeting Attendance .....	84
Number of People Client Can Count on for Recovery Support.....	84
Chances of Staying Off Drugs/alcohol .....	85
What Will Be Most Useful in Staying Off Drugs/alcohol .....	85
<b>Section 7. Multidimensional Recovery.....</b>	<b>87</b>
<b>Section 8. Client Satisfaction with Opioid Treatment Programs.....</b>	<b>89</b>
Client Involvement in the Program .....	89
Recommendation to the Program.....	89
Client Ratings of Program Experiences .....	89
Positive and Negative Aspects of Program.....	91
<b>Section 9. Conclusion and Implications.....</b>	<b>93</b>
Who Do the Opioid Treatment Programs Serve? .....	93
Areas of Success .....	94
Areas of Concern .....	96
Study Limitations.....	99
Conclusion .....	101
<b>Appendix A. Methods .....</b>	<b>102</b>
<b>Appendix B. Locating Efforts For The 2015 KORTOS Follow-Up Study .....</b>	<b>104</b>
<b>Appendix C. Client Characteristics At Intake For Those Who Completed A Follow-Up Interview And Those Who Did Not Complete A Follow-Up Interview.....</b>	<b>106</b>
<b>Appendix D. Trends In Prescription Opioid, Methadone, Bup-Nx, And Heroin Use Among Kortos Clients With A Completed Follow-Up Interview .....</b>	<b>113</b>
<b>Appendix E. Comparison Of Clients Who Were Currently Involved With The Clinic At Follow-Up And Those Who Were Not Involved On Targeted Factors.....</b>	<b>114</b>

## EXECUTIVE SUMMARY

Opioid treatment programs (OTPs) may play a unique and important role in addressing opioid abuse in Kentucky, where non-medical use of prescription opioids is a continuing health concern.<sup>1,2</sup> In 2007, Kentucky OTPs began collecting outcome data on opioid treatment programs. The outcome project is conducted in collaboration with the Kentucky Division of Behavioral Health and Narcotic Treatment Authority. The Kentucky Opioid Replacement Treatment Outcome Study (KORTOS) is an evidence-based data collection system designed to examine opioid treatment outcomes over time.

The goal of KORTOS is to examine client satisfaction and client outcomes for several targeted factors including: (1) substance use, (2) mental and physical

health, (3) criminal justice involvement, (4) quality of life, (5) education, economic status, and living situation, and (6) recovery supports. This report describes outcomes for 34 clients who: (1) attended one of twelve Kentucky OTPs eligible to participate in the study, (b) completed an intake interview between January 1, 2020 and December 31, 2020, (c) agreed to do the follow-up about 6 months later, and (d) completed a follow-up interview between July 1, 2020 and June 30, 2021. Of those individuals, 61.8% (n=21) were still involved in the treatment clinic at follow-up and 38.2% (n=13) were not involved in the treatment clinic at follow up. In previous years, clients who were no longer involved in the clinic were not eligible for the follow-up sample. However, beginning this report year, the decision was made to include these individuals in the eligible follow-up sample, but to analyze their outcomes separately. Therefore, for the follow-up analysis, only clients who reported that they were still involved with the clinic at follow-up (n = 21) were included; however, the 21 clients who were still involved in the clinic are compared to clients who completed a follow-up but were not still involved in the clinic at follow-up (n = 13) for

each section.<sup>3,4</sup>

### Who Do the Opioid Treatment Programs Serve?

Overall, in CY 2020, 192 clients from 6 of the 12 participating Kentucky OTPs completed the KORTOS intake interview.<sup>5</sup> Information from those intake interviews indicate that clients were an average of 39 years old ranging from 22 to 70 years old. Less than half (41.1%) were female and 58.3% were male. The majority of clients (60.9%) self-reported they decided get help on their own and 30.2% reported that they were referred to the OTP by a family member, partner,

<sup>3</sup>Of the 13 clients who were no longer involved in the treatment clinic at follow-up reasons for not being involved include: cost of treatment or insurance problems (n = 1), too many requirements from clinic or doctor to say on MAT (n = 1), trouble staying with the medication schedule (n = 1), didn't want to take medication for their drug problem (n = 6), no particular issues (n = 6), other reason (n = 1).

<sup>4</sup>See Appendix E for the complete comparison of clients who were still involved in the clinic at follow-up and clients who were not.

<sup>5</sup>For more information, see: Logan, T., Cole, J., Miller, J., & Scrivner, A. (2020). *Evidence Base for the Kentucky Opioid Program Treatment Outcome Study (KORTOS) Assessment and Methods*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

<sup>1</sup>Substance Abuse and Mental Health Services Administration (SAMHSA). (2015). *2013-2014 National Survey on Drug Use and Health: Model-based prevalence estimates (50 states and the District of Columbia)*. Rockville, MD: Substance Abuse and Mental Health Services Administration, Center for Mental Health Statistics and Quality.

<sup>2</sup>World Health Organization (2004). *Substitution maintenance therapy in the management of opioid dependence and HIV/AIDS prevention*. Geneva, Switzerland: United Nations Office on Drugs and Crime.

or friend. Less than half (47.9%) were unemployed, and of those unemployed clients, 38.0% reported they were looking for work.

In the six months before entering treatment, 98.4% of clients reported illegal drug use, 15.1% reported alcohol use, and 79.7% reported smoking tobacco. About one-fifth of clients (19.8%) reported using only opioids, 73.9% reported using opioids and at least one other class of drugs, and 4.7% of clients reported no opioids use (only other classes of drugs). In the past 30 days at intake, 98.4% of clients reported illegal drug use, 11.6% reported alcohol use, and 78.9% reported smoking tobacco. Clients were asked, at intake, how old they were when they first began to use illegal drugs, when they had their first alcoholic drink (more than just a sip), and when they began smoking cigarettes regularly. Trend outcomes show the age for having their first alcoholic drink at about 13 years old, first illegal drug use was 17 to 18 years old, and first tobacco use was 13 to 14 years old. Results of KORTOS drug trends show that although the majority of clients report illicit use of prescription opioids when they entered the program, the percent of clients who reported using heroin and methamphetamine in the 30 days before entering treatment has increased since CY 2011.

At intake, clients were asked 17 items about ten types of adverse childhood experiences from the Adverse Childhood Experiences (ACE). Results indicated that the majority of clients (76.0%) reported at least one adverse childhood experience. Significantly more men than women reported experiencing no adverse childhood experiences, whereas significantly more women than men reported experiencing 7 or more types of adverse childhood experiences. Significantly more women (32.9%) than (17.0%) men reported physical maltreatment. About 44% of women and 19.6% of men reported experiencing emotional neglect, which was also significantly different. Significantly more women than men also reported sexual abuse (36.7% vs. 8.0%) before the age of 18. In terms of interpersonal victimization experiences, compared to men, significantly more women reported having ever been the victim of assault (other than IPV), being stalked by someone who scared them, sexually assaulted or raped, abused by a dating or intimate partner, and verbally, sexually or otherwise harassed in a way that made the client afraid for their safety. About 29% percent of women and men reported having ever been directly or indirectly threatened with a gun or held at gunpoint.

In the six months before entering the program, 47.9% of clients met study criteria for depression, and 59.4% met study criteria for generalized anxiety. About 10% reported suicidal thoughts or attempts of suicide in the 6 months before entering the program. In addition, 16.1% had post-traumatic stress disorder (PTSD) scores that indicated risk of PTSD. About 46% of clients reported chronic pain in the 6 months before entering the program. Over half of clients (56.8%) reported they had at least one of the 15 chronic health problems listed on the intake interview. Trend analysis shows that from CY 2013 to CY 2019 the percent of clients who reported chronic medical problems has increased from just under half of clients to over half of clients.

## Change in Targeted Factors from Intake to Follow-up for Clients Still Involved in the Clinic at Follow-up

### Substance Use

When examining client change from past 6 months at intake to the 6-month follow-up period, clients who were still involved in the clinic at follow-up (n = 21) reported significant decreases in illicit drug use. Overall, 52.4% of clients reported illegal use of prescription opioids in the past 6 months at

intake, whereas 4.8% of clients reported illegal use of prescription opioids at any point during the 6 months before the follow-up assessment. Almost three-quarters of clients (71.4%) reported past-6-month heroin use at intake and that percent decreased to 38.1% at follow-up. Not only did clients' use of overall opioids decrease significantly, but also their use of non-opioid drugs (such as marijuana, tranquilizers, benzodiazepines, and stimulants) decreased from 85.7% to over one-third (38.1%). The majority of clients (90.0%) reported experiencing problems with drugs or alcohol (such as craving, withdrawal, wanting to quit and being unable, or worrying about relapse) at intake compared to 15.0% at follow-up. In addition, the number of clients who reported an ASI drug composite score that met the cut-off score for severe substance use disorder (SUD) decreased from 95.2% at intake to 19.0% at follow-up.

### Mental and Physical Health

There were also improvements in clients' overall past-6-month mental health. Two-thirds of clients met study criteria for depression at intake compared to 14.3% of clients at follow-up. Trend reports over the past 8 years indicate that, overall, the percent of clients who met criteria for depression at intake has fluctuated

between 59% and 75%. At follow-up, the percent of clients meeting study criteria for depression has been on the rise since 2017 before decreasing to 14.3% in 2022. Close to three-quarters of clients (71.4%) met study criteria for generalized anxiety at intake compared to 19.0% at follow-up. Similar to trends in depression, the percent of clients meeting study criteria for generalized anxiety has remained steady in the past 6 months at intake (around three-quarters) but has increased over the past 8 years to 31.7% in 2021 before decreasing to 19.0% in 2022. In addition, there was a decrease in clients who met study criteria for comorbid depression and generalized anxiety from intake (66.7%) to follow-up (9.5%). Further, 14.3% of clients reported suicidal ideation or attempts at intake compared to 0.0% at follow-up. Trends of suicidal ideation or attempts show that the percent of clients reporting suicide ideation in the past 6 months at intake appeared to peak in 2018 before decreasing again. At follow-up, the percent of clients reporting suicide ideation was stable at less than 5% over the past 8 years. The majority of clients at intake (81.0%) reported using alcohol, prescription drugs, or illegal drugs to reduce stress, anxiety, worry, sadness, or fear which decreased to 23.8% at follow-up.

**Overall, Kentucky opiate treatment program clients made significant strides in all of the targeted areas**



**REPORTED ANY ILLEGAL DRUG USE**

**100%** | **62%**  
at intake | at follow-up



**MET STUDY CRITERIA FOR COMORBID DEPRESSION AND ANXIETY\*\*\***

**67%** | **10%**  
at intake | at follow-up



**CURRENTLY EMPLOYED FULL-TIME**

**38%** | **43%**  
at intake | at follow-up



**REPORTED DIFFICULTY MEETING HEALTH CARE NEEDS**

**48%** | **29%**  
at intake | at follow-up



**AVERAGE NUMBER OF RECOVERY SUPPORT PERSONS\*\*\***

**3.8** | **8.0**  
at intake | at follow-up

Further, physical health was better for clients at follow-up. Specifically, clients reported significantly fewer number of days of poor physical (1.7 days compared to 5.4 days at intake) and mental health (4.3 days compared to 13.0 days at intake) in the past 30 days at follow-up. Significantly fewer clients reported they had experienced chronic pain in the 6 months before follow-up (47.6% at follow-up compared to 19.0% at intake). The majority of clients at intake (85.7%) reported using alcohol, prescription drugs, or illegal drugs to reduce their physical pain. At follow-up, 19.0% of clients reported using alcohol, prescription drugs, or illegal drugs to reduce their physical pain, which was a significant decrease of 66.7%.

Clients rated their quality of life as significantly higher after they began participating in the program. Trend analyses show that these high ratings for quality of life at follow-up have been consistent over the past 8 years.

### **Criminal Justice Involvement**

A minority of KORTOS clients reported criminal justice system involvement. In the 6 months before the intake, 4.8% of clients reported being arrested compared to 9.5% at follow-up. Further, 9.5% of clients reported being incarcerated in the 6 months before

treatment and 4.8% of the clients reported being incarcerated in the past 6 months at follow-up. Since 2015, a trend report shows that the percent of clients reporting an arrest had been stable with approximately 15-20% of clients reporting an arrest in the past 6 months at intake with a slight increase in 2021. Trend analyses show that the percent of clients who spent at least one night in jail were also consistent over the past 8 years at both intake and follow-up.

### **Economic Status and Living Circumstances**

KORTOS clients showed improvements in past-6-month economic and living circumstances from intake to follow-up. Specifically, there was a significant increase in the percent of clients reporting attending/completing vocational school, college, or graduate school from 42.9% at intake to 76.2% at follow-up. At intake and follow-up, 47.6% of clients reported having difficulty meeting basic living needs (e.g., food, shelter, utilities, and telephone) for financial reasons in the past 6 months. Trend reports over 8 years at intake indicate that the number of clients reporting difficulty meeting basic living needs has decreased overall at intake but has increased over the years at follow-up. The number of clients who reported they had difficulty obtaining health care

(e.g., doctor visits, dental visits, and prescription medications) for financial reasons decreased (but not significantly) from 47.6% at intake to 28.6% at follow-up. Overall, trends show that the percent of clients reporting difficulty meeting basic health care needs slowly decreased at intake from 48.9% in 2015 to 29.0% in 2019 but has increased back to 47.6% over the past few years. At follow-up, the percent of clients reporting difficulty meeting basic health care needs has increased from 4.6% in 2017 to 29.3% in 2021.

### **Recovery Supports**

At intake, 9.5% of clients reported going to mutual help recovery group meetings (e.g., AA, NA, or faith-based) in the past 30 days compared to 14.3% of clients at follow-up, which was not a significant increase. The average number of people clients said they could count on for recovery support increased significantly from 3.8 at intake to 8.0 follow-up. At intake and follow-up, clients were asked what, other than medication-assisted treatment, they believed would be most useful in helping them quit or stay off drugs/alcohol. The most common responses at intake were employment, the need to stay out of jail, and support from friends. At follow-up, the most common responses were the need to stay out of jail, employment, and support from friends.



## Multidimensional Recovery

Recovery goes beyond relapse or return to occasional drug or alcohol use. The multidimensional recovery measure items from the intake and follow-up surveys to create one measure of recovery. At intake, none of the clients had all positive dimensions of recovery, whereas at follow-up, almost half (47.6%) had all positive dimensions.

## Treatment Program Satisfaction

The majority of clients reported that the program started good (76.2%) and 95.2% reported it was currently going good. In addition, all clients reported that the treatment episode is working pretty well or extremely well for them. Furthermore, the majority of clients (95.2%) indicated they would refer a close friend or family member to their treatment provider. Of the clients who reported they would refer a close friend or family member to the program (n = 20), 15.0% reported they would warn their friend or family member about certain things or tell them who to work with or who to avoid.

On a scale from 1 representing the worst possible experience to 10 representing the best possible experience, clients rated their experience an 9.1 with 95.3% of clients giving a highly positive

rating of 8 through 10. The majority of clients reported that when they told their counselor or program staff personal things, they felt listened to and that the treatment approach and that the program staff believed in them and that treatment would work for them. In addition, clients felt that their expectations and hopes for treatment and recovery were met and that they felt they had input into their treatment goals, plans, and progress. Clients reported many positive aspects of their participation in the program including reduced substance use, improved mental health and their feelings about themselves, improved relationships with others, the quality of the treatment, and improved financial situation.

## Comparison of Clients Who Were No Longer Involved in the Treatment Clinic at Follow-up with Clients Who Were Still Involved

Of the 13 clients who were no longer involved in the treatment clinic at follow-up reasons for not being involved include: cost of treatment or insurance problems (n = 1), too many requirements from clinic or doctor to say on MAT (n = 1), trouble staying with the medication schedule (n = 1), didn't want to take medication for their drug problem (n = 6), no

particular issues (n = 6), and other reason (n = 1). There were a few significant demographic differences between clients who were still involved with the clinic at follow-up (n=21) and clients who were not (n=13). Clients who were involved with the clinic at follow-up were all White and clients who were no longer involved in the clinic at follow-up were more likely to be married. Clients who were not still involved in the treatment program reported significantly more years of education (15.6) compared to clients who were still involved in the clinic (13.9). Five percent of clients who were still involved in the clinic at follow-up and none of the clients who were no longer involved in the clinic reported less than a high school diploma or GED at follow-up. There were no differences between clients who were not involved in the clinic at follow-up and clients who were still involved at follow-up on employment in the past 30 days at follow-up.

Overall, significantly more clients who involved in the clinic at follow-up reported illicit drug use at follow-up, specifically, clients who were still involved in the clinic at follow-up reported stimulants or cocaine use in the 6 months before follow-up (19.0% vs. 0.0%). Significantly more clients who were still involved in the clinic at follow-up reported e-cigarette use in the past 6 months at

follow-up compared to clients who were no longer involved in the clinic. Close to 15% of clients who were not involved in the clinic at follow-up and 23.8% of clients who were still involved in the clinic at follow-up met or surpassed the Addiction Severity Index (ASI) composite score cutoff for alcohol and/or drug severe SUD at follow-up with no difference by group. There were significant differences between clients who were still involved in the clinic and clients who were not involved in the clinic at follow-up for the percent of clients attending mutual help recovery meetings in the past 30 days at follow-up (14.3% vs. 61.5%, respectively).

#### Areas of Concern

Several findings suggest opportunities to provide or target additional support for clients. First, 61.9% of KORTOS clients reported using illegal drugs in the 6 months before follow-up. While this year's follow-up sample size is small and may be affecting the results, this is still a sizable increase in illegal drug use at follow-up compared to last year (34.1%). Specifically, 38.1% of clients reported using heroin at follow-up and 38.1% of clients reported non-opioid drug use at follow-up. Additionally, 19.0% of clients still met criteria for 6 or more DSM-5 severity of substance use symptoms which classifies them as having a severe

substance use disorder. Rates tobacco smoking were high for clients (71.4%) and did not change from the past 6 months at intake to the past 6 months at follow-up.

While there were significant reductions in clients meeting study criteria for generalized anxiety, clients who met criteria for generalized anxiety at follow-up still reported an average of 6.5 (out of 7) symptoms in the past 6 months at follow-up. In addition, almost one-quarter of clients (23.8%) of clients reported using alcohol, prescription drugs, or illegal drugs to reduce stress, anxiety, worry, sadness, or fear in the past 6 months at follow-up. Further, 38.1% of clients were worried about their personal safety at follow-up. Also, while the number of clients reporting chronic pain decreased significantly from intake (47.6%) to follow-up (19.0%), clients still reported 26.3 average days of chronic pain at follow-up.

Further, almost half of clients (47.6%) at follow-up still reported having difficulty meeting basic living needs and 28.6% still reported difficulty meeting health care needs in the past 6 months. Similarly, almost half of clients (47.6%) remained unemployed at follow-up. Trends in economic difficulties show that the number of clients who

reported they had difficulty meeting basic living needs and/or health care needs has increased at follow-up since 2017.

There were only a few gender differences in targeted factors. Significantly more men reported heroin use in the past 30 days at follow-up and smokeless tobacco in the past 6 months at intake and in the past 30 days at intake.

It is important to keep in mind that this year, the follow-up sample size is small and should be considered when interpreting the findings. To increase the statistical power to detect change in this small sample size, the alpha for statistical tests was increased to  $p < .10$ , instead of  $p < .05$ .

The 2021 KORTOS evaluation indicates that opioid treatment programs in Kentucky have been successful in facilitating positive changes in clients' lives in a variety of ways, including decreased substance use, decreased mental health symptoms, decreased involvement with the criminal justice system, improved quality of life, improved health status, decreased economic hardship, and more support for recovery. This report also suggests there are a number of things clients continue to struggle with 6 months into their program participation.

## INTRODUCTION AND OVERVIEW

While prescription opioids are instrumental to reducing pain, misuse can lead to serious negative consequences such as addiction or even overdose. Non-medical use of prescription opioids is a continuing health concern in Kentucky where 6.5% of adults report nonmedical use of prescription opioids.<sup>6</sup> In 2019, Kentucky health care providers dispensed 72.3 prescriptions for opioids per 100 people compared to the United States which was 46.7 prescriptions per 100: the state with the fifth highest rate.<sup>7</sup> In addition, compared to the United States which had a rate of 14.6 opioid-involved overdose deaths, there were 23.4 overdose deaths involving opioids per 100,000 people in Kentucky.<sup>8</sup> Further, from September 2019 to September 2020, the number of Kentuckians who died from drug overdose increased 50% compared to the previous 12 months while there was a 28.8% increase nationwide.<sup>9</sup> Among the drugs found in toxicology reports in Kentucky drug overdose cases in 2020, the most frequent substance found was fentanyl found in 71% of all overdose cases.<sup>10</sup> Heroin-related emergency department visits and inpatient hospitalizations increased from the first quarter of 2020 through the second quarter of 2020 by 66.2% and 101.4% respectively.<sup>11</sup>

One of the key methods for treating persons addicted to opioids is through medication assisted therapy (or treatment, MAT) primarily with methadone or buprenorphine-naloxone (bup-nx). One of three priority areas of the United States Health and Human Services' (HHS) launched initiative in 2015 to reduce prescription opioid- and heroin-related overdose, death, and dependence is to expand the use of medication-assisted therapy.<sup>12</sup> These federally regulated opioid treatment programs (OTPs) provide evidence-based, clinically monitored, medication-assisted therapy with methadone or bup-nx.<sup>13</sup> Research evidence supports the effectiveness of methadone maintenance and bup-nx maintenance in retaining clients in treatment and reducing opioid use as well as reducing overdose deaths.<sup>14, 15</sup> The number of persons receiving methadone in substance use treatment in Kentucky rose from 2009 to

<sup>6</sup> [https://www.americashealthrankings.org/explore/annual/measure/drug\\_use/population/drug\\_use\\_presc\\_opioids/state/KY](https://www.americashealthrankings.org/explore/annual/measure/drug_use/population/drug_use_presc_opioids/state/KY)

<sup>7</sup> Center for Disease Control and Prevention. U.S. Opioid Prescribing Rate Maps. Retrieved from <https://www.cdc.gov/drugoverdose/maps/rxrate-maps.html> on April 19, 2021.

<sup>8</sup> NIDA. 2020, April 3. *Kentucky: Opioid-Involved Deaths and Related Harms*. Retrieved from <https://www.drugabuse.gov/drug-topics/opioids/opioid-summaries-by-state/kentucky-opioid-involved-deaths-related-harms> on April 19, 2021.

<sup>9</sup> Ahmad FB, Rossen LM, Sutton P. (2021). *Provisional drug overdose death counts*. National Center for Health Statistics. Retrieved from <https://www.cdc.gov/nchs/nvss/vsrr/drug-overdose-data.htm#data-tables> on April 19, 2021.

<sup>10</sup> Kentucky Office of Drug Control Policy. 2020 Overdose Fatality Report. <https://odcp.ky.gov/Documents/2020%20KY%20ODCP%20Fatality%20Report%20%28final%29.pdf>

<sup>11</sup> Kentucky Substance Use Research & Enforcement. (2021, January). Five major overdose-related substances in Kentucky, January 1, 2017-June 30, 2020. K-SURE Brief (No. 11). Retrieved on July 6, 2021 from [https://kiprc.uky.edu/sites/default/files/2021-02/K.SURE%20Product%2011%2C%202020\\_0.pdf](https://kiprc.uky.edu/sites/default/files/2021-02/K.SURE%20Product%2011%2C%202020_0.pdf).

<sup>12</sup> Office of the Assistant Secretary for Planning and Evaluation. (2015, March 26). Opioid abuse in the U.S. and HHS actions to address opioid-drug related overdoses and deaths. Washington, DC: U.S. Department of Health and Human Services, Office of the Assistant Secretary for Planning and Evaluation.

<sup>13</sup> Mattick, R., Breen, C., Kimber, J., & Davoli, M. (2009). Methadone maintenance therapy versus no opioid replacement therapy for opioid dependence. *Cochrane Database System Review*, Jul 8 (3).

<sup>14</sup> Kakko, J. Svanborg, K. D., Kreek, M J., & Heilig, M. (2003). 1-year retention and social function after buprenorphine-assisted relapse prevention treatment for heroin dependence in Sweden: A randomised, placebo-controlled trial. *Lancet*, 361, 662-668.

<sup>15</sup> Mattick, R., Kimber, J., Breen, C., & Davoli, M. (2008). Methadone maintenance therapy versus placebo or methadone maintenance for opioid dependence. *Cochrane Database of Systematic Reviews*, 2. CD002207.

2012, but decreased in 2013 while the number of persons receiving bup-nx multiplied by 5 from 2011 to 2013.<sup>16</sup>

In 2007, Kentucky OTPs began collecting state-specific outcome data on medication-assisted therapy. The outcome evaluation project is conducted in collaboration with the Kentucky Division of Behavioral Health, which is part of the Department of Behavioral Health, Developmental, and Intellectual Disabilities (DBHDID). The Kentucky Opioid Replacement Treatment Outcome Study (KORTOS) is conducted by the Behavioral Health Outcome Study team at the University of Kentucky Center on Drug and Alcohol Research (UK CDAR) and is an important part of the DBHDID Division of Behavioral Health's performance-based measurement of treatment outcomes in Kentucky's communities. The KORTOS project collects data from clients receiving medication-assisted treatment with methadone or bup-nx at licensed OTPs because they follow clinical monitoring protocols; thus, this report does not include data from all programs in Kentucky or independent physicians who prescribe bup-nx outside of an OTP. In calendar year 2020, six Kentucky licensed OTPs submitted data for KORTOS.<sup>17</sup>

In previous years, clients who were no longer involved in the clinic were not included in the eligible follow-up sample. However, beginning this report year, the decision was made to include these individuals in the eligible follow-up sample, but to analyze them separately. Therefore, for the follow-up analysis, only clients who reported that they were still involved with the clinic at follow-up (n = 21) were included; however, the 21 clients who were still involved in the clinic are compared to clients who completed a follow-up but were not still involved in the clinic at follow-up (n = 13) for each section.<sup>18,19</sup> This report describes outcomes for 21 adults who participated in a Kentucky OTP, completed an intake interview and then a follow-up telephone interview about 5-6 months (an average of 160 days) after the intake interview was completed.

Results are reported within nine main sections for the overall sample and separately by gender where there were significant differences.

**Section 1. Overview and Description of KORTOS Clients.** This section describes KORTOS including a description of clients who were involved in Kentucky's participating licensed OTPs in calendar year 2020 and who had completed an intake (n = 192) as well as clients who completed a 6-month follow-up interview and were still involved in the opiate treatment program (n = 21).

**Section 2. Substance Use.** This section examines change in substance use (any illegal drugs, alcohol, and tobacco) for 6-month and 30-day periods at intake and follow-up. Specific classes of illegal drugs examined include misuse of prescription opioids, non-prescribed methadone, non-prescribed bup-nx, heroin, and other illegal drugs. In addition, self-reported

<sup>16</sup> Substance Abuse and Mental Health Services Administration. (2015). *Mental health barometer: Kentucky, 2014*. HHS Publication No. SMA-15-4895KY. Rockville, MD: Substance Abuse and Mental Health Services Administration.

<sup>17</sup> In CY 2020, 6 of 12 OTPs submitted intake surveys for clients: Center for Behavioral Health –Bowling Green, Center for Behavioral Health –Elizabethtown, Center for Behavioral Health –Louisville, Pikeville Treatment Center, Ultimate Treatment Center, and Western Kentucky Medical.

<sup>18</sup> Of the 13 clients who were no longer involved in the treatment clinic at follow-up reasons for not being involved include: cost of treatment or insurance problems (n = 1), too many requirements from clinic or doctor to stay on MAT (n = 1), trouble staying with the medication schedule (n = 1), didn't want to take medication for their drug problem (n = 6), no particular issues (n = 6), other reason (n = 1).

<sup>19</sup> See Appendix E for the complete comparison of clients who were still involved in the clinic at follow-up and clients who were not.

severity of alcohol and drug use based on the DSM-5 criteria for severity of substance use disorder (SUD) and the Addiction Severity Index (ASI) alcohol and drug use composite scores are compared at intake and follow-up. Further, this section also examines change in problems experienced with alcohol/drug use, readiness for treatment, and medication-assisted treatment history.

**Section 3. Mental and Physical Health.** This section examines changes in mental health, physical health status, and quality of life from intake to follow-up. Specifically, this section examines: (1) depression, (2) generalized anxiety, (3) comorbid depression and generalized anxiety, (4) suicidal ideation and attempts, (5) posttraumatic stress disorder, (6) general health status, (7) perceptions of physical and mental health, (8) chronic pain, (9) health insurance, and (10) quality of life. The mental and physical health questions on the KORTOS intake and follow-up interviews were self-report measures.

**Section 4. Criminal Justice System Involvement and Interpersonal Victimization.** This section describes change in client involvement with the criminal justice system during the 6-month period before entering treatment and the 6-month period before the follow-up interview. Specifically, results include changes in: (1) any arrest, (2) the number of times arrested, among clients with any arrests, (3) any incarceration, (4) the number of nights incarcerated, among clients with any incarceration, (5) criminal justice supervision status, (6) interpersonal victimization, and (7) personal safety.

**Section 5. Education, Economic Status, and Living Circumstances.** This section examines changes in education, economic status, and living circumstances from intake to follow-up including: (1) highest level of education completed, (2) the number of months clients were employed full-time or part-time in the past 6 months, (3) current employment status, (4) hourly wage, (5) homelessness, (6) living situation, and (7) economic hardship (i.e., difficulty meeting living and health care needs for financial reasons).

**Section 6. Recovery Supports.** This section focuses on four main changes in recovery supports: (1) mutual help recovery group meeting attendance, (2) the number of people the client said they could count on for recovery support, (3) what will be most useful to the client in staying off drugs/alcohol, and (4) clients' perceptions of their chances of staying off drugs/alcohol.

**Section 7. Multidimensional Recovery.** This section examines multidimensional recovery that takes into account severity of substance use disorder, employment, homelessness, criminal justice system involvement, suicide ideation, overall health, recovery support, and quality of life. Change in recovery status from intake to follow-up is presented. Furthermore, a multivariate analysis was conducted to examine the intake indicators of recovery status and their association with having all eight dimensions of recovery at follow-up.

**Section 8. Client Satisfaction with the Opioid Treatment Programs.** The items measured in this report include: (1) client involvement in the program, (2) if the client would refer someone else to the program, (3) client ratings of program experiences, and (4) positive and negative aspects of program participation.

**Section 9. Conclusion and Implications.** This section summarizes the highlights from the evaluation results and suggests implications from these findings for the state.

It is important to keep in mind that this year, the follow-up sample size is small and should be considered when interpreting the findings. To increase the statistical power to detect change in this small sample size, the alpha for statistical tests was increased to  $p < .10$ , instead of  $p < .05$ .

## SECTION 1. KORTOS CLIENT CHARACTERISTICS

*This section briefly describes the Kentucky Opioid Replacement Treatment Outcome Study (KORTOS) including how clients are selected into the outcome evaluation. In addition, this section describes characteristics of clients who participated in federally licensed Kentucky opioid treatment programs in calendar year 2020 and who had completed an intake assessment (n = 192), including clients who also completed a 6-month follow-up interview (n = 34), of which 61.8% (n = 21) were still involved in a MAT program at follow up and 38.2% (n = 13) were not.*

KORTOS includes a face-to-face interview with program staff at the beginning of a new medication-assisted treatment episode. The interview is an evidence-based assessment<sup>20</sup> that asks about targeted factors such as substance use, mental health, involvement in the criminal justice system, quality of life, health status, and economic and living circumstances prior to entering treatment (submitted to UK CDAR from January 1, 2020 to December 31, 2020). In 2020, 192 adults completed an intake interview<sup>21</sup> that was submitted by one of 6 Kentucky licensed OTPs to UK CDAR.<sup>22</sup> The following section describes characteristics for all clients from those programs with a completed and submitted intake assessment.

### DESCRIPTION OF KORTOS CLIENTS AT TREATMENT INTAKE

#### DEMOGRAPHICS

Table 1.1 shows that over half of clients were male (58.3%) and most were White (91.7%). Clients were, on average, 39 years old, with the youngest client being 22 and the oldest being 70 years old. Overall, 56.8% were married or cohabiting, 22.9% of clients had never been married, 17.2% were separated or divorced, and 3.1% were widowed. Over 40% of clients reported they had at least one child under the age of 18 who was living with them in the 6 months before they entered the program. The majority of clients (88.4%) indicated they lived in a metropolitan community, 10.4% lived in a nonmetropolitan community, and 1.2% were from a very rural community.

<sup>20</sup>Logan, TK, Cole, J., Miller, J., Scrivner, A., & Walker, R. (2020). *Evidence Base for the Kentucky Opioid Replacement Treatment Outcome Study (KORTOS) Assessment and Methods*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

<sup>21</sup>When a client had more than one intake survey in the same fiscal year, the survey with the earliest submission date was kept in the data file and the other intake surveys were deleted so that each client was represented once and only once in the data set.

<sup>22</sup>In CY 2020, 6 of 12 OTPs submitted intake surveys for clients: Center for Behavioral Health –Bowling Green, Center for Behavioral Health –Elizabethtown, Center for Behavioral Health –Louisville, Pikeville Treatment Center, Ultimate Treatment Center, and Western Kentucky Medical.

TABLE 1.1. DEMOGRAPHICS FOR ALL KORTOS CLIENTS AT INTAKE (N = 192)

<b>Age<sup>23</sup></b> .....	39.1 years ( <i>Min. = 22, Max. = 70</i> )
<b>Gender</b>	
Male .....	58.3%
Female.....	41.1%
Transgender .....	0.5%
<b>Race</b>	
White/Caucasian.....	91.7%
Black/African American.....	5.7%
Other or multiracial .....	2.6%
<b>Marital status</b>	
Never married.....	22.9%
Separated or divorced .....	17.2%
Married or cohabiting.....	56.8%
Widowed .....	3.1%
<b>Have children under the age of 18 who live with them..</b>	43.8%
<b>Type of community<sup>24</sup></b>	
Metropolitan.....	88.4%
Nonmetropolitan	10.4%
Very rural	1.2%

## EDUCATION


About 17% of clients had less than a high school diploma or GED at intake (see Figure 1.1). About 43% of the sample had a high school diploma or GED and 25.0% of clients had completed some vocational/technical school or college. Only a minority of clients had completed vocational/technical school (5.2%), an associate's degree (4.2%), or a bachelor's degree or higher (6.3%).

<sup>23</sup>Two clients had incorrect birthdates and, therefore, age could not be determined.

<sup>24</sup>19 clients either did not indicate a county of residence or lived in another state.




FIGURE 1.1. HIGHEST LEVEL OF EDUCATION COMPLETED AT INTAKE (N = 192)



### SELF-REPORTED REFERRAL SOURCE

Figure 1.2 shows the self-reported treatment referral source for all KORTOS clients. Less than two-thirds of clients (60.9%) decided to get help on their own and 30.2% of clients reported they were referred by a family member, partner or friend. A small percentage of clients (3.6%) were transferred or referred by another OTP, 2.6% were referred by a health care or mental health care worker, 0.5% were referred by the justice system, 1.5% were referred by other sources, and 0.5% were referred by a substance abuse treatment facility.

FIGURE 1.2 SELF-REPORTED REFERRAL SOURCE FOR ALL KORTOS CLIENTS AT INTAKE (N = 192)



### EMPLOYMENT

Over one-third of clients (36.5%) reported they had not worked in the past 6 months, 15.6% had worked 1 to 3 months, and 47.9% had worked 4 or more months (not depicted in figure). In the 30 days before entering the program, almost half of clients (47.9%) reported being unemployed, 40.1% reported they were employed full-time, and 12.0% were employed part-time or had occasional or seasonal employment (see Figure 1.3). Among those who reported being employed full or part-time at intake (n = 100), the median hourly wage was \$15.88.

FIGURE 1.3. EMPLOYMENT STATUS IN THE PAST 30 DAYS AT INTAKE (N = 192)





Figure 1.4 shows that of the individuals who were currently unemployed at intake (n = 92), 38.0% stated they were looking for work, 27.2% were on disability, 10.9% were unemployed and not looking for work, 9.8% were keeping the house or taking care of children full-time at home, 4.3% were on furlough from their job or temporarily laid off, 1.1% were students, and the remaining 8.7% gave other reasons for not being employed (e.g., retired, or in a controlled environment).


FIGURE 1.4. OF THOSE UNEMPLOYED, REASONS FOR BEING UNEMPLOYED (N = 92)



## ADVERSE CHILDHOOD EXPERIENCES AND VICTIMIZATION

At intake, clients were asked 17 items about ten types of adverse childhood experiences from the Adverse Childhood Experiences (ACE).<sup>25, 26, 27</sup> In addition to providing the percent of clients who reported each of the ten types of adverse childhood experiences before the age of 18 years old captured in ACE, the number of types of experiences was computed such that items clients answered affirmatively were added to create a score equivalent to the ACE score. A score of 0 means the client answered “No” to the five abuse and neglect items and the five household dysfunction items in the intake interview. A score of 10 means the client reported all five forms of child maltreatment and neglect, and all five types of household dysfunction before the age of 18. Figure 1.5 shows that 24.0% reported they did not experience any of the ACE included in the assessment. Almost 41% of clients reported experiencing 1 to 3 ACE, 21.4% reported experiencing 4 – 6 ACE, and 12.5% reported experiencing 7 – 9 ACE. Only 1.6% of clients reported experiencing all 10 types of adverse childhood experiences.

FIGURE 1.5. NUMBER OF TYPES OF ADVERSE CHILDHOOD EXPERIENCES REPORTED AT BASELINE (N = 192)



There was a significant difference in the proportion of men and women<sup>28</sup> classified by number of types of ACE (see Figure 1.6). Women had a higher average number of ACE compared of men. In addition, more women than men reported experiencing 7 or more ACE.


<sup>25</sup> Felitti, V. J., Anda, R. F., Nordenberg, D., Williamson, D. F., Spitz, A. M., Edwards, V., Koss, M. P., & Marks, J. S. (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine*, 14(4), 245-258.

<sup>26</sup> Centers for Disease Control and Prevention. (2014). *Prevalence of individual adverse childhood experiences*. Atlanta, GA: National Center for Injury Prevention and Control, Division of Violence Prevention. <http://www.cdc.gov/violenceprevention/acestudy/prevalence.html>.

<sup>27</sup> The intake assessment asked about 10 major categories of adverse childhood experiences: (a) three types of abuse (e.g., emotional maltreatment, physical maltreatment, and sexual abuse), (b) two types of neglect (e.g., emotional neglect, physical neglect), and (c) five types of family risks (e.g., witnessing partner violence victimization of parent, household member who was an alcoholic or drug user, a household member who was incarcerated, a household member who was diagnosed with a mental disorder or had committed suicide, and parents who were divorced/separated).

<sup>28</sup> Because there was only one transgender person, they were not included in the analysis for confidentiality reasons.


FIGURE 1.6. THE NUMBER OF CATEGORIES OF ADVERSE CHILDHOOD EXPERIENCES BY GENDER\*\*



\*\*p < .01, \*\*\*p < .001.

There were several significant differences between men and women for maltreatment and neglect experiences in childhood. Significantly more women (44.3%) reported they had experienced emotional maltreatment in their childhood, compared to 25.9% of men (see Figure 1.7). Significantly more women (32.9%) than (17.0%) men reported physical maltreatment. About 44% of women and 19.6% of men reported experiencing emotional neglect, which was also significantly different. Significantly more women than men also reported sexual abuse (36.7% vs. 8.0%) before the age of 18.


FIGURE 1.7. MALTREATMENT AND NEGLECT EXPERIENCES IN CHILDHOOD BY GENDER (n = 191)



\*p < .05, \*\*p < .01, \*\*\*p < .001.

Over half of men and women (63.3% and 51.8%, respectively) reported their parents were divorced or lived separately (see Figure 1.8). Over one-quarter of women and 15.2% of men reported they had witnessed the intimate partner violence of their mother or stepmother. Significantly more women (50.6%) than men (34.8%) reported they lived with someone who was a problem drinker or alcoholic or used street drugs. Significantly more women (40.5%) than men (25.9%) also reported they had a household member with a mental illness or who had committed suicide. About 14% of women and 12.5% of men reported a household member had been incarcerated.

FIGURE 1.8. HOUSEHOLD RISKS IN CHILDHOOD BY GENDER (n = 192)




\*p < .05.

## VICTIMIZATION EXPERIENCES

Clients were also asked about victimization experiences (including when they may have been the victim of a crime, harmed by someone else, or felt unsafe) they had experienced in their lifetime and in the 6 months before entering the treatment program. The results of the most commonly reported experiences are presented by gender in Figure 1.9. Compared to men, significantly more women reported having ever been the victim of assault (other than IPV), being stalked by someone who scared them, sexually assaulted or raped, abused by a dating or intimate partner, and verbally, sexually, or otherwise harassed in a way that made them afraid for their safety. Close to 30% of women and men reported having ever been assaulted and directly or indirectly threatened with a gun or held at gunpoint.


FIGURE 1.9. LIFETIME CRIME AND INTERPERSONAL VICTIMIZATION BY GENDER (n = 191)



\*p < .05, \*\*\*p < .001.

Smaller percentages of clients reported experiencing crime and interpersonal victimization in the 6 months before entering programs than in their lifetime (see Figure 1.10). Significantly more women than men reported having been sexually assaulted or raped, and abused by a dating or intimate partner.

FIGURE 1.10. PAST-6-MONTH CRIME AND INTERPERSONAL VICTIMIZATION BY GENDER (n = 191)




\*p < .05, \*\*p < .01.

## MENTAL HEALTH

At intake, 47.9% of KORTOS clients met study criteria for depression in the past 6 months (see Figure 1.11). Additionally, 59.4% of clients met study criteria for generalized anxiety at intake. One in ten clients reported suicidal thoughts or attempts in the 6 months before entering the program and 16.1% of clients had PTSD scores that indicated a risk of PTSD.<sup>29</sup>

<sup>29</sup>Price, M., Szafranski, D. D., van Stolk-Cooke, K., & Gros, D. F. (2016). Investigation of abbreviated 4 and 8 item versions of the PTSD Checklist 5. *Psychiatry Research*, 239, 124-130.

FIGURE 1.11. DEPRESSION, GENERALIZED ANXIETY, SUICIDALITY, AND POST TRAUMATIC STRESS DISORDER IN THE PAST 6 MONTHS AT INTAKE (N = 192)



PHYSICAL HEALTH

At intake, clients reported an average of 7.5 days of poor physical health in the past 30 days and an average of 11.2 days of poor mental health in the past 30 days (see Table 1.2). About 46% of clients reported chronic pain in the 6 months before entering the program. Over half of clients (56.8%) reported they had at least one of the 16 chronic health problems listed on the intake interview. Of those clients (n = 109), the most common medical problems were hepatitis C, arthritis, cardiovascular disease, and asthma.


The most common insurance provider reported at intake was Medicaid (55.2%; see Table 1.2). Nineteen percent of clients did not have any insurance. Small percentages of clients had insurance through an employer, including through a spouse, partner, or self-employment (16.0%) and Medicare (9.9%).

TABLE 1.2. HEALTH-RELATED CONCERNS FOR ALL KORTOS CLIENTS AT INTAKE (N = 192)

Average number of poor health days in past 30 days .....	7.5
Average number of poor mental health days in past 30 days .....	11.2
Average number of days poor physical or mental health limited activities.....	8.6
Chronic pain .....	46.4%
<b>At least one chronic medical problem.....</b>	<b>56.8%</b>
Hepatitis C .....	39.4%
Arthritis .....	27.5%
Cardiovascular/heart disease .....	22.9%
Asthma .....	19.3%
<b>Medical insurance</b>	
No insurance .....	18.8%
Medicaid .....	55.2%
Through employer (including spouse’s employer, parents’ employer, and self-employed) .....	16.0%
Medicare .....	9.9%
VA/Champus/Tricare .....	0.0%
Through health exchange .....	0.0%

Figure 1.12 shows the percent of clients who reported having different numbers of chronic medical problems at intake. Over two-fifths (43.2%) of clients reported no problems, and less than one-third (31.3%) reported one chronic medical problem. About 26% reported two chronic medical problems and none reported having three or more chronic medical problems.


FIGURE 1.12. NUMBER OF CHRONIC MEDICAL PROBLEMS AT INTAKE FOR TOTAL SAMPLE (N = 192)



**TREND ALERT: CHRONIC MEDICAL PROBLEMS AT INTAKE**

At intake, clients were asked if, in their lifetime, they have been told by a doctor they had any of the chronic medical problems listed (e.g., diabetes, arthritis, asthma, heart disease, cancer, hepatitis B or C, cirrhosis of the liver). The percent of clients reporting at least one chronic health problem in their lifetime remained steady from CY 2013 (48.5%) to CY 2016 (51.9%) and increased to 65.4% in CY 2018. In CY 2020, 56.8% of clients reporting at least one chronic health problem.

FIGURE 1.13. TRENDS IN CHRONIC MEDICAL PROBLEMS AT INTAKE






**TREND ALERT: CHRONIC PAIN AT INTAKE**

The percent of clients who reported chronic pain at intake has increased slowly, but minimally, over time. In CY 2013 37.4% of clients reported experiencing chronic pain and in CY 2020 46.4% reported experiencing chronic pain.

FIGURE 1.14. TRENDS IN CHRONIC PAIN AT INTAKE




CY 2013 (N = 703)	CY 2014 (N = 717)	CY 2015 (N = 517)	CY 2016 (N = 514)	CY 2017 (N = 625)	CY 2018 (N = 384)	CY 2019 (N = 233)	CY 2020 (N = 192)
-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------

**SUBSTANCE USE**

The majority of KORTOS clients who completed an intake interview reported using illegal drugs (98.4%) and smoking tobacco (79.7%) while 15.1% reported using alcohol in the 6 months before intake (see Figure 1.15). The drug classes reported by the greatest number of clients were heroin (72.4%), prescription opioids/opiates<sup>30</sup> (62.5%), marijuana (52.6%), and amphetamines (42.7%; not represented in a figure).

Similarly, 98.4% reported using illegal drugs, 78.9% reported smoking tobacco, and 11.6% reported using alcohol in the 30 days before entering treatment.

FIGURE 1.15 ALCOHOL, DRUG, AND TOBACCO USE 6 MONTHS AND 30 DAYS BEFORE TREATMENT<sup>31</sup>



<sup>30</sup>For brevity's sake, the class of substances including prescription opiates and opioids will be referred to as opioids.

<sup>31</sup>Because being in a controlled environment reduces opportunities for substance use, only clients who were not incarcerated for the entire time period were included in the substance use analysis; therefore, none of the clients were excluded from the past-6-month substance use but 2 clients were excluded from the past-30-day use.

Figure 1.16 presents the percentage distribution of individuals who used alcohol and/or illegal drugs in the 6 months before entering the program. Among the individuals who were not incarcerated all 180 days before entering the program, 1.6% reported no alcohol or drug use, none reported alcohol use only, 83.3% reported illegal drug use only, and 15.1% reported both alcohol and illegal drug use. None of the clients reported only alcohol use.

FIGURE 1.16. PAST-6-MONTH ALCOHOL AND ILLEGAL DRUG USE AT INTAKE FOR THOSE NOT INCARCERATED ALL 180 DAYS BEFORE ENTERING THE PROGRAM (N = 192)





Figure 1.17 presents the distribution of clients who reported using no drugs, alcohol only, only opioids (including prescription opioids, bup-nx, methadone, heroin) and other drug classes from the following: marijuana, CNS depressants (such as benzodiazepines, sedatives, tranquilizers, barbiturates), stimulants (including amphetamines and cocaine), and other classes such as hallucinogens, synthetic marijuana, and inhalants. KORTOS clients who were not incarcerated all 180 days before entering the program are predominately polysubstance users. Close to 20% of clients reported only using opioids (19.8%) while 73.9% reported using opioids and at least one other class of drug.

FIGURE 1.17. PAST-6-MONTH POLYSUBSTANCE USE AT INTAKE FOR THOSE NOT INCARCERATED ALL 180 DAYS BEFORE ENTERING THE PROGRAM (N = 192)




TREND REPORT IN SPECIFIC PAST-30-DAY DRUG USE

When looking at trends over time for all clients with completed intake interviews, the percent of clients using prescription opioids in the past 30 days peaked in CY 2008 and steadily dropped until CY 2018 when it increased to 65.6% before decreasing again in CY 2019. The percent of clients who reported using non-prescribed methadone before entering treatment showed a decline from CY 2007 to 2011 and again from CY 2012 to 2019. The percent of clients who reported using bup-nx slowly increased from CY 2007 through 2015, dropped slightly in CY 2016 and 2017 before increasing again in CY 2018. In CY 2020, the percent of clients who reported using bup-nx decreased further to 14.7%.

The most notable change in substance use among KORTOS clients, however, is for heroin. Small percentages of KORTOS clients reported using heroin from CY 2007 through 2011. Then, the percent tripled from 8.8% in CY 2011 to 26.3% and then nearly doubled from 26.3% in CY 2012 to 48.1% in CY 2013. The percent of KORTOS clients reporting heroin use at intake in CY 2014 increased again to 50.7% and further still to 56.5% in CY 2015. The number of KORTOS clients reporting heroin use at intake in CY 2017 increased further to 69.6%. While heroin use decreased in CY 2018 and again in CY 2019, there was an increase to the highest point of 71.6% in CY 2020.

In addition, the use of methamphetamine among clients has gradually been increasing since CY 2008. In CY 2018 and CY 2019, about one-quarter of clients were using methamphetamine when they entered the program, which was in increase from CY 2017. In CY 2020 there was a sharp increase in methamphetamine use to 40.7%. These trends are very similar when examining only those clients who were followed-up (see Appendix D).<sup>32</sup>

FIGURE 1.18. PERCENT OF ALL CLIENTS WITH A COMPLETED INTAKE INTERVIEW REPORTING NON-PRESCRIBED USE OF PRESCRIPTION OPIOIDS, METHADONE, BUP-NX, HEROIN, AND METHAMPHETAMINE IN THE 30 DAYS BEFORE ENTERING TREATMENT AT THE OTP (N = 9,456)<sup>33, 34</sup>




<sup>32</sup>Due to the proximity of the trend lines, only the most recent year’s data is labeled.  
<sup>33</sup>Clients who reported being in a controlled environment all 30 days before entering treatment were not included in this analysis.  
<sup>34</sup>Two clients in CY 2015 had missing data for past-30-day heroin use at intake.

### TRENDS IN AGE OF FIRST USE


Clients were asked, at intake, how old they were when they first began to use illegal drugs, when they had their first alcoholic drink (more than just a sip), and when they began smoking cigarettes regularly (see Figure 1.19). The age at which KORTOS clients reported initiating drug use was steady for the past 8 years, around 17 to 18 years old. Clients generally reported having their first alcoholic drink in their early teens (around 13 years old). The age of first regular smoking tobacco use was steady for the past 8 years, typically between 13 and 14 years old.

FIGURE 1.19. AGE OF FIRST USE REPORTED AT INTAKE



At intake, clients were asked how important it was to them to help others who have had substance use problems. The majority of clients (60.9%) reported they agreed or strongly agreed that it is important while 5.7% disagreed or strongly disagreed.


FIGURE 1.20. CLIENTS REPORTING THE IMPORTANCE OF HELPING OTHERS WHO HAVE A SUBSTANCE USE PROBLEM (N = 192)



### CRIMINAL JUSTICE INVOLVEMENT

Less than 13% of clients reported being arrested at least once and 12.5% reported being incarcerated at least one night in the 6 months before entering treatment (see Figure 1.21). Among those who were arrested (n = 24), they were arrested an average of 2.0 times. Among those who were incarcerated (n = 24), they were incarcerated an average of 32.8 nights (not depicted in the figure). In addition, 9.4% of clients reported being under supervision by the criminal justice system.

FIGURE 1.21. CRIMINAL JUSTICE INVOLVEMENT 6 MONTHS BEFORE TREATMENT AT INTAKE  
(N = 192)



## KORTOS FOLLOW-UP SAMPLE

Follow-up interviews are conducted with a selected sample of KORTOS clients targeted for 6 months after the intake interview is completed. At the completion of the intake interview, program staff inform individuals about the KORTOS follow-up study and ask if they are interested in participating. Clients who agree to participate are asked to provide contact information. All individuals who agree to be contacted by UK CDAR for the follow-up interview and have given at least one mailing address and one phone number, or two phone numbers if they do not have a mailing address in their locator information, are eligible for the follow-up component of the study. All eligible individuals are then selected by the month in which they completed intake interviews.<sup>35</sup>

Of the 192 clients who completed an intake interview, 73 agreed to be contacted for the follow-up interview (38.0% agreement rate). Of these clients, a total of 61 provided the minimum amount of contact information at the end of the intake interview, had intake interviews that were submitted to UK CDAR within 30 days of completion, and were selected into the follow-up sample. Of the 61 clients included in the follow-up sample, none were ineligible for participating in the follow-up interview. Of these 61 eligible clients, UK CDAR interviewers completed follow-up assessments with 34 clients (55.7% follow-up rate). This means that roughly 44% of eligible individuals included in the sample to be followed up were not successfully contacted within the targeted eligibility time period.<sup>36</sup>

Follow-up procedures for the outcome study use several best practices. First, the follow-up assessments are conducted independently from the treatment programs by UK CDAR staff. Second, UK CDAR has over 20 years of extensive experience following up study participants and staff are highly trained, supervised, and monitored. Third, the confidentiality of clients is protected through specific study procedures, UK human subjects' protections, and through a federal certificate of confidentiality. Clients are provided with full information about their rights as a research subject and the protections for confidentiality provided by the study. Clients must consent to the study twice: once at the completion of the intake interview and once when on the phone for the follow-up interview.

In previous years, clients who were no longer involved in the clinic were not included in the eligible follow-up sample. However, beginning in the 2021 report year, the decision was made to include these individuals in the eligible follow-up sample, but to analyze them separately. Therefore, for the follow-up analysis, only clients who reported that they were still involved with the clinic at follow-up ( $n = 21$ ) were included; however, the 21 clients who were still involved in the clinic are compared to clients who completed a follow-up, but were

<sup>35</sup> If a person has more than one intake interview in a given year, the interview with the earliest date will be selected into the follow-up sample.

<sup>36</sup> Clients are not contacted for a variety of reasons including follow-up staff are not able to find a working address or phone number or are unable to contact any friends or family members of the client.

not still involved in the clinic at follow-up (n = 13) for each section.<sup>37, 38</sup> This report describes outcomes for 21 adults who were still participating in a Kentucky OTP, completed an intake interview and then a follow-up telephone interview about 5-6 months (an average of 188 days) after the intake interview was completed. Detailed information about the methods and follow-up efforts can be found in Appendices A and B.

Of the 21 adults who completed a 6-month follow-up interview, 66.7% were female. All follow-up clients were White and an average of 37.6 years old and between the ages of 23 and 60. Around 29% of clients had never been married at intake, 19.0% were separated or divorced, 47.6% were married or cohabiting, and 4.8% were widowed. About 38% of follow-up clients had at least one child under age 18 who was living with them and all clients indicated they lived in a metropolitan community.

TABLE 1.3. DEMOGRAPHICS FOR KORTOS FOLLOW-UP CLIENTS AT INTAKE (N = 21)

	Clients still in the clinic at follow-up (n = 21)	Clients not in the clinic at follow-up (n = 13)
Age .....	37.6 years ( <i>range of 23 - 60</i> )	43.4 years ( <i>range 24 - 70</i> )
<b>Gender</b>		
Male .....	33.3%	53.8%
Female.....	66.7%	46.2%
<b>Race</b>		
White/Caucasian.....	100.0%	84.6%
Black/African American.....	0.0%	15.4%
Other race or multiracial .....	0.0%	0.0%
<b>Marital status</b>		
Never married.....	28.6%	23.1%
Separated or divorced .....	19.0%	53.8%
Married or cohabiting.....	47.6%	7.7%
Widowed .....	4.8%	0.0%
Have children under the age of 18 who live with them .....	38.1%	38.5%
<b>Type of community<sup>39</sup></b>		
Metro.....	100.0%	91.7%
Non-metro .....	0.0%	8.3%
Very rural	0.0%	0.0%

<sup>37</sup> Of the 13 clients who were no longer involved in the treatment clinic at follow-up reasons for not being involved include: cost of treatment or insurance problems (n = 1), too many requirements from clinic or doctor to stay on MAT (n = 1), trouble staying with the medication schedule (n = 1), didn't want to take medication for their drug problem (n = 6), no particular issues (n = 6, other reason (n = 1).

<sup>38</sup> See Appendix E for the complete comparison of clients who were still involved in the clinic at follow-up and clients who were not.

<sup>39</sup> One client was missing data for type of community at intake.

When clients who completed a follow-up interview were compared with those who did not have a follow-up interview on a variety of intake variables, there were only two significant differences for demographics, socio-economic status, substance use, physical health, and mental health. More clients who completed a follow-up were female and, among clients who reported having been arrested in the past 6 months at intake, clients who completed a follow-up reported more arrests.<sup>40</sup>

TABLE 1.4. FOLLOWED-UP VERSUS NOT FOLLOWED-UP

	Followed up	
	No (n = 158)	Yes (n = 34)
Demographic	More clients were male	More clients were female
Socio-economic status indicators (e.g., education, employment, living situation, inability to meet basic needs)	No differences	
Substance use, severity of alcohol and drug use	No differences	
Treatment history	No differences	
Health (e.g., overall health status, chronic medical problems, chronic pain)	No differences	
Mental health (e.g., depression, generalized anxiety, suicidality)	No differences	
Criminal justice involvement (e.g., arrested, incarcerated)	Reported more arrests among clients who reported an arrest	

<sup>40</sup> See Appendix C for detailed comparisons of clients who completed a follow-up interview and were included in the follow-up analysis (n = 34) and clients who did not complete a follow-up interview or were not included in the follow-up analysis (n = 158).

## SECTION 2. SUBSTANCE USE

*This section describes change in illegal drug, alcohol, and tobacco use from intake to follow-up (n = 41). Past-6-month substance use is examined as well as past-30-day substance use for clients who were not in a controlled environment all 30 days before entering treatment or the follow-up interview. In addition, this section includes problems experienced with substance use in the past 30 days, readiness for treatment, self-reported severity of alcohol and drug use, and medication-assisted treatment. Results for each targeted factor are presented for the overall sample and by gender when there were significant gender differences.*

Changes in illegal drug, alcohol, and tobacco use before entering the program and during the 6-month follow-up period are presented in this section. In addition to examining the overall use of illegal drugs, several specific categories of illegal drugs were examined including: (a) prescription opioid misuse (including opioids such as morphine, Percocet, Oxycontin, Lortab), (b) non-prescribed methadone, (c) non-prescribed buprenorphine-naloxone (bup-nx), (d) heroin, and (e) non-opioid drugs other than those mentioned above (including marijuana, cocaine, amphetamines, tranquilizers, hallucinogens, inhalants, and barbiturates). Analysis is presented in detail for KORTOS study participants who were not in a controlled environment for the entire period of 6 months and/or 30 days before entering treatment. Changes in substance use from intake to follow-up are presented in 4 main subsections and organized by type of substance use:

1. **Change in past-6-month substance use from intake to follow-up.** Comparison of any illegal drugs, prescription opioid misuse, non-prescribed methadone, non-prescribed bup-nx, heroin, other non-opioid drugs, alcohol, and tobacco use in the 6 months before the client entered the program and use of these substances during the 6-month follow-up period (n = 21) are presented.
2. **Average number of months clients used substances at intake and follow-up.** For those who used any illegal drugs, alcohol, or tobacco, the average number of months of use before program entry and during the follow-up period are reported.
3. **Change in 30-day substance use from intake to follow-up.** Comparison of any illegal drugs, prescription opioid misuse, non-prescribed methadone, non-prescribed bup-nx, heroin, other non-opioid drugs, alcohol, and tobacco use in the 30 days before the client entered the program and during the follow-up period (n = 21) is presented.<sup>41</sup> In addition, this section examines the number of days clients experienced alcohol/drug problems in the past 30 days, how troubled or bothered clients were by alcohol/drug problems in the past 30 days, and how important treatment is for these alcohol/drug problems at intake and follow-up.
4. **Change in self-reported severity of alcohol and drug use from intake to follow-up.** There are two indices of substance use severity presented in this report. One way to examine overall change in degree of severity of substance use is to ask participants to self-report whether they met the 11 criteria included in the DSM-5 for diagnosing substance use disorder in the past 6 months. Under DSM-5, anyone meeting any two of the 11 criteria during the same 6-month period would receive a diagnosis of substance use disorder (SUD) as long as their symptoms were causing clinically

<sup>41</sup>None of the individuals were in a controlled environment all 30 days before intake or before follow-up; therefore, all clients were included in the analysis.



significant impairments in functioning. The severity of the substance use disorder (i.e., none, mild, moderate, or severe) in this report is based on the number of criteria met. The percent of individuals in each of the four categories at intake and follow-up is presented.


The Addiction Severity Index (ASI) composite scores are examined for change over time for illegal drugs (n = 21). The ASI composite score assesses self-reported addiction severity even among those reporting no substance use in the past 30 days. The drug composite score is computed from items about 30-day drug use and the number of days individuals used multiple drugs in a day, as well as the impact of substance use on the individual's life, such as number of days individuals had drug problems, how troubled or bothered individuals were by their drug problems, and how important treatment was to them.

## ALCOHOL AND/OR DRUG USE

### PAST-6-MONTH ALCOHOL AND/OR DRUG USE

All clients reported using alcohol and/or illegal drugs in the 6 months before entering the program, which decreased to 66.7% at follow-up (see Figure 2.1).

FIGURE 2.1. PAST 6-MONTH ALCOHOL AND/OR DRUG USE AT INTAKE AND FOLLOW-UP (N = 21)



a – No measures of association could be computed for alcohol and/or illegal drug use because the value at intake was 0.


*“I am a chronic relapser and I have tried other programs, This is my fifth time in treatment and nothing else has ever worked. It has been 100% no using and no relapse since I have started this program.”*

KORTOS FOLLOW-UP CLIENTS

### TRENDS IN ANY ALCOHOL AND/OR DRUG USE

The number of KORTOS clients reporting alcohol and/or drug use in the 6 months before treatment was consistently high (about 97%) until 2021 when it briefly decreased to 85.4%. At follow-up, from 2015 – 2018, less than half of clients reported any alcohol and/or drug use. In 2019, 53.1% of clients reported alcohol and/or drug use compared to 39.0% in 2021. In 2022, the number of clients reporting any alcohol and/or drug use at follow-up increased to 66.7%.


FIGURE 2.2. TRENDS IN ANY ALCOHOL AND/OR ILLEGAL DRUG USE AT INTAKE AND FOLLOW-UP, REPORTS 2015-2022<sup>42, 43</sup>



### PAST-30-DAY ALCOHOL AND/OR DRUG USE

All clients reported using alcohol and/or illegal drugs in the 30 days before entering the program, which decreased to 42.9% at follow-up (see Figure 2.3).

FIGURE 2.3. PAST 30-DAY ALCOHOL AND/OR DRUG USE AT INTAKE AND FOLLOW-UP (N = 21)



a - No measures of association could be computed for alcohol and/or illegal drug use because the value at intake was 0.

<sup>42</sup>For each trend report presented, the years correspond to years in which the annual reports were published. In addition, all trend analyses present only annual report data at intake and follow-up and do not include between-year statistical analysis.


<sup>43</sup>In 2015, 3 cases had missing data for drug use at intake.

## ANY ILLEGAL DRUGS

### PAST-6-MONTH ANY ILLEGAL DRUG USE

All clients reported using illegal drugs in the 6 months before entering the program, which decreased to 61.9% at follow-up (see Figure 2.4).

FIGURE 2.4. PAST-6-MONTH ILLEGAL DRUG USE AT INTAKE AND FOLLOW-UP (N = 21)




a - No measures of association could be computed for illegal drug use because the value at intake was 0.

### AVERAGE NUMBER OF MONTHS USED ANY ILLEGAL DRUGS

Clients who reported any illegal drug use at intake (n = 21) reported an average maximum of 5.6 months of use. Among clients who reported any illegal drug use in the 6 months before follow-up (n = 13), the maximum number of months they reported using any drug was, on average, 4.5 months (see Figure 2.5).


FIGURE 2.5. AVERAGE NUMBER OF MONTHS CLIENTS USED ILLEGAL DRUGS



### TRENDS IN PAST-6-MONTH ILLEGAL DRUG USE

The percent of KORTOS clients reporting any illegal drug use in the 6 months before treatment was consistently high (about 97%) until 2021 when it decreased to 85.4%. The percent of clients who reported any illegal drug use at follow-up increased from 31.5% in 2015 to 49.0% in 2019. In 2021, the percent of clients reporting any illegal drug use at follow-up decreased to 34.1%, but increased to 61.9% report year 2022, which corresponds to intake in 2020, the beginning of the pandemic.


FIGURE 2.6. TRENDS IN ILLEGAL DRUG USE AT INTAKE AND FOLLOW-UP, REPORTS 2015-2022<sup>44</sup>



### PAST-30-DAY ANY ILLEGAL DRUG USE

At intake, all clients reported any illegal drug use in the 30 days before entering the program and at follow-up, 42.9% of clients reported any illegal drug use in the past 30 days.

FIGURE 2.7. PAST-30-DAY USE OF ANY ILLEGAL DRUGS AT INTAKE AND FOLLOW-UP (N = 21)



a - No measures of association could be computed for illegal drug use because the value at intake was 0.


<sup>44</sup>The small sample size could be affecting the increase in illegal drug use at follow-up.

## PRESCRIPTION OPIOID MISUSE

### PAST-6-MONTH PRESCRIPTION OPIOID MISUSE

Over half of clients (52.4%) reported misusing prescription opioids (such as morphine, Percocet, Oxycontin, Lortab) in the 6 months before treatment entry. At follow-up, 4.8% of clients reported misusing prescription opioids (see Figure 2.8). This means there was a 47.6% significant decrease in the percent of clients reporting prescription opioid misuse.


FIGURE 2.8. PAST-6-MONTH PRESCRIPTION OPIOID MISUSE AT INTAKE AND FOLLOW-UP (N = 21)



### AVERAGE NUMBER OF MONTHS MISUSED PRESCRIPTION OPIOIDS

Figure 2.9 shows the average number of months prescription opioid users reported misusing prescription opioids at intake and during the 6-month follow-up. Among the clients who reported misusing prescription opioids before entering the program (n = 11), clients reported using prescription opioids an average of 4.6 of the 6 months. Only one client reported misusing opioids at follow-up and used 4.0 of the 6 months before follow-up.

FIGURE 2.9. AVERAGE NUMBER OF MONTHS CLIENTS USED PRESCRIPTION OPIOIDS




### PAST-30-DAY PRESCRIPTION OPIOID MISUSE

At intake, 52.4% of clients reported past-30-day misuse of prescription opioids and at follow-up, 4.8% of clients reported misuse of prescription opioids (see Figure 2.10). This reflects a significant decrease of 47.6% in the percent of clients reporting misuse of

prescription opioids in the past 30 days.

FIGURE 2.10. PAST-30-DAY PRESCRIPTION OPIOID MISUSE AT INTAKE AND FOLLOW-UP (N = 21)




\*\*\*p < .01.

## NON-PRESCRIBED USE OF METHADONE

### PAST-6-MONTH NON-PRESCRIBED USE OF METHADONE

Almost 10% of clients reported using non-prescribed methadone in the 6 months before intake (see Figure 2.11). At follow-up, none of the clients reported non-prescribed use of methadone. Clients who reported non-prescribed methadone use at intake reporting using an average of one month in the past 6 months.

FIGURE 2.11. PAST-6-MONTH NON-PRESCRIBED METHADONE USE AT INTAKE AND FOLLOW-UP (N = 21)




a - No measures of association could be computed for methadone use because the value at follow-up was 0.

### PAST-30-DAY NON-PRESCRIBED USE OF METHADONE

About 10% of clients reported using non-prescribed methadone in the 30 days before entering the program (see Figure 2.12). At follow-up, none of the clients reported past-30-day use of non-prescribed methadone.

FIGURE 2.12. PAST-30-DAY NON-PRESCRIBED METHADONE USE AT INTAKE AND FOLLOW-UP (N = 21)




a - No measures of association could be computed for methadone use because the value at follow-up was 0.

## NON-PRESCRIBED USE OF BUP-NX

### PAST-6-MONTH NON-PRESCRIBED USE OF BUP-NX

Figure 2.13 shows that 14.3% of clients reported using non-prescribed bup-nx in the 6 months before intake. At follow-up, none of the clients reported using non-prescribed bup-nx. Clients who reported using bup-nx reported they reported using non-prescribed bup-nx an average of 2.7 of the 6 months.

FIGURE 2.13. PAST-6-MONTH NON-PRESCRIBED USE OF BUP-NX AT INTAKE AND FOLLOW-UP (N = 21)




a - No measures of association could be computed for non-prescribed buprenorphine because the value at follow-up was 0.

### PAST-30-DAY NON-PRESCRIBED USE OF BUP-NX

About 14% of clients reported using non-prescribed bup-nx in the 30 days before entering the program (see Figure 2.14). At follow-up, none of the clients reported past-30-day use of non-prescribed bup-nx.

FIGURE 2.14. PAST-30-DAY NON-PRESCRIBED BUP-NX USE AT INTAKE AND FOLLOW-UP (N = 21)




a - No measures of association could be computed for non-prescribed buprenorphine because the value at follow-up was 0.

## HEROIN

### PAST-6-MONTH HEROIN USE

Almost three-quarters of clients (71.4%) reported using heroin in the 6 months before entering treatment, which significantly decreased 33.3% to 38.1% at follow-up (see Figure 2.15).

FIGURE 2.15. PAST-6-MONTH HEROIN USE AT INTAKE AND FOLLOW-UP (N = 21)




\*\*p < .05.

### AVERAGE NUMBER OF MONTHS USED HEROIN

Among the clients who reported using heroin in the 6 months before entering treatment (n = 15), they reported using heroin, on average, 4.8 months (see Figure 2.16). Among clients who reported using heroin in the 6 months before follow-up (n = 8), they reported using, on average, 4.4 months.

FIGURE 2.16. AVERAGE NUMBER OF MONTHS CLIENTS USED HEROIN




### PAST-30-DAY HEROIN USE

Almost three-quarters of clients (71.4%) reported using heroin in the 30 days before intake. At follow-up, 19.0% reported using heroin in the past 30 days, a significant decrease of 52.4% (see Figure 2.17).



FIGURE 2.17. PAST-30-DAY HEROIN USE AT INTAKE AND FOLLOW-UP (N = 21)




\*\*\*p < .01.

**GENDER DIFFERENCES IN PAST-30-DAY HEROIN USE**

Significantly more men than women using heroin in the past 30 days at follow-up, 42.9% vs. 7.1%. The percent of women who reported heroin use significantly decreased from intake to follow-up (see Figure 2.18).

FIGURE 2.18. GENDER DIFFERENCES IN PAST-30-DAY HEROIN USE AT INTAKE AND FOLLOW-UP<sup>a</sup>




a—Significant difference by gender at follow-up; p < .05.  
\*\*\*p < .01.

**NON-OPIOID DRUG USE**

**PAST-6-MONTH USE OF NON-OPIOID DRUGS**

The majority of clients (85.7%) used illegal drugs other than prescription opioids, non-prescribed methadone, non-prescribed bup-nx, or heroin in the 6 months before entering the program (see Figure 2.19). Drugs in this category include marijuana, cocaine, amphetamines, tranquilizers/benzodiazepines/sedatives, hallucinogens, inhalants, barbiturates, and synthetic drugs like synthetic marijuana or bath salts. The percent of clients who reported use of non-opioid drugs decreased to 38.1% at follow-up (a significant decrease of 47.6%).

FIGURE 2.19. PAST-6-MONTH NON-OPIOID DRUG USE AT INTAKE AND FOLLOW-UP (N = 21)




\*\*\*p < .01.

### AVERAGE NUMBER OF MONTHS USED NON-OPIOID DRUGS

Figure 2.20 shows the maximum number of months clients that used non-opioid drugs reported using these illegal drugs (e.g., marijuana, cocaine, amphetamine, tranquilizers, barbiturates, inhalants, hallucinogens, synthetic drugs).<sup>45</sup> Among the clients who reported using non-opioid drugs at intake (n = 18), the maximum number of months clients reported using any of these drugs was an average of 4.6 months. Among clients who reported using non-opioid drugs at follow-up (n = 8), the maximum average number of months clients reported using any of these drugs was 3.8 months.

FIGURE 2.20. AVERAGE MAXIMUM NUMBER OF MONTHS CLIENTS USED NON-OPIOID DRUGS




### PAST-30-DAY USE OF NON-OPIOID DRUGS

About 86% of clients reported using non-opioid drugs in the 30 days before intake (see Figure 2.21). At follow-up, 28.6% of clients reported non-opioid drug use, which is a 57.1% significant decrease.

<sup>45</sup> Because number of months of use of each class of substance was measured separately (e.g., marijuana, cocaine, amphetamines, tranquilizers, barbiturates, inhalants, hallucinogens, synthetic drugs), the value is a calculation of the maximum number of months clients used any substance class.

FIGURE 2.21. PAST-30-DAY NON-OPIOID DRUG USE AT INTAKE AND FOLLOW-UP (N = 21)



### INJECTION DRUG USE

At intake, 52.4% of clients reported having ever injected any drug in their lifetime. Of those clients (n = 11), 45.5% reported having ever used a Needle Exchange Program in Kentucky and all reported they were in Jefferson County/Louisville. At follow-up, 14.3% of clients reported injecting drugs in the past 6 months. Of those clients (n = 3), one reported having used a Needle Exchange Program in Kentucky, which was in Jefferson County.


### ALCOHOL USE

There were three measures of alcohol use including: (1) any alcohol use, (2) alcohol use to intoxication, and (3) binge drinking. Binge drinking was defined as having 5 or more (4 or more if the client was female) alcoholic drinks in a period of about 2 hours.<sup>46</sup>

#### PAST-6-MONTH ALCOHOL USE

Nineteen percent of clients reported using alcohol in the 6 months before entering treatment while 14.3% of clients reported alcohol use in the 6 months before follow-up (see Figure 2.22). Very few clients reported using alcohol to intoxication and binge drinking in the past 6 months at intake (9.5% and 4.8%, respectively). None of the clients reported alcohol use to intoxication or binge drinking at follow-up.

FIGURE 2.22. PAST-6-MONTH ALCOHOL USE AT INTAKE AND FOLLOW-UP (N = 21)




<sup>46</sup>National Institute on Alcohol Abuse and Alcoholism. (2004, Winter). NIAAA council approves definition of binge drinking. *NIAAA Newsletter, Winter 2004* (3). Rockville, MD: Department of Health and Human Services, National Institutes of Health, national Institute on Alcohol Abuse and Alcoholism.

### AVERAGE NUMBER OF MONTHS USED ALCOHOL

Figure 2.23 shows the average number of months alcohol users reported using alcohol at intake and follow-up. Among the clients who reported using alcohol in the 6 months before entering treatment (n = 4), they reported using alcohol, on average, 4.0 months. Among clients who reported using alcohol in the 6 months before follow-up (n = 3), they reported using an average number of 5.3 months.


FIGURE 2.23. AVERAGE NUMBER OF MONTHS OF ALCOHOL USE



### TRENDS IN PAST-6-MONTH ALCOHOL USE

Less than one-third of clients reported any alcohol use in the 6 months before entering treatment. The percent of clients reporting alcohol use decreased to 22.9% in 2017, but increased to 27.5% at intake in 2018. In 2019, the percent of clients reporting alcohol use decreased to 23.3% and continued to decrease in 2021 to 4.9%. In 2022, however, the number of clients reporting alcohol use at intake increased to 19.0%. At follow-up, the percent of clients who reported alcohol use increased to 19.1% in 2016, but decreased to 12.6% by 2018. The percent of clients reporting alcohol use at follow-up remained relatively stable from 2015 to 2019. In 2020, the percent of clients reporting past-6-month alcohol use at intake or at follow-up decreased compared to 2019 and increased slightly to 12.2% in 2021 and again in 2022 to 14.3%.

FIGURE 2.24. TRENDS IN ALCOHOL USE AT INTAKE AND FOLLOW-UP, REPORTS 2015-2022<sup>47</sup>




### PAST-30-DAY ALCOHOL USE

About 10% of clients reported using alcohol use in the 30 days before intake and follow-up (see Figure 2.25). Close to 5% of clients reported they either drink alcohol to intoxication and none of the clients reported alcohol to intoxication at follow-up. None of the clients reported binge drinking in the past 6 months at intake or follow-up.

<sup>47</sup>In 2015, 5 cases had missing data for alcohol use at intake.


FIGURE 2.25. PAST-30-DAY ALCOHOL USE AT INTAKE AND FOLLOW-UP (N = 21)



## PROBLEMS EXPERIENCED WITH SUBSTANCE USE IN THE PAST 30 DAYS

In the past 30 days at intake, 90.0% of clients reported they experienced problems with drugs or alcohol such as craving, withdrawal, wanting to quit but being unable, or worrying about relapse (see Figure 2.26). In the past 30 days at follow-up, 15.0% of clients reported experiencing problems with drugs or alcohol (a significant decrease of 75.0%).

FIGURE 2.26 CLIENTS EXPERIENCING PROBLEMS WITH ILLEGAL DRUGS OR ALCOHOL AT INTAKE AND FOLLOW-UP (N = 21)




\*\*\*p < .01.

## READINESS FOR SUBSTANCE ABUSE TREATMENT

Figure 2.27 shows that 85.7% of clients reported they were considerably or extremely troubled or bothered by drug or alcohol problems in the past 30 days at intake. In the past 30 days at follow-up, 4.8% of clients reported that they were considerably or extremely troubled or bothered by drug or alcohol problems (a significant decrease of 80.9%).

The figure below also shows that 95.2% of clients in the past 30 days at intake and 81.0% of clients in the past 30 days at follow-up reported that treatment for drug or alcohol problems was considerably or extremely important.

FIGURE 2.27. READINESS FOR TREATMENT FOR ILLEGAL DRUG OR ALCOHOL USE AT INTAKE AND FOLLOW-UP (N = 21)



\*\*\*p &lt; .01.

## SUBSTANCES THAT CLIENT HAD THE MOST TROUBLE WITH

Of the clients who reported any drug or alcohol use in the past 6 months at intake (n = 21), the majority of clients (66.7%) reported that heroin was the most problematic substance for them followed by prescription opiate misuse (28.6%). The second most problematic substance reported was marijuana (23.8%) followed by stimulants (14.3%).

Of the clients who reported any drug or alcohol use in the past 6 months at follow-up (n = 14), 50.0% reported heroin was the most problematic and 21.4% reported they had no preference. Half of clients reported there was no other substance that was the second most problematic and 14.3% reported stimulants.

## SELF-REPORTED SEVERITY OF ALCOHOL AND DRUG USE

### DSM-5 CRITERIA FOR SUBSTANCE USE DISORDER, PAST 6 MONTHS

One way to examine overall change in degree of severity of substance use is to ask participants to self-report whether they met the 11 criteria included in the DSM-5 for diagnosing substance use disorder (SUD) in the past 6 months. The DSM-5 diagnostic criteria for substance use disorders included in the KORTOS intake and follow-up interviews are similar to the criteria for DSM-IV, which has evidence of excellent test-retest reliability and validity.<sup>48,49</sup> However, the DSM-5 eliminates the distinction between substance abuse and dependence, substituting severity ranking instead. In addition, the DSM-5 no longer includes the criterion about legal problems arising from substance use but adds a new criterion about craving and compulsion to use.<sup>50</sup> Under DSM-5, anyone meeting any two

<sup>48</sup>Hasin, D., & Paykin, A. (1999). Alcohol dependence and abuse diagnoses: Concurrent validity in a nationally representative sample. *Alcoholism: Clinical and Experimental Research*, 23(1), 144-150.


<sup>49</sup>Hasin, D., Trautman, K., Miele, G., Samet, S., Smith, M., & Endicott, J. (1996). Psychiatric Research Interview for Substance and Mental Disorders (PRISM): Reliability for substance abusers. *American Journal of Psychiatry*, 153(9), 1195-1201.

<sup>50</sup>Malone, M., & Hoffmann, N. (2016). A comparison of DSM-IV versus DSM-5 substance use disorder diagnoses in adolescent populations. *Journal of Child & Adolescent Substance Abuse*, 25(5), 399-408.

of the 11 criteria during the same 6-month period for either alcohol or drugs would receive a diagnosis of substance use disorder as long as their symptoms were causing clinically significant impairments in functioning. The severity of the substance use disorder (i.e., none, mild, moderate, or severe) in this report is based on the number of criteria met. Clients who report 2 or 3 DSM-5 symptoms are considered to have a mild substance use disorder, 4 or 5 symptoms is considered a moderate substance use disorder, and 6 or more symptoms is considered severe.

Change in the severity of SUD in the prior 6 months was examined for clients at intake and follow-up. Figure 2.28 displays the change in the percent of individuals in each SUD severity classification, based on self-reported criteria in the preceding 6 months. At intake, none of the clients met criteria for no substance use disorder (meaning they reported 0 or 1 DSM-5 criteria for SUD), while at follow-up, 71.4% of clients met criteria for no SUD, a significant increase of 71.4%. At the other extreme of the continuum, the vast majority of clients (95.2%) met criteria for severe SUD at intake, while at follow-up, only 19.0% met criteria for severe SUD, a significant decrease of 76.2%.

FIGURE 2.28. DSM-5 SUD SEVERITY AT INTAKE AND FOLLOW-UP (N = 21)<sup>a</sup>



a – Significance tested with the Stuart-Maxwell Test for Marginal Homogeneity ( $p < .001$ ).  
 \*\*\* $p < .01$ .

*“They’re really good and take everything into consideration. It feels good to talk to someone who doesn’t look down on you.”*

KORTOS FOLLOW-UP CLIENTS

## ADDICTION SEVERITY INDEX CRITERIA FOR SUBSTANCE USE DISORDER, PAST 30 DAYS


Another way to examine overall change in degree of severity of substance use is to calculate the Addiction Severity Index (ASI) composite scores for alcohol and drug use. These composite scores are computed based on self-reported severity of past 30-day alcohol and drug use, taking into consideration several issues including:

- The number of days of alcohol (or drug) use,
- Money spent on alcohol,
- The number of days individuals used multiple drugs (for drug use composite score),
- The number of days individuals experienced problems related to their alcohol (or drug) use,
- How troubled or bothered they are by their alcohol (or drug) use, and
- How important treatment is to them for their alcohol (or drug) problems (see sidebar).

Change in the average ASI composite score for drug use was examined for clients who were not in a controlled environment all 30 days before entering treatment. Clients who reported abstaining from drugs at both intake and follow-up were not included in the analysis of change in drug composite score.

Figure 2.29 displays the change in past 30-day average composite scores from intake to follow-up. The average for the drug composite score decreased significantly from 0.34 to 0.14.

FIGURE 2.29. AVERAGE ASI DRUG COMPOSITE SCORES OF THOSE WHO USED SUBSTANCES AT INTAKE AND/OR FOLLOW-UP<sup>51</sup>



\*\*\*p < .01.

<sup>51</sup> Because so few clients reported only alcohol use (n = 2), the alcohol composite score was skewed. Therefore, it is not presented in this year's report.

## ASI Alcohol and Drug Composite Scores and Substance Dependence

Rikoon et al. (2006) conducted two studies to determine the relationship between the ASI alcohol and drug use composite scores and DSM-IV substance dependence diagnosis. They identified alcohol and drug use composite score cutoffs that had 85% sensitivity and 80% specificity with regard to identifying DSM-IV substance dependence diagnosis: .17 for alcohol composite score and .16 for drug composite score. These composite score cutoffs can be used to estimate the number of individuals who are likely to meet criteria for active alcohol or drug dependence and to show reductions in self-reported severity of substance use. In previous years we have used the ASI composite scores to estimate the number and percentage of clients who met a threshold for alcohol and drug dependence. However, recent changes in the diagnostics for substance abuse call into question the distinction between dependence and abuse. Thus, ASI composite scores that met the threshold can be considered indicative of severe substance use disorder to be compatible with current thinking about substance use disorders in the DSM-V (American Psychiatric Association, 2013), where we would have previously referred to them as meeting the threshold for dependence. Change from intake to follow-up in the severity rating has the same clinical relevance as moving from dependence to abuse in the older criteria.


Rikoon, S., Cacciola, J., Carise, D., Alterman, A., McLellan, A. (2006). Predicting DSM-IV dependence diagnoses from Addiction Severity Index composite scores. *Journal of Substance Abuse Treatment*, 31(1), 17-24.

American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders (5th ed.)*. Arlington, VA: American Psychiatric Publishing.



ASI drug composite scores that met the cutoff for severe substance use disorder (SUD) significantly decreased from 95.2% at intake to 19.0% at follow-up (see Figure 2.30).


FIGURE 2.30. OF THOSE WHO USED SUBSTANCES, INDIVIDUALS WITH ASI COMPOSITE SCORES MEETING THE CUTOFF FOR SEVERE SUBSTANCE USE DISORDER IN THE PAST 30 DAYS AT INTAKE AND FOLLOW-UP



\*\*\*p < .01.

Analyses were also conducted to determine if clients who had a drug composite score indicative of severe SUD at intake and follow-up differed by gender or age (see Figure 2.31).<sup>52</sup> There were no significant gender, or age group differences from intake to follow-up for clients who had a drug composite score indicative of severe SUD.

FIGURE 2.31. DRUG-USING CLIENTS WITH A DRUG COMPOSITE SCORE INDICATIVE OF SEVERE SUD IN THE PAST 30 DAYS AT INTAKE AND FOLLOW-UP BY DEMOGRAPHIC FACTORS (N = 21)



\*\*\*p < .01.

<sup>52</sup> All follow-up clients were White and, therefore, drug composite scores were not compared by race.

## SUBSTANCE ABUSE TREATMENT HISTORY

### LIFETIME SUBSTANCE ABUSE TREATMENT

Prior to the current admission, 61.9% of clients reported at intake that they had received services for substance abuse (including detox, drug court, and recovery programs). Overall, clients reported receiving services or substance abuse an average of 2.1 times in their lifetime.

### OVERDOSE HISTORY


Over one-third of clients (38.1%,  $n = 8$ ) reported that they have overdosed on drugs (and required interventions by someone to recover) in their lifetime (an average of 1.8 times, among the clients who reported they had overdosed on drugs). In the past 6 months at intake, 9.6% of clients ( $n = 2$ ) reported having an overdose (an average of 1.5 time among the clients who reported having an overdose in the past 6 months at intake). At follow-up, one of the clients reported having an overdose in the past 6 months (2 times in the past 6 months).

### MEDICATION-ASSISTED TREATMENT

#### MEDICATION-ASSISTED TREATMENT AT INTAKE


Two-thirds of clients who were not incarcerated all 180 days before treatment, reported at intake they had ever received medication from a clinic or doctor's office to help with their substance abuse. Of these clients ( $n = 14$ ), 64.3% reported it was dispensed in a clinic, 28.6% were prescribed the medication by a doctor in a specialty clinic, and 7.1% were prescribed the medication by a doctor in a general medical practice (Figure 2.32). They also reported using the prescribed medication for 1.9 months in the 6 months before they began involvement at the OTP and 3.9 days in the past 30 days at intake.

FIGURE 2.32. CLIENTS REPORTING WHERE MEDICATION WAS RECEIVED (N=14)



The majority of clients (92.9%) reported that they think their use of medication-assisted treatment helped treat their drug problem, 7.1% reported it had no effect on their drug problem, and none reported medication-assisted treatment made their drug problems worse (Figure 2.33).

FIGURE 2.33. CLIENTS REPORTING HOW MUCH THEY BELIEVE THE MEDICATION HELPED THEIR OPIOID USE PROBLEMS (N=14)



### *MEDICATION-ASSISTED TREATMENT AT FOLLOW-UP*

The majority of clients (95.2%) who were not incarcerated all 180 days before treatment entry or in the past 6 months at follow-up, reported that they received methadone in the past 6 months at follow-up. About 5% of clients reported receiving Suboxone, and none of the clients received Vivitrol.


At follow-up, clients reported using the medication prescribed to them for an average of 5.8 months in the past 6 months. In addition, clients reported using the medication prescribed for an average of 26.3 days in the past 30 days. In the past 6 months at follow-up, the majority of clients (85.7%) reported they had not participated in any medication-assisted treatment program other than the one they've been most recently involved with. Overall, at follow-up, all clients reported that they think their use of medication-assisted treatment helped treat their drug problem.

## TOBACCO USE

### PAST-6-MONTH SMOKING, E-CIGARETTES, AND SMOKELESS TOBACCO USE

There was no significant change in either smoking, e-cigarettes, or smokeless tobacco use from intake to follow-up (see Figure 2.34). Most clients reported smoking tobacco in the 6 months before entering the program (71.4%) and in the 6 months before follow-up (71.4%). About 24% of clients reported the use of e-cigarettes (e.g., battery-powered nicotine delivery devices that vaporize a liquid mixture consisting of propylene glycol, glycerin, flavorings, nicotine, and other chemicals) at intake and follow-up. Around 10% of clients at intake and 4.8% of the clients at follow-up reported using smokeless tobacco in the past 6 months.


FIGURE 2.34. PAST-6-MONTH SMOKING TOBACCO, E-CIGARETTE, AND SMOKELESS TOBACCO USE AT INTAKE AND FOLLOW-UP (n = 21)



*GENDER DIFFERENCES IN PAST-6-MONTH SMOKELESS TOBACCO USE*

Significantly more men than women reported smokeless tobacco use in the past 6 months at intake (28.6% vs. 0.0%) and follow-up (14.3% vs. 0.0%; see Figure 2.35).

FIGURE 2.35. GENDER DIFFERENCES IN PAST-30-DAY SMOKELESS TOBACCO USE AT INTAKE AND FOLLOW-UP<sup>a</sup>




a—Significant difference by gender at intake; p < .05.

*AVERAGE NUMBER OF MONTHS OF SMOKING TOBACCO*

Figure 2.36 shows that among clients who reported smoking tobacco in the 6 months before entering treatment (n = 15), they reported smoking tobacco, on average, 5.5 months. In the 6 months before follow-up, there was no change in the average number of months clients reported smoking tobacco among clients who reported smoking tobacco (6.0 months; n = 15).


FIGURE 2.36. AVERAGE NUMBER OF MONTHS OF SMOKING TOBACCO USE



### TRENDS IN PAST-6-MONTH SMOKING TOBACCO USE

The majority of KORTOS clients at intake and follow-up reported smoking tobacco. The only significant change in the use of smoking tobacco from intake to follow-up was in report year 2018 when 83.5% of clients reported smoking tobacco at intake and 76.9% of clients reported smoking tobacco at follow-up.


FIGURE 2.37. TRENDS IN SMOKING TOBACCO USE AT INTAKE AND FOLLOW-UP, REPORTS 2015-2022<sup>53</sup>



### TRENDS IN PAST-6-MONTH E-CIGARETTE USE

Each year slightly more clients have reported e-cigarette use at follow-up compared to intake. The percent of clients reporting e-cigarette use at follow-up increased from 15.0% in the 2019 report to 29.8% in the 2020 report. In 2021, however, the percent of clients reporting e-cigarette use decreased at both intake and follow-up compared to 2020.

FIGURE 2.38 TRENDS IN E-CIGARETTE USE AT INTAKE AND FOLLOW-UP, REPORTS 2018-2022<sup>54</sup>




<sup>53</sup>In 2015, 5 cases had missing data for smoking tobacco use at intake.

<sup>54</sup>E-cigarette use, specifically, was not included in the instrument until the 2018 report.

### AVERAGE NUMBER OF CIGARETTES SMOKED

The average number of cigarettes clients reported smoking daily decreased slightly from intake to follow-up (see Figure 2.39). Of those who smoked tobacco at intake (n = 15), clients reported smoking an average of 23.0 cigarettes per day. At follow-up, among clients who reported smoking tobacco (n = 15), they reported smoking an average of 16.6 cigarettes per day.


FIGURE 2.39. NUMBER OF CIGARETTES SMOKED IN AN AVERAGE DAY AMONG CLIENTS WHO SMOKED TOBACCO



### PAST-30-DAY USE SMOKING, E-CIGARETTE, AND SMOKELESS TOBACCO USE


The percent of clients who reported any smoking or smokeless tobacco use, or e-cigarette use in the past 30 days did not change significantly from intake to follow-up (see Figure 2.40).

FIGURE 2.40. PAST-30-DAY SMOKING, E-CIGARETTE AND SMOKELESS TOBACCO USE AT INTAKE AND FOLLOW-UP (n = 21)



### GENDER DIFFERENCES IN PAST-30-DAY SMOKELESS TOBACCO USE

Significantly more men than women smokeless tobacco use in the past 30 days at intake (28.6% vs. 0.0%) and follow-up (14.3% vs. 0.0%; see Figure 2.41).

FIGURE 2.41. GENDER DIFFERENCES IN PAST-30-DAY SMOKELESS TOBACCO USE AT INTAKE AND FOLLOW-UP<sup>a</sup>

a—Significant difference by gender at intake;  $p < .05$ .

### SUBSTANCE USE COMPARISONS BETWEEN CLIENTS WHO WERE STILL INVOLVED IN THE TREATMENT PROGRAM AT FOLLOW-UP AND CLIENTS WHO WERE NOT STILL INVOLVED

Use of illegal drugs in past the 6 months at follow-up was compared for clients who were still involved in the treatment program at follow-up ( $n = 21$ ) and clients who were not ( $n = 13$ ). Overall, significantly more clients who were involved in the clinic at follow-up reported illicit drug use at follow-up, specifically, clients who were still involved in the clinic at follow-up reported stimulants or cocaine use in the 6 months before follow-up compared to clients who were not involved in the clinic at follow-up (19.0% vs. 0.0%). There were no significant differences between clients who were involved with the clinic at follow-up and clients who were not for alcohol use at follow-up.

In the past 6 months at follow-up, the majority of clients reported smoking tobacco products, with no difference between clients who were involved in the clinic at follow-up and those who were not. Significantly more clients who were still involved in the clinic at follow-up reported e-cigarette use in the past 6 months at follow-up compared to clients who were no longer involved in the clinic.

Close to 15% of clients who were not involved in the clinic at follow-up and 23.8% of clients who were still involved in the clinic at follow-up met or surpassed the Addiction Severity Index (ASI) composite score cutoff for alcohol and/or drug severe SUD at follow-up with no difference by group. The average score for the drug severity composite score was 0.04 for clients who were not involved in the clinic at follow-up and 0.13 for clients who were still involved in the clinic, which was not a significant difference.

## SECTION 3: MENTAL AND PHYSICAL HEALTH

*This section examines changes in mental health, physical health status, and quality of life from intake to follow-up. Specifically, this section examines: (1) depression, (2) generalized anxiety, (3) comorbid depression and generalized anxiety, (4) suicidal ideation and attempts, (5) posttraumatic stress disorder, (6) general health status, (7) perceptions of physical and mental health, (8) chronic pain, (9) health insurance, and (10) quality of life. The mental and physical health questions on the KORTOS intake and follow-up interviews were self-report measures.*

### DEPRESSION SYMPTOMS

To assess depression, participants were first asked two screening questions:

“Did you have a two-week period when you were consistently depressed or down, most of the day, nearly every day?” and

“Did you have a two-week period when you were much less interested in most things or much less able to enjoy the things you used to enjoy most of the time?”

If participants answered “yes” to at least one of these two screening questions, they were then asked seven additional questions about symptoms of depression (e.g., sleep problems, weight loss or gain, feelings of hopelessness or worthlessness).


Two-thirds of clients met study criteria for depression in the 6 months before they entered treatment (see Figure 3.1). At follow-up, 14.3% met study criteria for depression—a significant decrease of 52.4%.

Of those who met study criteria at intake (n = 14), they had an average of 7.3 symptoms out of 9. At follow-up, among those who met study criteria for depression (n = 3), clients reported an average of 6.3 symptoms out of 9.

#### Study Criteria for Depression

To meet study criteria for depression, clients had to say “yes” to at least one of the two screening questions and at least 4 of the 7 symptoms. Thus, the minimum score to meet study criteria: 5 out of 9.

FIGURE 3.1. MEETING STUDY CRITERIA FOR PAST-6-MONTH DEPRESSION AT INTAKE AND FOLLOW-UP (N = 21)




\*\*\*p < .01.



## TRENDS IN PAST-6-MONTH DEPRESSION

The percent of clients who met criteria for depression at intake has fluctuated over the past 8 years between 59% and 75%. The percent of clients who met criteria for depression at follow-up has been on the rise since 2017 (10.9%) to 29.3% in 2021 before decreasing to 14.3% in 2022.

FIGURE 3.2. TRENDS IN THE NUMBER OF CLIENTS WHO MET STUDY CRITERIA FOR PAST-6-MONTH DEPRESSION AT INTAKE AND FOLLOW-UP, REPORTS 2015-2022



## ANXIETY SYMPTOMS

To assess for generalized anxiety symptoms, participants were first asked:

“In the 6 months before you entered this program, did you worry excessively or were you anxious about multiple things on more days than not for all 6 months (like family, health, finances, school, or work difficulties)?”

Participants who answered “yes” were then asked 6 additional questions about anxiety symptoms (e.g., felt restless, keyed up or on edge, have difficulty concentrating, feel irritable).


In the 6 months before entering treatment, 71.4% of clients reported symptoms that met study criteria for generalized anxiety and 19.0% reported symptoms at follow-up – a significant decrease of 52.4% (see Figure 3.3).

Of those who met study criteria for anxiety at intake (n = 15), they had an average of 6.7 symptoms out of 7. At follow-up, among those who met study criteria for anxiety (n = 4), clients reported an average of 6.5 symptoms out of 7.

### Study Criteria for General Anxiety Disorder

To meet study criteria for general anxiety disorder, clients had to say “yes” to the one screening question and at least 3 of the other 6 symptoms. Thus, minimum score to meet study criteria: 4 out of 7.

FIGURE 3.3. CLIENTS MEETING STUDY CRITERIA FOR PAST-6-MONTH GENERALIZED ANXIETY AT INTAKE AND FOLLOW-UP (N = 21)




\*\*\*p < .01.

### TRENDS IN PAST-6-MONTH GENERALIZED ANXIETY

The percent of clients who met criteria for generalized anxiety at intake has remained consistent over from 2015 to 2018 (around three-quarters each year). In 2019, however, the percent of clients who met criteria for generalized anxiety at intake decreased to 68.3% before increasing again to 78.1% in 2020.

The percent of clients who met criteria for generalized anxiety at follow-up has fluctuated over time. About 15% of clients in 2015 and 21.6% of clients in 2016 met criteria for generalized anxiety at follow-up. In 2017, only 8.0% of clients met criteria for generalized anxiety at follow-up compared to 25.0% of clients in 2018. In 2020, the percent of clients who met criteria for generalized anxiety increased again (35.1%). Nearly one-third of clients met criteria for generalized anxiety at follow-up in 2021 (31.7%) and in 2022 the percent had decreased to 19.0%.


FIGURE 3.4. TRENDS IN THE NUMBER OF CLIENTS WHO MET STUDY CRITERIA FOR PAST-6-MONTH GENERALIZED ANXIETY AT INTAKE AND FOLLOW-UP, REPORTS 2015-2022



## COMORBID DEPRESSION AND ANXIETY SYMPTOMS

Figure 3.5 shows that at intake, 66.7% of clients met study criteria for both depression and generalized anxiety. There was a significant decrease of 57.2% to 9.5% at follow-up.

FIGURE 3.5. CLIENTS MEETING STUDY CRITERIA FOR COMORBID DEPRESSION AND GENERALIZED ANXIETY IN THE PAST 6 MONTHS AT INTAKE AND FOLLOW-UP (N = 21)




\*\*\*p < .01.

## SUICIDAL IDEATION AND/OR ATTEMPTS

Suicidal ideation and attempts were measured with self-reported questions about thoughts of suicide and actual attempts to commit suicide in the past 6 months. Close to 14% of clients reported suicidal ideation or attempts at intake and none of the clients reporting suicide ideation and/or attempts at follow-up (see Figure 3.6).

FIGURE 3.6. CLIENTS REPORTING SUICIDAL IDEATION AND/OR ATTEMPTS IN THE PAST 6 MONTHS AT INTAKE AND FOLLOW-UP (N = 21)



a - No measures of association could be computed for suicidal ideation and/or attempts because the value at follow-up was 0.


*“They have helped me with everything I have asked them to, they have never judged me, always answered all my questions, and helped with referrals.”*

KORTOS FOLLOW-UP CLIENTS

### TRENDS IN PAST-6-MONTH SUICIDE IDEATION AND/OR ATTEMPTS

The percent of clients who reported suicidal ideation and attempts at intake increased between 2015 and 2018 from 16.6% to 27.2%. In 2019, however, 13.2% of clients reported suicidal ideation and attempts at intake, but increased in 2020 before decreasing again in 2021. At follow-up, the percent of clients reporting suicidal ideation and attempts increased from 0.6% in 2017 to 4.9% in 2021. In 2022, none of the clients reported suicidal ideation and attempts at follow-up.


FIGURE 3.7. TRENDS IN THE NUMBER OF CLIENTS REPORTING SUICIDAL IDEATION AND/OR ATTEMPTS AT INTAKE AND FOLLOW-UP, REPORTS 2015-2022



### SELF-MEDICATION FOR MENTAL HEALTH SYMPTOMS

The majority of clients at intake (81.0%) reported using alcohol, prescription drugs, or illegal drugs to reduce stress, anxiety, worry, sadness, or fear. At follow-up, 23.8% of clients reported using alcohol, prescription drugs, or illegal drugs to reduce stress, anxiety, worry, sadness, or fear, which was a significant decrease of 57.2%.

FIGURE 3.8. CLIENTS WHO REPORTED THEY USED ALCOHOL, PRESCRIPTION DRUGS, OR ILLEGAL DRUG USE TO REDUCE STRESS, ANXIETY, WORRY, SADNESS, OR FEAR AT INTAKE AND FOLLOW-UP (N = 21)




\*\*\*p < .01.

## POST-TRAUMATIC STRESS DISORDER

Clients were asked to answer the four-item PTSD checklist about how bothered they had been about the symptoms in the prior 6 months.<sup>55</sup>

At intake, 19.0% of clients screened positive for PTSD symptoms at intake and 9.5% of clients screen positive for PTSD symptoms at follow-up (see Figure 3.9).

FIGURE 3.9. CLIENTS WHO SCREENED POSITIVE FOR POST-TRAUMATIC STRESS DISORDER SYMPTOMS IN THE PAST 6 MONTHS AT INTAKE AND FOLLOW-UP (n = 21)



### MENTAL HEALTH COMPARISONS BETWEEN CLIENTS WHO WERE STILL INVOLVED IN THE TREATMENT PROGRAM AT FOLLOW-UP AND CLIENTS WHO WERE NOT STILL INVOLVED

There was no significant difference between clients who were not involved in the clinic at follow-up and clients who were involved in the clinic for the percent of clients who met criteria for depression: 7.7% vs. 14.3%, respectively. Further, there was no significant difference between clients who were not involved in the clinic at follow-up and clients who were involved in the clinic for the percent of clients who met criteria for generalized anxiety: 19.0% vs. 7.7%, respectively. In addition, there was no significant difference between clients who were not involved in the clinic at follow-up and clients who were involved in the clinic for thoughts of suicide or suicide attempts with neither group reporting have thoughts of suicide or suicide attempts.


## GENERAL HEALTH STATUS

### OVERALL HEALTH

At both intake and follow-up, clients were asked to rate their overall health in the past 6 months from 1 = poor to 5 = excellent. Clients rated their health, on average, as 2.9 at intake and 3.2 at follow-up (not depicted in figure). Figure 3.10 shows that only 23.8% of clients rated their overall physical health as very good or excellent and 42.9% did at follow-up.

<sup>55</sup>Price, M., Szafranski, D., van Stolk-Cooke, K., & Gros, D. (2016). Investigation of an abbreviated 4 and 8-item version of the PTSD Checklist 5. *Psychiatry Research*, 239, 124-130.

FIGURE 3.10. CLIENTS' SELF-REPORT OF OVERALL HEALTH STATUS AT INTAKE AND FOLLOW-UP (N = 21)<sup>a</sup>




a - Significance tested with the Stuart-Maxwell Test for Marginal Homogeneity.

### PERCEPTIONS OF PHYSICAL AND MENTAL HEALTH

Clients were asked how many days in the past 30 days their physical and mental health were not good at intake and follow-up (see Figure 3.11). The number of days clients reported their physical health was not good decreased significantly from an average of 5.4 days to 1.7 days. The number of days clients' mental health was not good also decreased significantly from intake (13.0) to follow-up (4.3).

FIGURE 3.11. PERCEPTIONS OF POOR PHYSICAL HEALTH AND MENTAL HEALTH IN THE PAST 30 DAYS AT INTAKE AND FOLLOW-UP (N = 21)



\*p < .10, \*\*\*p < .01.


*It lets you go at your own pace depending on how you feel, you decide the dosage, counselors are always available to talk.*

KORTOS FOLLOW-UP CLIENTS

### TRENDS IN PERCEPTIONS OF POOR PHYSICAL HEALTH

The average number of days clients reported their physical health was poor in the past 30 days at intake has fluctuated over the past 8 years, ranging from 10.1 days to 14.4 days. The average number of days clients reported their physical health was poor in the past 30 days at follow-up decreased from 2.1 days in 2015 to 1.3 days in 2018, but increased in 2020 to 7.6 days before decreasing slightly to 1.7 days in 2022.


FIGURE 3.12. TRENDS IN PERCEPTIONS OF PHYSICAL HEALTH AT INTAKE AND FOLLOW, REPORTS 2015-2022



### TRENDS IN PERCEPTIONS OF POOR MENTAL HEALTH

The average number of days clients reported their mental health was not good in the past 30 days at intake has fluctuated from a low of 13.0 in 2022 to a high of 19.2 in 2018. The average number of days clients reported their mental health was poor in the past 30 days at follow-up has increased from 2.3 days in 2015 to 8.9 days in 2018. In 2019, however, the number of days clients reported their mental health was not good decreased at follow-up, but increased to 8.3 days in 2020. In 2021, the average number of days clients reported their mental health was poor in the past 30 days at follow-up decreased to 6.5 and continued to decrease in 2022 to 4.3.

FIGURE 3.13. TRENDS IN PERCEPTIONS OF MENTAL HEALTH AT INTAKE AND FOLLOW, REPORTS 2015-2022<sup>56</sup>




<sup>56</sup>In 2015, 3 cases had missing data for perceptions of mental health at intake, 2019 had one client with missing data, and 2020 had one client with missing data.

PERCEPTIONS OF POOR PHYSICAL OR MENTAL HEALTH LIMITING ACTIVITIES

Clients were also asked to report the number of days in the past 30 days poor physical or mental health had kept them from doing their usual activities. The number of days clients reported their physical or mental health kept them from doing their usual activities decreased significantly from 8.6 days at intake to 3.5 days at follow-up (see Figure 3.14).

FIGURE 3.14. PERCEPTIONS OF POOR PHYSICAL HEALTH AND MENTAL HEALTH LIMITING ACTIVITIES IN THE PAST 30 DAYS AT INTAKE AND FOLLOW-UP (N = 21)




\*p < .10.

TRENDS IN NUMBER OF DAYS POOR PHYSICAL OR MENTAL HEALTH KEPT CLIENT FROM DOING USUAL ACTIVITIES

The average number of days in the past 30 days at intake clients reported their poor physical or mental health kept them from doing their usual activities has decreased from a high of 14.8 in 2018 to a low of 8.6 in 2022. The average number of days in the past 30 days at follow-up clients reported their poor physical or mental health kept them from doing their usual activities was a low of 1.8 in 2015 to a high of 4.5 in 2020. Nonetheless, the average number of days poor physical or mental health kept clients from doing their usual activities decreased significantly from intake to follow-up each report year.

FIGURE 3.15. TRENDS IN THE NUMBER OF DAYS POOR PHYSICAL OR MENTAL HEALTH KEEP CLIENT FROM DOING USUAL ACTIVITIES AT INTAKE AND FOLLOW, REPORTS 2015-2022<sup>57</sup>



<sup>57</sup>In 2015, one case had a missing value, in 2019 five cases had a missing value, and in 2020 two cases had a missing value for this item at follow-up.




## CHRONIC PAIN

The percent of clients who reported chronic pain that was persistent and lasted at least 3 months decreased significantly from intake to follow-up by 28.6% (see Figure 3.16). At intake, 47.6% of clients reported chronic pain and that percent dropped to 19.0% by follow-up. Of those clients who reported chronic pain at intake (n = 10), clients reported that the pain started around the age of 29. In addition, clients reported experiencing chronic pain for almost 26 of the 30 days before entering the program. On a scale of 0 (no pain) to 10 (pain as bad as you can image), clients reported an average of 7.2 intensity in the 30 days before entering the program (not shown in the figure).

Of those clients who reported chronic pain at follow-up (n = 4), clients reported experiencing chronic pain for about 26 of the past 30 days. On a scale of 0 (no pain) to 10 (pain as bad as you can image), clients reported an average of 6.5 intensity in the past 30 days (not shown in the figure).

FIGURE 3.16. CLIENTS REPORTING CHRONIC PAIN AT INTAKE AND FOLLOW-UP (N = 21)




\*p < .10.

### TRENDS IN CHRONIC PAIN

Overall, the percent of clients who reported chronic pain increased over time at intake. In the 2015 report, 39.9% of clients reported chronic pain compared to 46.6% in 2016. In 2017, the percent of clients reporting chronic pain decreased slightly to 42.3% and then increased to 52.7% in 2018 and increased again in 2021 to 56.1%. In 2022, the number of clients reporting chronic pain decreased to 47.6%.

At follow-up, 18.8% of clients reported chronic pain in 2015 and 20.3% of clients reported chronic pain in 2016. The percent of clients reporting chronic pain at follow-up decreased to 12.6% in 2017, but more than doubled in 2018 with 30.4% of clients reporting chronic pain. In 2019, the percent of clients reporting chronic pain at follow-up decreased to 25.2%, but increased in 2020 to 35.1%. In 2022, the percent of clients reporting chronic pain at follow-up decreased to 19.0%.

FIGURE 3.17. TRENDS IN THE NUMBER OF CLIENTS REPORTING CHRONIC PAIN AT INTAKE AND FOLLOW-UP, REPORTS 2015-2022<sup>58</sup>



### Prescription Opioid Misuse and Chronic Pain

Of those who were not incarcerated all 180 days before entering the program or in the 6 months before follow-up and misused prescription opioids at intake (n = 11), 63.6% reported chronic pain in the 6 months before entering the program and 27.3% experienced chronic pain at follow-up.


Additionally, among clients who were not incarcerated all 180 days before entering the program or at follow-up, reported misusing prescription opioids, and experienced chronic pain at intake (n = 7), 42.9% (n = 3) reported chronic pain in the past 6 months at follow-up and 14.3% (n = 1) reported past-6-month misuse of prescription opioids.

### SELF-MEDICATION FOR PHYSICAL PAIN

The majority of clients at intake (85.7%) reported using alcohol, prescription drugs, or illegal drugs to reduce their physical pain. At follow-up, 19.0% of clients reported using alcohol, prescription drugs, or illegal drugs to reduce their physical pain, which was a significant decrease of 66.7%.

<sup>58</sup>In 2018, one client was missing information on chronic pain at follow-up.

FIGURE 3.18. CLIENTS WHO REPORTED THEY USED ALCOHOL, PRESCRIPTION DRUGS, OR ILLEGAL DRUG USE TO REDUCE PHYSICAL PAIN AT INTAKE AND FOLLOW-UP (N = 21)




\*\*\*p < .01.

## HEALTH INSURANCE

The majority of KORTOS clients reported they had health insurance through Medicaid at intake (57.1%) and follow-up (61.9%; see Figure 3.19). Around 15% of clients at intake and 4.8% of clients at follow-up did not have any insurance. Nineteen percent of clients at intake and 28.6% of clients at follow-up reported they had health insurance through an employer (including spouse’s, parents’, and self-employed). A minority of clients had insurance at either point through Medicare and none had VA/Champus/Tricare.

FIGURE 3.19. HEALTH INSURANCE FOR KORTOS CLIENTS AT INTAKE AND FOLLOW-UP (N = 21)



Of those clients who were employed full-time at intake (n = 8), 37.5% had insurance through an employer (including a spouse’s or parent’s employer). In addition, 50.0% of clients who were employed full-time at intake indicated they had insurance through Medicaid. At follow-up, of those clients employed full-time (n = 9), 66.7% had insurance through an employer. One-third of clients, however, had insurance through Medicaid.


**PHYSICAL HEALTH COMPARISONS BETWEEN CLIENTS WHO WERE STILL INVOLVED IN THE TREATMENT PROGRAM AT FOLLOW-UP AND CLIENTS WHO WERE NOT STILL INVOLVED**

There was not a significant difference for experiencing chronic pain (i.e., pain that lasted more than 3 months) at follow-up between clients who were not involved in the clinic at follow-up and clients who were. In addition, there were no significant differences for the average number of days physical health or mental health was poor in the past 30 days at follow-up between the two groups. In addition, there was no difference between the groups for medical insurance with the majority of clients having Medicaid at follow-up.

**QUALITY OF LIFE RATINGS**

At intake and follow-up, clients were asked to rate their quality of life at the time of the interview. Ratings were from 1 = ‘Worst imaginable’ to 5 = ‘Good and bad parts were about equal’ to 10 = ‘Best imaginable’. KORTOS clients rated their quality of life as a 4.7 at intake (see Figure 3.20). The average quality of life rating significantly increased to 8.3 at follow-up.

FIGURE 3.20. PERCEPTION OF QUALITY OF LIFE AT INTAKE AND FOLLOW-UP (N = 21)



\*\*\*p < .01.


*“They’re not perfect but no place is. They really do make you feel like a human, not like a junkie. They do try to help you, I love all the staff there.”*

KORTOS FOLLOW-UP CLIENTS

### TRENDS IN QUALITY OF LIFE RATING

Clients were asked to rank their overall quality of life on a scale from 1 (worst imaginable) to 10 (best imaginable) at both intake and follow-up. At intake, KORTOS clients have consistently rated their quality of life, on average, close to 4.0. At follow-up, that rating has significantly increased to an average of about a 7.7 (see Figure 3.21).

FIGURE 3.21. TRENDS IN QUALITY OF LIFE RATING AT INTAKE AND FOLLOW, REPORTS 2015-2022



### QUALITY OF LIFE AND WELL-BEING COMPARISONS BETWEEN CLIENTS WHO WERE STILL INVOLVED IN THE TREATMENT PROGRAM AT FOLLOW-UP AND CLIENTS WHO WERE NOT STILL INVOLVED

There was no significant difference between clients who were still involved in the clinic and follow-up and those who were not for average quality of life rating (8.3 vs 8.1, respectively).

## SECTION 4. CRIMINAL JUSTICE SYSTEM INVOLVEMENT AND INTERPERSONAL VICTIMIZATION


This section describes change in client involvement with the criminal justice system during the 6-month period before entering treatment and the 6-month period before the follow-up interview. Specifically, results include changes in: (1) any arrest, (2) the number of times arrested, among clients with any arrests, (3) any incarceration, (4) the number of nights incarcerated, among clients with any incarceration, (5) criminal justice supervision status, (6) interpersonal victimization, and (7) personal safety.

### ARRESTS

#### ANY ARRESTS IN THE PAST 6 MONTHS

Less than 5% of clients (n = 1) reported any arrests in the 6 months before entering treatment and 9.5% of clients (n = 2) reported any arrests in the 6 months before follow-up (see Figure 4.1).


FIGURE 4.1. PAST-6 MONTH ARRESTS AT INTAKE AND FOLLOW-UP (N = 21)



#### TRENDS IN PAST-6-MONTH ARRESTS

While the percent of clients reporting an arrest in the past 6 months at intake has been stable overall in the report years 2015 through 2020, the percent of clients reporting an arrest in the past 6 months at intake increased to 24.4% in 2021. At follow-up, the percent of clients reporting arrest has remained low over since 2015 (see Figure 4.2).

FIGURE 4.2. TRENDS IN THE NUMBER OF CLIENTS REPORTING ANY ARRESTS IN THE PAST-6-MONTHS AT INTAKE AND FOLLOW-UP, REPORTS 2015-2022<sup>59, 60</sup>



<sup>59</sup>In 2019, one client refused to answer criminal justice system involvement questions at follow-up and one client was missing data on criminal justice questions at follow-up.


<sup>60</sup>The small sample size in report year 2022 may be affecting the number of clients reporting an arrest.

## INCARCERATION

### INCARCERATION IN THE PAST 6 MONTHS

About 10% of clients reported they had spent at least one night in jail or prison at intake. At follow-up, 4.8% of clients reported they had spent at least one night in jail or prison in the past 6 months (see Figure 4.3).


FIGURE 4.3. CLIENTS REPORTING INCARCERATION AT INTAKE AND FOLLOW-UP (N = 21)



### TRENDS IN PAST-6-MONTH INCARCERATION

The percent of clients reporting spending at least one night in jail or prison has been relatively steady since 2015 with less than 2 in 10 clients reporting an incarceration at intake. At follow-up, relatively few clients reported being incarcerated in the past 6 months and in 2021, none of the clients reported being incarcerated (see Figure 4.4).


FIGURE 4.4. TRENDS IN THE NUMBER OF CLIENTS REPORTING ANY INCARCERATION IN THE PAST-6-MONTHS AT INTAKE AND FOLLOW-UP, REPORTS 2015-2022<sup>61</sup>



## CRIMINAL JUSTICE SYSTEM SUPERVISION

Close to 10% of clients at intake and none of the clients at follow-up self-reported they were under criminal justice system supervision (e.g., probation or parole; see Figure 4.5).

FIGURE 4.5. CLIENTS REPORTING CRIMINAL JUSTICE SYSTEM SUPERVISION AT INTAKE AND FOLLOW-UP (N = 21)



<sup>61</sup>In 2019, one client declined to answer criminal justice system involvement questions at follow-up and one client was missing data on criminal justice questions at follow-up.

### CRIMINAL JUSTICE SYSTEM INVOLVEMENT COMPARISONS BETWEEN CLIENTS WHO WERE STILL INVOLVED IN THE TREATMENT PROGRAM AT FOLLOW-UP AND CLIENTS WHO WERE NOT STILL INVOLVED


Around 8% of clients who not involved in the clinic at follow-up and none of the clients who were still involved in the clinic were under supervision by the criminal justice system at follow-up (e.g., probation, parole), with no significant difference between groups.

There was no difference between the groups for the number of clients who were arrested for any charge or incarcerated for at least one night in the past 6 months at follow-up.

### INTERPERSONAL VICTIMIZATION EXPERIENCES

Clients were asked about several types of interpersonal victimization<sup>62</sup> (including when they may have been the victim of a crime, harmed by someone else, or felt unsafe) in the 6 months before entering programs and in the 6 months before follow-up (see Figure 4.6). Because relatively small percentages of clients reported each specific type of victimization experience in the 6-month periods, the items were collapsed. The percent of clients who reported experiencing any victimization in the past 6 months decreased from 9.5% at intake to 0.0% at follow-up.

FIGURE 4.6. CLIENTS REPORTING PAST-6-MONTH CRIME AND INTERPERSONAL VICTIMIZATION (N = 21)



a - No measures of association could be computed for victimization use because the value at follow-up was 0.


### WORRY ABOUT PERSONAL SAFETY

At intake, 52.4% of clients reported they were worried about their personal safety. None of clients, however, were worried about personal safety a great deal (not displayed in a figure). At follow-up, 38.1% of clients were worried about their personal safety (which was not a significant decrease). Only 4.8% of clients reported they were worried about personal safety a great deal at follow-up (not displayed in a figure).

<sup>62</sup>Victimization includes being robbed or mugged by force, assaulted with or without a weapon, threatened with a gun, intimate partner violence, stalking, sexually harassed or assaulted.



FIGURE 4.7. CLIENTS WHO WORRIED ABOUT PERSONAL SAFETY AT INTAKE AND FOLLOW-UP  
(N = 21)




## SECTION 5. EDUCATION, ECONOMIC STATUS, AND LIVING CIRCUMSTANCES

*This section examines changes in education, economic status, and living circumstances from intake to follow-up including: (1) highest level of education completed, (2) the number of months clients were employed full-time or part-time in the past 6 months, (3) current employment status, (4) hourly wage, (5) homelessness, (6) living situation, and (7) economic hardship (i.e., difficulty meeting living and health care needs for financial reasons).*

### EDUCATION

At intake, the average number of years of education was 13.6 years and at follow-up client reported 13.9 years, where 12 = High school diploma or GED (not depicted in a figure). Another way to examine change in education is to examine change in the percent of clients who reported different levels of education. There was not a significant change in the percent of clients who reported attending or completing vocational school, college, or graduate school from intake to follow-up (see Figure 5.1).

FIGURE 5.1. HIGHEST LEVEL OF EDUCATION COMPLETED AT INTAKE AND FOLLOW-UP (n = 21)<sup>a63</sup>



a - Significance tested with the Stuart-Maxwell Test of Overall Marginal Homogeneity


### EMPLOYMENT

#### AVERAGE NUMBER OF MONTHS EMPLOYED IN THE PAST 6 MONTHS

At both intake and follow-up, clients were asked to report the number of months in the past 6 months they were employed at least part-time. Figure 5.2 shows there was not a significant change in the average number of months clients reported they were employed from intake (3.0) to follow-up (3.1).

<sup>63</sup>One client was missing data for education level at follow-up because of data inconsistencies compared to intake.

FIGURE 5.2. AVERAGE NUMBER OF MONTHS EMPLOYED IN THE PAST 6 MONTHS AT INTAKE AND FOLLOW-UP (N = 21)




The majority of clients at intake (81.0%) and follow-up (85.7%) reported that they expected to be employed in the next 6 months.

### TRENDS IN EMPLOYMENT BY GENDER

For the past 8 years, significantly fewer women reported being employed (full- or part-time) at least one month in the past 6 months at intake compared to men. In the 2016 report year, only 37.9% of women were employed at least one month in the past 6 months at intake while 75.0% of men reported employment. In 2018, however, the gap narrowed with 67.4% of men reporting employment compared to 53.3% of women. Less than half of women in the 2019 sample reported being employed at intake compared to 64.1% of men and in 2020 and 2021 the gap between men and women reporting employment widened further. However, in 2022, the gap decreased and was the smallest it has been over the 8 years.

By follow-up, around half of women reported they were employed full-time or part-time at least one month in the past 6 months but significantly more men reported employment during that same time frame. Since 2016, the percent of women who reported being employed at least one month at follow-up has increased. In 2019, almost two-thirds of women reported being employed at least one month at follow-up. In 2020, however, less than half of female clients reported being employed at follow-up compared to over three-quarters of men. In 2021, the gap between men and women for employment at follow-up decreased slightly compared to 2020. For the first time in the 8 years, in 2022, the percent of women who reported being employed at follow-up was higher than the percent of men.

FIGURE 5.3. TRENDS IN GENDER DIFFERENCES IN CLIENTS EMPLOYED AT LEAST ONE MONTH AT INTAKE AND FOLLOW-UP<sup>64</sup>




<sup>64</sup> The small sample size in report year 2022 may be affecting the number of clients who reported being employed at least one month.

### CURRENT EMPLOYMENT STATUS


At intake, 57.1% of clients were not employed (see Figure 5.4) in the 30 days before they entered the program and at follow-up, the percent of clients who were not employed was 47.6%. Additionally, the percent of clients who were employed full-time increased slightly, but not significantly, from 38.1% at intake to 42.9% at follow-up.

FIGURE 5.4. CURRENT EMPLOYMENT STATUS AT INTAKE AND FOLLOW-UP (N = 21)



Of those not employed at each point, clients were asked why they were not currently employed. At intake (n =12), 58.3% were unemployed, but were looking for a job, 16.7% were keeping house or caring for children, and 8.3% of clients reported they were unemployed, but not looking for work. Of clients not employed at follow-up (n = 9), 44.4% reported they were on disability or had applied for disability, 33.3% were keeping house or caring for children, and 22.2% were unemployed, but looking for work.

FIGURE 5.5. REASONS FOR UNEMPLOYMENT STATUS AT EACH POINT




### HOURLY WAGE

Of those clients who were employed at intake (n = 9), the median hourly wage was \$15.75. Of those employed at follow-up and who reported their hourly wage (n = 9),<sup>65</sup> the median hourly wage was \$17.60 (see Figure 5.6).

<sup>65</sup> While 11 clients were employed at follow-up, 2 clients were missing data for hourly wage at follow-up.


FIGURE 5.6. CURRENT MEDIAN HOURLY WAGE AT INTAKE AND FOLLOW-UP, AMONG EMPLOYED CLIENTS



GENDER DIFFERENCES IN OCCUPATION TYPE

In previous reports, median hourly wages were significantly higher for men compared to women. No statistically significant difference was found this year, likely due to the small sample size. At least part of the reason for the marked difference in hourly wages between men and women in previous reports was due to the difference in occupation type by gender for employed clients. The same difference by gender was found in this year’s sample.<sup>66</sup> At intake, 40.0% of employed women reported having a service job (e.g., waiter/waitress, childcare, housekeeping, hair stylist, etc.) while none of the employed men reported having a service job (see Figure 5.7a). Half of men reported working in production, transportation, and material moving e.g., factory production line, power plant, bus driver, welder, sanitation worker, etc.) compared to none of the women. In addition, 20.0% of women reported having a professional job while none of the men reported having a professional job.


FIGURE 5.7a. AMONG EMPLOYED INDIVIDUALS, TYPE OF OCCUPATION BY GENDER AT INTAKE



At follow-up, the difference in occupation type was similar. Two-thirds of employed women reported having a service job whereas none of the employed men had a service job (see Figure 5.7b). Identical to intake, half of employed men and none of the employed women reported having a production, transportation, and material moving jobs (e.g., factory production line, power plant, bus driver, welder, sanitation worker, etc.).

<sup>66</sup>Occupation type was asked only of individuals who reported they were currently employed at intake and at follow-up.

FIGURE 5.7B. AMONG EMPLOYED INDIVIDUALS, TYPE OF OCCUPATION BY GENDER AT FOLLOW-UP




## LIVING CIRCUMSTANCES

### HOMELESSNESS

None of the clients reported at treatment intake or in the past 6 months at follow-up that they were homeless at some point in the past 6 months (see Figure 5.8).

FIGURE 5.8. CLIENTS REPORTING HOMELESSNESS IN THE PAST 6 MONTHS AT INTAKE AND FOLLOW-UP (N = 21)



*It has worked really well for me, other programs haven't been successful.*


KORTOS FOLLOW-UP CLIENTS

### TRENDS IN PAST-6-MONTH HOMELESSNESS

From 2015 to 2017, the percent of clients reporting that they considered themselves homeless in the past 6 months at intake remained steady at around 16% of clients. In 2018, the percent of clients who considered themselves homeless at intake increased slightly to 23.9%, but decreased again in 2019 to 15.7%. In 2020, the percent of clients reporting that they considered themselves homeless increased again to 24.6%, but decreased to 9.8% in 2021. None of the clients considered themselves homeless at intake in 2022.

At follow-up each year, very few clients reported that they considered themselves homeless. In both 2015 and 2018, only 2.2% of clients considered themselves homeless in the past 6 months at follow-up. The percent of clients who considered themselves homeless at follow-up increased to 8.3% in 2019 and again to 8.8% in 2020. In 2021, however, the percent of clients who considered themselves homeless at follow-up decreased to 4.9% and decreased further to 0.0% in 2022.


FIGURE 5.9. TRENDS IN THE NUMBER OF CLIENTS REPORTING HOMELESSNESS IN THE PAST-6-MONTHS AT INTAKE AND FOLLOW-UP, REPORTS 2015-2022



### LIVING SITUATION IN THE PAST 6 MONTHS

Figure 5.10 shows that about 43% of clients in the past 6 months at intake and 71.4% of clients at follow-up reported they were living in their own home or apartment, which was not a significant increase. Over half of clients at intake (52.4%) and 28.6% of clients at follow-up reported living in someone else’s home or apartment. Only 4.8% of client at intake lived in another situation such as jail, a treatment program, shelter, or on the street.

FIGURE 5.10. TYPE OF TYPICAL LIVING SITUATION IN THE PAST 6 MONTHS AT INTAKE AND FOLLOW-UP (N = 21)






## ECONOMIC HARDSHIP

Less than 5% of clients at intake and 9.5% of clients at follow-up reported they were currently receiving SSI or SSDI benefits (not presented in a figure).

Clients were also asked eight items, five of which asked about the clients' difficulty meeting basic living needs such as food, shelter, utilities, and telephone, while three items asked about the clients' difficulty obtaining health care for financial reasons.

Less than half of clients (47.6%) reported difficulty meeting basic living needs (e.g., shelter, utilities, phone, food) at intake and follow-up (see Figure 5.11). The percent of clients who reported difficulty meeting health care needs (e.g., doctor visits, dental visits, and prescription medications) for financial reasons decreased, but not significantly, from 47.6% at intake to 28.6% at follow-up.

FIGURE 5.11. DIFFICULTY MEETING BASIC LIVING NEEDS AND HEALTH CARE NEEDS FOR FINANCIAL REASONS IN THE PAST 6 MONTHS AT INTAKE AND FOLLOW-UP (N = 21)



### TRENDS IN DIFFICULTY MEETING BASIC LIVING AND HEALTH CARE NEEDS FOR FINANCIAL REASONS

For each of the past 8 years, there has been a significant decrease in the percent of KORTOS clients who reported they had difficulty meeting basic living needs and/or health care needs in the past 6 months from intake to follow-up.


In general, half of clients reported having difficulty meeting basic living needs (e.g., shelter, utilities, phone, food) at intake. In 2019, 46.1% of clients reported having difficulty which increased to 54.0% in 2020 and again in 2021 to 56.1%. At follow-up, overall, trends show that the number of clients who have reported having difficulty meeting basic living needs is increasing over time. In 2018, the percent of clients who reported having difficulty meeting basic living needs increased to 33.2% from 15.2% in 2017 and continued to increase through 2022.

From 2015 to 2017, the percent of clients reporting difficulty meeting basic health care needs (i.e., doctor visits, dental visits, and prescription medications) at follow-up decreased; however, beginning in 2018, the percent of clients reporting difficulty meeting basic needs has increased. Whereas less than 5% of clients reported difficulty meeting basic health care needs in 2017, 29.3% of clients reported difficulty meeting basic health care needs in 2021 and 28.6% in 2022.

FIGURE 5.12. TRENDS IN THE NUMBER OF CLIENTS REPORTING ECONOMIC DIFFICULTY IN THE PAST-6-MONTHS AT INTAKE AND FOLLOW-UP, REPORTS 2015-2022

Difficulty Meeting Basic Living Needs

Difficulty Meeting Basic Health Care Needs



### EDUCATION, EMPLOYMENT, AND LIVING SITUATION COMPARISONS BETWEEN CLIENTS WHO WERE STILL INVOLVED IN THE TREATMENT PROGRAM AT FOLLOW-UP AND CLIENTS WHO WERE NOT STILL INVOLVED

There was a significant difference in the average number of years of education at follow-up where clients who were not still involved in the treatment program reported more years of education (15.6) compared to clients who were still involved in the clinic at follow-up (13.9). Five percent of clients who were still involved in the clinic at follow-up and none of the clients who were no longer involved in the clinic reported less than a high school diploma or GED at follow-up. There were no differences between clients who were not involved in the clinic at follow-up and clients who were still involved at follow-up on employment in the past 30 days at follow-up.

The majority of clients in both groups reported that their usual living arrangement in the past 6 months at follow-up was living in their own home or apartment. There were also no differences between the groups on clients who considered themselves homeless at follow-up.

Finally, there were no significant difference between clients who were not involved in the clinic at follow-up and clients who were on difficulty meeting basic living needs or for being unable to receive needed health care for financial reasons at follow-up.

## SECTION 6. RECOVERY SUPPORTS


*This section focuses on four main changes in recovery supports: (1) percent of clients attending mutual help recovery group meetings, (2) the number of people the client said they could count on for recovery support, (3) what will be most useful to the client in staying off drugs/alcohol, and (4) clients' perceptions of their chances of staying off drugs/alcohol.*

### MUTUAL HELP RECOVERY GROUP MEETING ATTENDANCE

At intake, 9.5% of clients reported going to mutual help recovery group meetings (e.g., AA, NA, or faith-based) in the past 30 days (see Figure 6.1). At follow-up, 14.3% of clients reported they had gone to mutual help recovery group meetings.

Among clients who had attended mutual help recovery group meetings at intake (n = 2), they reported attending an average of 8.5 meetings. Among clients who attended mutual help recovery group meetings at follow-up (n = 3), they reported attending an average of 17.7 meetings.


FIGURE 7.1. CLIENTS REPORTING MUTUAL HEALTH RECOVERY GROUP ATTENDANCE AT INTAKE AND FOLLOW-UP (N = 21)



### NUMBER OF PEOPLE CLIENT CAN COUNT ON FOR RECOVERY SUPPORT

The average number of people clients reported they could count on for recovery support increased from intake (3.8) to follow-up (8.0; see Figure 6.2).

FIGURE 6.2. AVERAGE NUMBER OF PEOPLE CLIENT COULD COUNT ON FOR RECOVERY SUPPORT AT INTAKE AND FOLLOW-UP (N = 20)<sup>67\*\*\*</sup>




\*\*\*p < .01.

<sup>67</sup>One client responded “don’t know” on how many people the client could on for recovery support.

## CHANCES OF STAYING OFF DRUGS/ALCOHOL

Clients were asked, based upon their situation, how good they believed their chances were of getting off and staying off drugs/alcohol using a scale from 1 (Very poor) to 5 (Very good). Clients rated their chances of getting off and staying off drugs/alcohol as a 4.4 at intake and a 4.6 at follow-up, which was a significant increase (not depicted in figure). Overall, 81.0% of clients at intake and 90.5% of clients at follow-up believed they had moderately or very good chances of staying off drugs/alcohol (see Figure 6.3).


FIGURE 6.3. CLIENTS REPORTING THEIR CHANCES OF GETTING OFF AND STAYING OFF DRUGS/ALCOHOL AT INTAKE AND FOLLOW-UP (N = 21)



## WHAT WILL BE MOST USEFUL IN STAYING OFF DRUGS/ALCOHOL

At intake and follow-up, clients were asked what, other than medication-assisted treatment, they believed would be most useful in helping them quit or stay off drugs/alcohol. Rather than conduct analysis on change in responses from intake to follow-up, the top categories during each time period are presented for descriptive purposes in Figure 6.4. The most common responses at intake were counseling, employment, and support from a partner. At follow-up, the most common responses were the need to stay out of jail, employment, and support from friends.

FIGURE 6.5. CLIENTS REPORTING WHAT WILL BE MOST USEFUL IN STAYING OFF DRUGS AND/OR ALCOHOL (N = 21)



### RECOVERY SUPPORT COMPARISONS BETWEEN CLIENTS WHO WERE STILL INVOLVED IN THE TREATMENT PROGRAM AT FOLLOW-UP AND CLIENTS WHO WERE NOT STILL INVOLVED

There were significant differences between clients who were still involved in the clinic and clients who were not involved in the clinic at follow-up for the percent of clients attending mutual help recovery meetings in the past 30 days at follow-up (14.3% vs. 61.5%, respectively). There was no significant difference between the two groups for the average number of people clients could count on for support at follow-up. There was also no significant difference for how good they believed their chances were of getting off and staying off drugs/alcohol; clients who were still involved in the clinic rated their chances of getting off and staying off drugs/alcohol as a 4.6 and clients who were not still involved in the clinic at follow-up rated their chances as 4.5.

## SECTION 7. MULTIDIMENSIONAL RECOVERY

*This section examines multidimensional recovery that takes into account severity of substance use disorder, employment, homelessness, criminal justice system involvement, suicide ideation, overall health, recovery support, and quality of life. Change in recovery status from intake to follow-up is presented. Furthermore, a multivariate analysis was conducted to examine the intake indicators of recovery status and their association with having all positive dimensions of recovery at follow-up. Furthermore, a multivariate analysis was conducted to examine the intake indicators of recovery status and their association with having all eight dimensions of recovery at follow-up.*

Recovery goes beyond relapse or return to occasional drug or alcohol use. Recovery from substance use disorders can be defined as “a process of change through which an individual achieves abstinence and improved health, wellness and quality of life: (p. 5).<sup>68</sup> The SAMHSA definition of recovery is similarly worded and encompasses health (including but not limited to abstinence from alcohol and drugs), having a stable and safe home, a sense of purpose through meaningful daily activities, and a sense of community.<sup>69</sup> In other words, recovery encompasses multiple dimensions of individuals’ lives and functioning. The multidimensional recovery measure uses items from the intake and follow-up surveys to create one index that can be used to classify individuals who all positive dimensions of recovery.

TABLE 7.1. MULTIDIMENSIONAL RECOVERY


INDICATOR	POSITIVE RECOVERY DIMENSIONS	NEGATIVE RECOVERY DIMENSIONS
Substance use .....	No or mild substance use disorder (SUD)	Mild, moderate, or severe substance use disorder (SUD)
Employment.....	Employed at least part-time or in school	Unemployed (not on disability, not going to school, not a caregiver)
Homelessness.....	No reported homelessness	Reported homelessness
Criminal Justice System Involvement.....	No arrest or incarceration	Any arrest or incarceration
Suicide ideation .....	No suicide ideation (thoughts or attempts)	Any suicide ideation (thoughts or attempts)
Overall health.....	Fair to excellent overall health	Poor overall health
Recovery support.....	Had at least one person he/she could count on for recovery support	Had no one he/she could count on for recovery support
Quality of life .....	Mid to high-level of quality of life	Low-level quality of life

At intake, none of the clients were classified as having all positive dimensions of recovery when entering treatment (see Figure 7.1). At follow-up, 47.6% of clients were classified as having all positive dimensions of recovery at follow-up.

<sup>68</sup>Center on Substance Abuse Treatment. (2007). *National summit on recovery: conference report* (DHHS Publication No. SMA 07-4276). Rockville, MD: Substance Abuse and Mental Health Services Administration.

<sup>69</sup>Laudet, A. (2016). *Measuring recovery from substance use disorders*. Workshop presentation at National Academies of Sciences, Engineering, and Medicine (February 24, 2016). Retrieved from [https://sites.nationalacademies.org/cs/groups/dbassesite/documents/webpage/dbasse\\_171025.pdf](https://sites.nationalacademies.org/cs/groups/dbassesite/documents/webpage/dbasse_171025.pdf)

FIGURE 7.1. MULTIDIMENSIONAL RECOVERY AT INTAKE AND FOLLOW-UP (N = 21)<sup>a</sup>



a- No measure of association could be computed for the cross tabulation because there was a value of 0 for the multidimensional recovery variable at intake.

Table 7.2 presents the frequency of clients who reported each of the specific components of the multidimensional recovery measure at intake and follow-up. At intake, the factors with the lowest percent of individuals indicated were no substance use disorder, a higher quality of life, having employment full-time and part-time. At follow-up, the factors with the lowest percent of individuals reporting the positive dimensions of recovery were no substance use disorder, and having employment full-time and part-time.

TABLE 7.2. PERCENT OF CLIENTS WITH SPECIFIC POSITIVE DIMENSIONS OF RECOVERY AT INTAKE AND FOLLOW-UP (N = 21)

Factor	Intake Yes	Follow-up Yes
Met DSM-5 criteria for no SUD in the past 6 months	0.0%	71.4%
Usual employment was employed full-time or part-time in the past 6 months (or unemployed because a student, home caregiver, on disability)	85.7%	71.4%
Reported no homelessness (or living in recovery center at follow-up)	100.0%	100.0%
Reported not being arrested and/or incarcerated in the past 6 months	90.5%	90.5%
Reported no thoughts of suicide or attempted suicide in the past 6 months	85.7%	100.0%
Self-rating of overall health at follow-up was fair, good, very good, or excellent	90.5%	95.2%
Reported having someone they could count on for recovery support	95.2%	100.0%
Reported a quality-of-life rating in the mid or higher range (rating of 5 or higher)	57.1%	100.0%

**MULTIDIMENSIONAL RECOVERY STATUS COMPARISONS BETWEEN CLIENTS WHO WERE STILL INVOLVED IN THE TREATMENT PROGRAM AT FOLLOW-UP AND CLIENTS WHO WERE NOT STILL INVOLVED**

There were no significant differences between clients who were still involved in the clinic and clients who were not involved in the clinic at follow-up for multidimensional recovery. Almost half of clients who were still involved in the clinic at follow-up (47.6%) and 69.2% of clients who were not still involved in the clinic at follow-up were classified as all positive dimensions of recovery.



## SECTION 8. CLIENT SATISFACTION WITH OPIOID TREATMENT PROGRAMS


At the beginning of the follow-up interview, clients were asked to give their opinions and feedback regarding their program experience. The items measured in this report include: (1) client involvement in the program, (2) if the client would refer someone else to the program, (3) client ratings of program experiences, and (4) positive and negative aspects of program participation.

### CLIENT INVOLVEMENT IN THE PROGRAM

Clients reported having been involved in the treatment program an average of 7.1 months at follow-up (range of 3 to 12 months).<sup>70</sup>

Figure 8.1 shows the percent of clients who reported the program started poor or good and ended poor or good. The majority of clients reported that the program started good (76.2%) and 95.2% reported it was currently good.

FIGURE 8.1. PERCENT OF CLIENTS WHO REPORTED AT FOLLOW-UP THE TREATMENT STARTED AND IS CURRENTLY POOR OR GOOD (N = 21)



All of the followed-up clients reported that the treatment episode is working pretty well or extremely well for them.

### RECOMMENDATION TO THE PROGRAM

The majority of clients (95.2%) indicated they would refer a close friend or family member to their treatment provider. Of the clients who reported they would refer a close friend or family member to the program (n = 20), 15.0% reported they would warn their friend or family member about certain things or tell them who to work with or who to avoid.

### CLIENT RATINGS OF PROGRAM EXPERIENCES

Clients were asked to report their perceptions of how the treatment program worked for them. The statements presented in Figure 8.2 had separate response options, with ratings

<sup>70</sup>One client did not know how long they were involved in the treatment program at follow-up.


ranging from 0 to 10. The higher values corresponded to the more positive responses and the lower values corresponded to the negative responses. For example, for the statement, “My expectations and hopes for treatment and recovery were met” the anchors were 0 “Not at all met” and 10 “Perfectly met.” Even the negatively worded items had anchors in which the higher values represented the more positive side of the continuum. For example, for the statement, “There were things I did not talk about or that I did not fully discuss with my counselor/program staff” the response option 0 corresponds to “I did not discuss lots of things, I held things back,” and 10 corresponds to “I discussed everything, I held back nothing.” Clients who rated each statement from 8 to 10 are shown in the figure below.

FIGURE 8.2. RATINGS OF PROGRAM TREATMENT EXPERIENCE (N = 21)



Clients rated their overall program experience, on average, as 9.1 (see Figure 8.3). Overall, 95.3% gave a rating between 8 and 10 and 42.9% of clients gave the highest possible rating, 10.


FIGURE 8.3. RATING OF EXPERIENCE AT THE PROGRAM (n = 21)



### POSITIVE AND NEGATIVE ASPECTS OF PROGRAM


Clients were asked to identify the three most positive aspects of their participation in the program (Figure 8.4). Almost two-thirds of clients (61.9%) reported that reduction in substance use was a positive outcome, 47.6% of clients reported improved mental health and feelings about themselves and improved interactions and relationships with others was a positive aspect. Close to 43% of clients reported the quality of treatment was a positive aspect. Nineteen percent of clients reported that changes in their financial situation were positive aspects and 19.0% reported other positive aspects (meeting other people in recovery, having a change in environment, and getting their lives back). Close to 10% of clients reported that changes in their physical health was a positive aspect.

FIGURE 8.4. PERCENT OF CLIENTS REPORTING POSITIVE ASPECTS OF THE PROGRAM (N = 21)



Aspects of treatment that clients identified as problematic or needing improvement are displayed in Figure 8.5. The negative aspects of the program suggest barriers that clients must overcome to participate in the program. Specifically, time away from work, household, or other responsibilities (33.3%), cost of the treatment (23.8%), problems with the facility (such as wait times, location, policies, dosage; 23.8%), and quality of counseling (e.g., not enough counseling; 19.0%). Less than 5% of clients reported transportation problems (4.8%).

FIGURE 8.5. PERCENT OF CLIENTS REPORTING NEGATIVE ASPECTS OF THE PROGRAM (N = 21)



### CLIENT INVOLVEMENT IN THE PROGRAM AND MANNER IN WHICH THE CLIENT LEFT

Clients who were not still involved in the clinic at follow-up were involved in the treatment program an average of 12.1 months (range of 1 to 48 months). Of those clients who were not currently involved in the opiate treatment program (n = 13), they were asked to report the main reason they were no longer using buprenorphine/naloxone or methadone. Less than half of clients (46.2%) reported there was no reason, and 23.1% of clients reported did not want to take the medication for their drug problem. Less than 10% of clients reported cost or insurance problems (7.7%), too many requirements from the clinic or doctor to stay on MAT (7.7%), and trouble staying with the medication schedule (7.7%).

Despite not being still involved in the clinic, 92.3% of clients reported they would refer a close friend or family member to the treatment provider though 66.7% of clients reported they would warn a friend or family member about certain things or who to work with. In addition, 23.1% (n = 3) reported having been in other treatment programs/episodes other than this program.

## SECTION 9. CONCLUSION AND IMPLICATIONS

*The KORTOS 2022 Annual Follow-Up Report describes characteristics of 192 clients who participated in opioid treatment programs during calendar year 2019 and completed intake interviews. In addition, outcomes are presented for 21 clients who completed a follow-up telephone interview 6 months after the intake interview and were still engaged in the treatment program.*

### WHO DO THE OPIOID TREATMENT PROGRAMS SERVE?

Of the clients with intake interviews (n = 192), they were an average of 39 years old ranging from 22 to 70 years old. Less than half (41.1%) were female and 58.3% were male. The majority of clients (60.9%) self-reported they decided get help on their own and 30.2% reported that they were referred to the OTP by a family member, partner, or friend. Less than half (47.9%) were unemployed, and of those unemployed clients, 38.0% reported they were looking for work.

Over three-quarters of KORTOS clients (76.0%) reported at least one adverse childhood experience before the age of 18. Specifically, about 41% reported 1-3 childhood experiences and 21.4% reported 4-6 childhood experiences. There were gender differences with adverse childhood experiences where women reported significantly more experiences. Significantly more men than women reported experiencing no adverse childhood experiences, whereas significantly more women than men reporting 7 - 9 types of adverse childhood experiences. Significantly more women (32.9%) than (17.0%) men reported physical maltreatment. About 44% of women and 19.6% of men reported experiencing emotional neglect, which was also significantly different. Significantly more women than men also reported sexual abuse (36.7% vs. 8.0%) before the age of 18. In terms of interpersonal victimization experiences, compared to men, significantly more women reported having ever been the victim of assault (other than IPV), being stalked by someone who scared them, sexually assaulted or raped, abused by a dating or intimate partner, and verbally, sexually or otherwise harassed in a way that made the client afraid for their safety. About 29% percent of women and men reported having ever been directly or indirectly threatened with a gun or held at gunpoint.

In the six months before entering the program, 47.9% of clients met study criteria for depression, and 59.4% met study criteria for generalized anxiety. About 10% reported suicidal thoughts or attempts of suicide in the 6 months before entering the program. In addition, 16.1% had post-traumatic stress disorder (PTSD) scores that indicated risk of PTSD.<sup>71</sup> About 46% of clients reported chronic pain in the 6 months before entering the program. Over half of clients (56.8%) reported they had at least one of the 15 chronic health problems listed on the intake interview. Trend analysis shows that from CY 2013 to CY 2019 the percent of clients who reported chronic medical problems has increased from just under half of clients to over half of clients.

Among the clients who were not incarcerated all 180 days before entering the program (n = 192), the majority reported using illegal drugs (98.4%) and smoking tobacco (79.7%) while 15.1% of clients reported using alcohol in the 6 months before intake. Close to 20% of clients reported only using opioids (19.8%) while 73.9% reported using opioids and at least one other class of drug.

<sup>71</sup>Among clients who were asked about PTSD symptoms.

Close to 13% of clients reported being arrested at least once and 12.5% reported being incarcerated at least one night in the 6 months before entering treatment. In addition, 9.4% of clients reported being under supervision by the criminal justice system.

## AREAS OF SUCCESS

The 2021 evaluation findings indicated that Kentucky opioid treatment programs have been successful in facilitating substantial positive changes in clients' lives. Results for those who were included in the followed-up analysis (n = 21) show that clients made substantial improvements from intake to follow-up in all four core components, including significant reductions in illegal drug and alcohol use, mental health problems, criminal justice system involvement, and a significant increase in quality of life. Improvements were also found for two supplemental areas: health status and economic and living circumstances.

### SUBSTANCE USE

Among clients who were not incarcerated all 180 days before entering the program or follow-up, there was a decrease in clients reporting past-6-month alcohol or illegal drug use with 100.0% of clients reporting any illegal drug use at intake compared to 61.9% at follow-up. Over half of clients (52.4%) reported misusing prescription opioids (such as morphine, Percocet, Oxycontin, Lortab) in the past 6 months at intake, whereas 4.8% of clients reported prescription opioid misuse in the past 6 months at follow-up. About three-quarters of followed-up clients (71.4%) also reported heroin use in the past 6 months at intake and that percent significantly decreased to 38.1% in the past 6 months at follow-up. The percent of clients using non-prescribed methadone and bup-nx also decreased. Not only did clients' use of overall opioids decrease significantly, but also their use of non-opioid drugs (such as marijuana, tranquilizers, benzodiazepines, and stimulants) decreased from 85.7% in the past 6 months at intake to 38.1% in the past 6 months at follow-up. In addition, the percent of clients with ASI drug composite scores that met the cut-off for severe substance use disorder (SUD) decreased from 95.2% in the past 30 days at intake to 19.0% in the past 30 days at follow-up.

### MEDICATION-ASSISTED TREATMENT

Two-thirds of clients who were not incarcerated all 180 days before treatment, reported at intake they had ever received medication from a clinic or doctor's office to help with their substance abuse. Of these clients (n =14), 64.3% reported it was dispensed in a clinic, 28.6% were prescribed the medication by a doctor in a specialty clinic, and 7.1% were prescribed the medication by a doctor in a general medical practice. The majority of clients (92.9%) reported that they think their use of medication-assisted treatment helped treat their drug problem.

The majority of clients, who were not incarcerated all 180 days before treatment entry or in the past 6 months at follow-up, (95.2%) reported that they received methadone in the past 6 months at follow-up. About 5% of clients reported receiving Suboxone, and none of the clients received Vivitrol. All of the clients reported that they think their use of medication-assisted treatment helped treat their drug problem.

### MENTAL HEALTH, PHYSICAL HEALTH, AND QUALITY OF LIFE

Clients' mental health also showed significant improvements. At follow-up, more than half as many clients had symptoms of depression, generalized anxiety, and comorbid

depression and anxiety in the past 6 months compared to intake. In addition, the percent of clients reporting past-6-month suicidal ideation or attempts decreased from 14.3% at intake to 0.0% at follow-up. The majority of clients at intake (81.0%) reported using alcohol, prescription drugs, or illegal drugs to reduce stress, anxiety, worry, sadness, or fear. At follow-up, 23.8% of clients reported using alcohol, prescription drugs, or illegal drugs to reduce stress, anxiety, worry, sadness, or fear, which was a significant decrease of 57.2%.

Clients' physical health was also better at follow-up. Clients rated their health, on average, as 2.9 at intake and this significantly increased to 3.2 at follow-up. In addition, the number of days clients reported their physical health was not good decreased significantly from an average of 5.4 days to 1.7 days. The number of days clients' mental health was not good also decreased significantly from intake (13.0) to follow-up (4.3). Clients reported significantly fewer days their poor physical or mental health kept them from doing their usual activities at intake (8.6 days) to follow-up (3.5 days). The percent of clients who reported chronic pain that was persistent and lasted at least 3 months decreased significantly from 47.6% at intake to 19.0% at follow-up as well.

On a scale from 1 to 10, clients rated their quality of life as significantly higher after participating in the program (8.3 at follow-up compared to 4.7 at intake).

### CRIMINAL JUSTICE SYSTEM INVOLVEMENT AND INTERPERSONAL VICTIMIZATION

A minority of KORTOS clients were involved in the criminal justice system in the past 6 months at intake and the percent of clients who reported being arrested or incarcerated was significantly lower at follow-up. Specifically, 4.8% of clients reported having been arrested in the 6 months before entering treatment and, at follow-up, 9.5% of clients reported an arrest in the past 6 months. Almost 10% of clients reported spending at least one night in jail or prison in the past 6 months at intake compared to 4.8% of clients at follow-up. Past-8-year trend analysis shows that the percent of clients reporting an arrest and clients reporting they spent at least one night in jail have been relatively stable.

Clients also had decreases in past-6-month crime and interpersonal victimization with fewer clients reporting being verbally harassed in public, burglarized, robbed or mugged, assaulted, or being a victim of intimate partner abuse at follow-up.

### ECONOMIC STATUS AND LIVING CIRCUMSTANCES

KORTOS clients showed improvements in past-6-month economic and living circumstances from intake to follow-up. Specifically, there was a significant increase in the percent of clients reporting attending/completing vocational school, college, or graduate school from 42.9% at intake to 76.2% at follow-up. About 43% of clients reported being employed full-time at follow-up compared to 38.1% at intake, which was not a significant increase. In addition, the percent of clients reporting difficulty meeting health care needs (e.g., doctor visits, dental visits, and prescription medications) for financial reasons in the past 6 months decreased (but not significantly) from 47.6% at intake to 28.6% at follow-up.

### RECOVERY SUPPORTS

At follow-up, clients reported having significantly more people they could count on for recovery support (8.0 vs. 3.8 at intake). The majority of clients (90.5%) stated they thought they had a moderately or very good chance of staying off drugs or alcohol at follow-up. At both intake and follow-up, clients reported that employment, the need to stay out of

jail, and support from friends would be most useful in helping them quit or stay off drugs/ alcohol.

## MULTIDIMENSIONAL RECOVERY

Recovery goes beyond relapse or return to occasional drug or alcohol use. The multidimensional recovery measure items from the intake and follow-up surveys to create one measure of recovery. At intake, none of the clients had all positive dimensions of recovery, whereas at follow-up, close to half (47.6%) had all positive dimensions.

## SATISFACTION WITH OPIOID TREATMENT PROGRAM

The majority of clients reported that the program started good (76.2%) and 95.2% reported it was currently good. In addition, all clients reported that the treatment episode was working pretty well or extremely well for them. Furthermore, the majority of clients (95.2%) indicated they would refer a close friend or family member to their treatment provider. Of the clients who reported they would refer a close friend or family member to the program (n = 20), 15.0% reported they would warn their friend or family member about certain things or tell them who to work with or who to avoid.

On a scale from 1 representing the worst possible experience to 10 representing the best possible experience, clients rated their experience an 9.1 with 95.3% of clients giving a highly positive rating of 8 through 10. In addition, all clients felt they had input into their treatment goals, plans, and progress. The majority of clients reported that their expectations and hopes for treatment and recovery were met, they worked on and talked about things that were most important to the client, and that treatment approach and method was a good fit. Clients reported many positive aspects of their participation in the program including reduced substance use, improved mental health and their feelings about themselves, improved relationships with others, the quality of the treatment, and improved financial situation.

## AREAS OF CONCERN

While there were many positive outcomes overall, there are also potential opportunities to make even more significant improvements in clients' functioning after they begin treatment.

## ILLEGAL DRUG USE

When looking at trends over time in past-30-day use at intake, results show that while prescription opioid and methadone use has decreased gradually over the past 14 years, heroin use has sharply increased since CY 2011. Beginning in CY 2018, heroin use decreased briefly but it still remains high relative to other past-30-day substance use. Compared to heroin and prescription opioids, methamphetamine use is relatively low; however, use has increased in the past couple of years from 13.7% in CY 2017 to 40.7% in CY 2020.

Also, almost two-thirds of KORTOS clients (61.9%) reported using illegal drugs in the 6 months before follow-up. While this year's follow-up sample size is small and may be affecting the results, this is still a sizable increase in illegal drug use at follow-up compared to last year (34.1%). Specifically, 38.1% of clients reported using heroin at follow-up and 38.1% of clients reported non-opioid drug use at follow-up. White et al. found that screening positive for just one non-prescribed drug doubled a client's dropout rate and screening for multiple drugs quadrupled it. In addition, continued drug use during medication-assisted



treatment has been associated with early program termination<sup>72, 73</sup> and longer treatment retention has been associated with more positive outcomes.<sup>74, 75</sup>

In addition, although the percent of clients who met DSM-5 criteria for severe SUD decreased at follow-up, 19.0% still met criteria for severe substance use disorder for drug use and 19.0% of clients still had ASI drug composite scores that met the cut-off for severe substance use disorder. While the percent of clients who reported substance use decreased from intake to follow-up, 15.0% of clients still reported experiencing problems associated with drugs and alcohol including cravings, withdrawal, wanting to quit but being unable, or worrying about relapse at follow-up. About 5% of clients reported that they were considerably or extremely troubled or bothered by drug or alcohol problems at follow-up.

## SMOKING RATES

Rates tobacco smoking were high for clients (71.4%) and did not change from the past 6 months at intake to the past 6 months at follow-up. Tobacco use is associated with increased mortality, and smoking cessation has been associated with lower alcohol and drug relapse.<sup>76</sup> Smoking has been associated with increased mental health symptoms and physical health problems.<sup>77, 78</sup> There is a commonly held belief that individuals should not attempt to quit smoking while in substance abuse treatment, because smoking cessation can endanger their sobriety. This belief has been refuted by recent empirical research studies.<sup>79</sup> Voluntary smoking cessation during substance abuse treatment has been associated with lower relapse.

In addition, almost 24% clients reported the use of e-cigarettes at follow-up. In fact, trend analysis shows that the percent of clients reporting e-cigarette use has generally increased over time at follow-up. While e-cigarettes are widely believed to be a mechanism for smoking cessation and/or a less dangerous alternative to conventional cigarettes, they also carry their own health risks. Although e-cigarettes contain lower levels of carcinogens compared to regular tobacco cigarettes,<sup>80</sup> the e-cigarette still contains potent cancer-

<sup>72</sup>Davstad, I., Stenbacka, M., Leifman, A., Beck, O., Kormaz, S., & Romelsjö, A. (2007) Patterns of illicit drug use and retention in a methadone program: A longitudinal study. *Journal of Opioid Maintenance* 3(1), 27-34.

<sup>73</sup>White, W., Campbell, M., Spencer, R., Hoffman, H., Crissman, B., & DuPont, R. (2014). Patterns of abstinence or continued drug use among methadone maintenance patients and their relation to treatment retention. *Journal of Psychoactive Drugs*, 46(2), 114-122.

<sup>74</sup>Hubbard, R., Craddock, S., & Anderson, J. (2003). Overview of 5-year follow-up outcomes in the drug abuse treatment outcome studies (DATOS). *Journal of Substance Abuse Treatment*, 25, 125-134.

<sup>75</sup>Gibson, A., Degenhardt, L., Mattick, R., Ali, R., White, J., & O'Brien, S. (2008). Exposure to opioid maintenance treatment reduces long-term mortality. *Addiction*, 103, 462-468.

<sup>76</sup>Proschaska, J. (2010). Failure to treat tobacco use in mental health and addiction treatment settings: A form of harm reduction? *Drug and Alcohol Dependence*, 110, 177-182.

<sup>77</sup>Patton, G., Coffey, C., Carlin, J., Sawyer, S., & Wakefield, M. (2006). The course of early smoking: A population-based cohort study over three years. *Addiction*, 93, 1251-1260.

<sup>78</sup>Kalman, D., Morissette, S., & George, T. (2005). Co-morbidity of smoking in patients with psychiatric and substance use disorders. *American Journal of Addictions*, 14(2), 106-123.

<sup>79</sup>Baca, C., & Yahne, C. (2009). Smoking cessation during substance abuse treatment: What you need to know. *Journal of Substance Abuse Treatment*, 36, 205-219.

<sup>80</sup>Goniewicz ML, Knysak J, Gawron M, Kosmider L, Sobczak A, et al. (2014). Levels of selected carcinogens and toxicants in vapour from electronic cigarettes. *Tobacco Control*, 23(2), 133-39.

causing toxins as well as chemicals that can trigger cardiovascular and lung disease,<sup>81, 82</sup> and gene mutation.<sup>83</sup>

## MENTAL AND PHYSICAL HEALTH

While there were significant reductions in clients meeting study criteria for generalized anxiety, the 19.0% of clients who reported generalized anxiety at follow-up still reported an average of 6.5 (out of 7) symptoms in the past 6 months at follow-up. In addition, almost one-quarter of clients (23.8%) of clients reported using alcohol, prescription drugs, or illegal drugs to reduce stress, anxiety, worry, sadness, or fear in the past 6 months at follow-up. Further, 38.1% of clients were worried about their personal safety at follow-up. Also, while the number of clients reporting chronic pain decreased significantly from intake (47.6%) to follow-up (19.0%), clients still reported 26.3 average days of chronic pain at follow-up.

## ECONOMIC HARDSHIP

Meeting basic needs including health, stable living arrangements, having a purpose with daily meaningful activities, and recovery community are the four key dimensions to recovery.<sup>84</sup> While the percent of clients who reported having difficulty meeting needs for financial reasons decreased at follow-up, almost half of clients (47.6%) at follow-up still reported having difficulty meeting basic living needs and 28.6% still reported difficulty meeting health care needs in the past 6 months. Similarly, almost half of clients (47.6%) remained unemployed at follow-up. Trends in economic difficulties show that the number of clients who reported they had difficulty meeting basic living needs and/or health care needs has increased at follow-up since 2017. Further, clients reported working an average of only 3.1 months in the past 6 months at follow-up. The resulting financial strain from these economic factors could lead to increased substance use to alleviate the stress.<sup>85</sup> Providing referrals and support for these factors may help improve basic living situations for many clients and support continued recovery living for long-term positive results after treatment.

## GENDER DIFFERENCES ON TARGETED FACTORS

There were only a few gender differences in targeted factors. Significantly more men reported heroin use in the past 30 days at follow-up and smokeless tobacco in the past 6 months at intake and in the past 30 days at intake.

## COMPARISON OF CLIENTS WHO WERE NOT LONGER INVOLVED IN THE TREATMENT CLINIC AT FOLLOW-UP WITH CLIENTS WHO WERE STILL INVOLVED

There were a few significant demographic differences between clients who were still involved with the clinic at follow-up (n=21) and clients who were not (n=13). Clients who were involved with the clinic at follow-up were all White and clients who were no longer involved in the clinic at follow-up were more likely to be married. Clients who were not

<sup>81</sup> Drummond, M. B., & Upson, D. (2014). Electronic cigarettes. Potential harms and benefits. *Annals of the American Thoracic Society*, 11(2), 236-242.

<sup>82</sup> Glantz, S. A., & Bareham, D. W. (2018). E-cigarettes: use, effects on smoking, risks, and policy implications. *Annual Review Of Public Health*, 39, 215-235.

<sup>83</sup> Canistro, D., Vivarelli, F., Cirillo, S., Marquillas, C. B., Buschini, A., Lazzaretti, M., et al. (2017). E-cigarettes induce toxicological effects that can raise the cancer risk. *Scientific reports*, 7(1), 1-9.

<sup>84</sup> <https://www.samhsa.gov/brss-tacs/recovery-support-tools-resources>

<sup>85</sup> Shaw, B. A., Agahi, N., & Krause, N. (2011). Are Changes in Financial Strain Associated with Changes in Alcohol Use and Smoking Among Older Adults? *Journal of Studies on Alcohol and Drugs*, 72(6), 917-925.

still involved in the treatment program reported significantly more years of education (15.6) compared to clients who were still involved in the clinic at follow-up (13.9). Overall, significantly more clients who involved in the clinic at follow-up reported illicit drug use at follow-up, specifically, more clients who were still involved in the clinic at follow-up reported stimulants or cocaine use in the 6 months before follow-up (19.0% vs. 0.0%). Significantly more clients who were still involved in the clinic at follow-up reported e-cigarette use in the past 6 months at follow-up compared to clients who were no longer involved in the clinic. About 15% of clients who were not involved in the clinic at follow-up and 23.8% of clients who were still involved in the clinic at follow-up met or surpassed the Addiction Severity Index (ASI) composite score cutoff for alcohol and/or drug severe SUD at follow-up with no difference by group. There were significant differences between clients who were still involved in the clinic and clients who were not involved in the clinic at follow-up for the percent of clients attending mutual help recovery meetings in the past 30 days at follow-up (14.3% vs. 61.5%, respectively).

## STUDY LIMITATIONS

The study findings must be considered within the context of the study's limitations. First, for this report year compared to previous years, a considerably smaller number of clients completed a follow-up and an intake because of the COVID-19 pandemic. A smaller number of clients at intake means that there is a smaller pool of eligible follow-up sample clients to pull from. In addition to fewer clients completing an intake assessment, the percent of baseline clients not agreeing to be contacted for follow-up has increased. In 2022, 62.0% of the intake sample did not agree to be contacted compared to only 33.0% in 2019.

TABLE C.1. NUMBER AND PERCENT OF CLIENTS CONSENTING TO FOLLOW-UP FOR EACH REPORT YEAR

Report year	Intake sample size	Clients NOT consenting to follow-up
2018	514	247 (48.1%)
2019	625	206 (33.0%)
2020	384	174 (45.3%)
2021	233	145 (62.2%)
2022	192	119 (62.0%)

Second, because there is no appropriate group of opioid dependent individuals who would like treatment but do not receive it to compare with the KORTOS individuals who participate in treatment, all changes from intake to follow-up cannot be attributed to medication-assisted treatment.

Third, because not all clients agree to participate in the 6-month follow-up interview, it is unclear how generalizable the findings are to the entire client population that completes an intake interview. Analysis comparing those individuals who completed a follow-up interview with those who did not complete a follow-up interview for any reason (for example, they did not agree to be in the follow-up study, they were not selected into the follow-up sample, or they were not successfully contacted for the follow-up interview) found a few significant differences between the two groups.

Fourth, in recent years, program intake data collection at the OTPs has decreased, which

limits the number of clients the follow-up research team can target for follow-up data collection. Decreasing numbers of participants in KORTOS reduces the generalizability of the outcome findings and the statistical power to detect the effects of the program on outcomes. Increasing efforts to conduct the intake surveys with OTP clients will net more data, more robust and meaningful findings, which help to support the mission of the programs.

Fifth, data included in this report was self-reported by clients. There is reason to question the validity and reliability of self-reported data, particularly with regard to sensitive topics, such as illegal behavior and stigmatizing issues such as mental health and substance use. However, some research has supported findings about the reliability and accuracy of individuals' reports of their substance use.<sup>86, 87, 88</sup> For example, in many studies that have compared agreement between self-report and urinalysis the concordance or agreement is acceptable to high.<sup>89, 90, 91</sup> In fact, in some studies, when there were discrepant results between self-report and urinalysis of drugs and alcohol, the majority were self-reported substance use that was not detected with the biochemical measures.<sup>92, 93, 94</sup> In other studies, higher percentages of underreporting have been found.<sup>95</sup> Prevalence of underreporting of substance use is quite varied in studies. Nonetheless, research has found that certain conditions facilitate the accuracy of self-report data such as assurances of confidentiality and memory prompts.<sup>96</sup> Moreover, the “gold standard” of biochemical measures of substance use have many limitations: short windows of detection that vary by substance; detection varies on many factors such as the amount of the substance consumed, chronicity of use, sensitivity of the analytic method used.<sup>97</sup> Therefore, the study method includes

<sup>86</sup> Del Boca, F.K., & Noll, J.A. (2000). Truth or consequences: The validity of self-report data in health services research on addictions. *Addiction*, 95, 347-360.

<sup>87</sup> Harrison, L. D., Martin, S. S., Enev, T., & Harrington, D. (2007). *Comparing drug testing and self-report of drug use among youths and young adults in the general population* (DHHS Publication No. SMA 07-4249, Methodology Series M-7). Rockville, MD: Substance abuse and Mental Health Services Administration, Office of Applied Studies.

<sup>88</sup> Rutherford, M.J., Cacciola, J.S., Alterman, A.I., McKay, J.R., & Cook, T.G. (2000). Contrasts between admitters and deniers of drug use. *Journal of Substance Abuse Treatment*, 18, 343-348.

<sup>89</sup> Rowe, C., Vittinghoff, E., Colfax, G., Coffin, P. O., & Santos, G. M. (2018). Correlates of validity of self-reported methamphetamine use among a sample of dependent adults. *Substance Use & Misuse*, 53(10), 1742-1755.

<sup>90</sup> Rygaard Hjorthoj, C., Rygaard Hjorthoj, A., & Nordentoft, M. (2012). Validity of timeline follow-back for self-reported use of cannabis and other illicit substances—Systematic review and meta-analysis. *Addictive Behaviors*, 37, 225-233.

<sup>91</sup> Wilcox, C. E., Bogenschutz, M. P., Nakazawa, M., & Woody, G. (2013). Concordance between self-report and urine drug screen data in adolescent opioid dependent clinical trial participants. *Addictive Behaviors*, 38, 2568-2574.

<sup>92</sup> Denis, C., Fatséas, M., Beltran, V., Bonnet, C., Picard, S., Combourieu, I., Daulouède, J., & Auriacombe, M. (2012). Validity of the self-reported drug use section of the Addiction Severity and associated factors used under naturalistic conditions. *Substance Use & Misuse*, 47, 356-363.

<sup>93</sup> Hilario, E. Y., Griffin, M. L., McHugh, R. K., McDermott, K. A., Connery, H. S., Fitzmaurice, G. M., & Weiss, R. D. (2015). Denial of urinalysis-confirmed opioid use in prescription opioid dependence. *Journal of Substance Abuse Treatment*, 48, 85-90.

<sup>94</sup> Williams, R. J., & Nowatzki, N. (2005). Validity of self-report of substance use. *Substance Use & Misuse*, 40, 299-313.

<sup>95</sup> Chermack, S. T., Roll, J., Reilly, M., Davis, L., Kilaru, U., Grabowski, J. (2000). Comparison of patient self-reports and urinalysis results obtained under naturalistic methadone treatment conditions. *Drug and Alcohol Dependence*, 59, 43-49.

<sup>96</sup> Del Boca, F. K., & Noll, J. A. (2000). Truth or consequences: the validity of self-report data in health services research on addictions. *Addiction*, 95 (Suppl. 3), S347–S360.

<sup>97</sup> Williams, R. J., & Nowatzki, N. (2005). Validity of self-report of substance use. *Substance Use & Misuse*, 40, 299-313.

several key strategies to facilitate accurate reporting of sensitive behaviors at follow-up including: (a) the follow-up interviews are conducted by telephone with a University of Kentucky Center on Drug and Alcohol Research (UK CDAR) staff person who is not associated with any opioid replacement treatment program; (b) the follow-up responses are confidential and are reported at a group level, meaning no individual responses are linked to participants' identity; (c) the study procedures, including data protections, are consistent with federal regulations and approved by the University of Kentucky Human Subjects Institutional Review Board; (d) confidentiality is protected under Federal law through a Federal Certificate of Confidentiality; (e) participants can skip any question they do not want to answer; and (f) UK CDAR staff are trained to facilitate accurate reporting of behaviors and are regularly supervised for quality data collection and adherence to confidentiality.

## CONCLUSION

The 2022 KORTOS evaluation indicates that opioid treatment programs in Kentucky have been successful in facilitating positive changes in clients' lives in a variety of ways, including decreased substance use, decreased severity of substance use, decreased mental health symptoms, and decreased economic hardship. Results also show that clients appreciate and value their experiences in treatment programs and have an improved quality of life after beginning participation in treatment. Overall, KORTOS clients had significant improvements in their lives that have been shown to be key factors that facilitate recovery<sup>98</sup>: meeting basic needs, stable living arrangements, having a purpose with daily meaningful activities, and recovery community. However, there were some areas of concern related to drug use, smoking rates, mental health, financial difficulties, and gender differences at follow-up.

---

<sup>98</sup><http://blog.samhsa.gov/2012/03/23/definition-of-recovery-updated/>

## APPENDIX A. METHODS

The KORTOS intake and follow-up interview instruments are modeled after the Kentucky Treatment Outcome Study (KTOS) assessment and are based on theory and research about substance use-related comorbidities relevant to substance use among clients in opioid treatment programs. The assessment's four core components (e.g., substance use, mental health, criminal justice involvement, and quality of life) and three supplemental components (e.g., health status, economic status and living circumstances, and recovery supports) have demonstrated validity and reliability<sup>99</sup> and have been developed in collaboration with key stakeholders to consider the context of Kentucky opioid treatment programs.

KORTOS intake interviews were conducted by a clinician or staff person at the OTP using a web-based interview tool, in which identifying data were encrypted and submitted to the master database on the UK CDAR secure server. At the end of the intake interview, clinicians explained the follow-up study to clients and gave them the opportunity to volunteer to participate. Clients who were interested gave electronic consent to be contacted by UK CDAR BHOS staff members about 6 months later. Follow-up interviews were conducted via telephone using a questionnaire with items and questions similar to the questions in the intake interview.

The target month for the follow-up interview was 6 months after the intake interview was completed. In other words, if a client completed an intake interview in December 2020, the target month for the follow-up interview was June 2021. The window for completing a follow-up interview with an individual selected into the follow-up sample began one month before the target month and spanned until two months after. Therefore, if the target month for a follow-up was June 2021, interviewers began working to locate and contact the individual in May and could work the file until the end of August.

Of the 192 clients who completed an intake interview in 2020, only 73 (38.0%) agreed to be contacted for the follow-up study, which is a lower agreement rate than in previous years. From this group of clients who voluntarily agreed to be contacted for the follow-up study, the research team pulled the follow-up sample by first identifying clients who had provided the minimum amount of contact information (e.g., two phone numbers or one phone number and one mailing address) and whose intakes were submitted to CDAR less than 30 days after the intake was completed, which left a sample of 63 clients. Of these clients, 34 completed a follow-up interview (see Table AA.1); thus, the follow-up rate was 54.0%.

The remaining clients were never successfully contacted, or if contacted they never completed the follow-up interview (n = 29, or 46.0% of the cases eligible for follow-up). Compared to previous reports, the number of individuals in the expired category has increased. First, the quality of contact information collected at the time of intake interviews has worsened over time. Second, the percent of individuals who ever answer the follow-up interviewers' phone calls is decreasing. The volume of scam-related or robocalls increased 35% in 2019, accounting for over one-third of personal calls in the U.S.<sup>100</sup> Both of these factors have necessitated changes in procedures and strategies, which the follow-up study team has developed and put into place. The only cases not considered accounted for are those individuals who are classified as expired.

<sup>99</sup>Logan, TK, Cole, J., Miller, J., Scrivner, A., & Walker, R. (2020). *Evidence Base for the Kentucky Opioid Replacement Treatment Outcome Study (KORTOS) Assessment and Methods*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

<sup>100</sup><https://www.usatoday.com/story/tech/news/2019/12/04/robocalls-us-eighth-most-spammed-country-report/2613528001/>

TABLE AA.1. FINAL CASE OUTCOMES FOR FOLLOW-UP EFFORTS (N = 63)

	Number of Records	Percent
Ineligible for follow-up interview .....	0	0.0%
	<b>Number of cases eligible for follow- up (n = 63)</b>	
Completed follow-up interviews .....	34	
Follow-up rate is calculated by dividing the number of completed interviews by the number of eligible cases and multiplying by 100 .....		54.0%
Expired cases (i.e., never contacted, did not complete the interview during the follow-up period) .....	29	
Expired rate ((the number of expired cases/eligible cases)*100) ..		46.0%
Refusal.....	0	
Refusal rate ((the number of refusal cases/eligible cases)*100).....		0.0%
Cases accounted for (i.e., records ineligible for follow-up + completed interviews + refusals).....	34	
Percent of cases accounted for ((# of cases accounted for/total number of records in the follow-up sample)*100).....		54.0%

Appendix B provides detailed information on the locating efforts for the 2015 KORTOS follow-up sample.

Appendix C presents analysis on comparisons between clients who completed a follow-up interview and clients who did not complete a follow-up interview for any reason on key variables included in the intake interview.

## APPENDIX B. LOCATING EFFORTS FOR THE 2015 KORTOS FOLLOW-UP STUDY

Project interviewers documented their efforts (e.g., mailings, phone calls, Internet searches, etc.) to locate each participant included in the sample of individuals to be followed up from July 2013 to June 2014 (n = 350), which is the follow-up period corresponding to the KORTOS 2015 report. All the locator files were examined and used to extract information about the efforts project interviewers made to locate and contact participants as well as the type of contact information provided by participants in the original locator information when the intake interview data was submitted to UK CDAR. A subsample of records was randomly selected and independently examined to check that the procedures for extracting data were reliable and valid. The extraction sheets were compared between the two raters for interrater reliability, which was high (96.1%). The following information is based on the data collected during this review of locator files.

For all 350 records, a total of 2,182 phone calls were made to client phone numbers and 773 calls to contact persons' phone numbers. As Table AB.1 shows, project interviewers made an average of about 6.2 calls to client phone numbers and 2.4 calls to contact persons' phone numbers. Fewer than 40% of clients called in at any point and only 4.3% called-in to complete the interview after receiving the initial mailing without project interviewers putting additional effort into contacting the clients. That means 95.7% of clients took considerable effort to try to locate, contact, and complete follow-up interviews.

A total of 649 mailings were sent to client addresses and 26 mailings were sent to contact persons, an average of 1.9 mailings to clients and 0.1 mailings to contact persons. The research team received returned mail for 12.3% of clients that received mailings to client addresses and 1.4% of clients that received mailings to contact addresses.



In cases where the client contact information was incorrect (i.e., mail was returned, phone number was disconnected), online public directory databases were used to try to verify that we had correct or updated information for the client. Because it had been six months since they provided contact information, we would like to be sure we are not calling or sending mailings to someone other than the client. Therefore, verifying the correct contact information is a critical interim step in the follow-up process to protect confidentiality. For 92.3% of the clients, the interviewers used public searches/directories to verify contact information. If the client information could not be verified, interviewers also used social media and more detailed public directory databases to find updated contact information (52.9%). In cases where very little contact information was given or clients were not successfully located in the ways listed above, more in-depth searching methods were used (28.9%). As a last resort, in the few cases where the client was not successfully located in any of the ways described above, interviewers worked to reach client contacts provided by them at intake (6.7%).



### KORTOS 2015 Quality of Data and Locator Efforts

For the 2015 follow-up study, 350 participants were included in the sample of individuals to be followed up from July 2013 to June 2014. Efforts to locate and contact these participants were examined.

Of these clients, 223 completed a follow-up survey for a follow-up rate of 82.6%.



# APPENDIX C. CLIENT CHARACTERISTICS AT INTAKE FOR THOSE WHO COMPLETED A FOLLOW-UP INTERVIEW AND THOSE WHO DID NOT COMPLETE A FOLLOW-UP INTERVIEW

*Clients who completed a follow-up interview are compared in this section with clients who did not complete a follow-up interview for any reason (e.g., did not agree to be contacted for the follow-up interview, not selected into the follow-up sample, ineligible for follow-up, not successfully located for the follow-up).*

## DEMOGRAPHICS

There were very few significant differences between clients who did and did not complete a follow-up on demographics (see Table AC.1). Significantly more clients who completed a follow-up interview were female (58.8%) compared to clients who did not complete a follow-up interview (37.3%).

TABLE AC.1. COMPARISON OF DEMOGRAPHICS FOR CLIENTS WHO WERE FOLLOWED UP AND CLIENTS WHO WERE NOT FOLLOWED UP

	FOLLOWED UP	
	NO n = 158	YES n = 34
<b>Age</b> <sup>101</sup> .....	38.9 years	39.8 years
<b>Gender*</b>		
Male .....	62.7%	41.2%
Female.....	37.3%	58.8%
<b>Race</b>		
White .....	91.1%	94.1%
African American .....	5.7%	5.9%
Other or Multiracial .....	3.2%	0.0%
<b>Marital status</b>		
Never married.....	22.2%	26.5%
Married.....	27.2%	26.5%
Separated or divorced	17.7%	14.7%
Widowed .....	3.2%	2.9%
Cohabiting .....	29.7%	29.4%

\*p < .05.

<sup>101</sup>7 clients had incorrect birthdates and, therefore, age could not be determined.

## SUBSTANCE USE AT INTAKE

Use of illegal drugs in the 6 months before entering treatment is presented by follow-up status in Table AC.2. The most frequently reported illegal drugs used in the 6 months before entering treatment were heroin, prescription opioids/opioids, and marijuana. There were no significant difference between clients who completed a follow-up and clients who did not complete a follow-up interview for illegal drug use in the 6 months before treatment.

TABLE AC.2. PERCENT OF CLIENTS REPORTING ILLEGAL DRUG USE IN THE 6 MONTHS BEFORE ENTERING TREATMENT

	FOLLOWED UP	
	NO n = 158	YES n = 34
Any illegal drug.....	98.7%	97.1%
Prescription opioid/opiate (illegal use).....	65.2%	50.0%
Heroin .....	71.5%	76.5%
Marijuana.....	51.3%	58.8%
CNS depressants.....	20.3%	23.5%
Cocaine .....	18.4%	20.6%
Stimulants (amphetamines, methamphetamine, prescription stimulants).....	43.7%	38.2%
Non-prescribed bup-nx.....	17.7%	17.6%
Non-prescribed methadone.....	9.5%	11.8%
Other illicit drugs (hallucinogens, inhalants, synthetic drugs) .....	6.3%	14.7%

There were no significant differences between clients who were followed up and clients who were not followed up for alcohol use (see Table AC.3).

TABLE AC.3. PERCENT OF CLIENTS REPORTING ALCOHOL USE IN THE 6 MONTHS BEFORE ENTERING TREATMENT

	FOLLOWED UP	
	NO n = 158	YES n = 34
Alcohol.....	13.3%	23.5%
Alcohol to intoxication.....	7.6%	11.8%
Binge drank alcohol (i.e., drank 5 or more [4 or more for women] drinks in 2 hours).....	4.4%	8.8%

In the 6 months before entering the program, the majority of clients reported smoking tobacco products, with no difference between those who completed a follow-up interview and those who did not (see Table AC.4). There were also no differences between the two groups for the use of smokeless tobacco or e-cigarettes.

TABLE AC.4. PERCENT OF CLIENTS REPORTING TOBACCO USE IN THE 6 MONTHS BEFORE ENTERING TREATMENT

	FOLLOWED UP	
	NO n = 158	YES n = 34
Smoked tobacco .....	81.0%	73.5%
Smokeless tobacco.....	7.6%	8.8%
E-cigarettes.....	11.4%	20.6%

Self-reported severity of alcohol and drug use was measured with Addiction Severity Index (ASI) alcohol and drug composite scores. Alcohol and drug composite scores are presented in Table AC.5 for those clients who were not in a controlled environment all 30 days before entering treatment.<sup>102</sup> The highest composite score is 1.0 for each of the two substance categories.

The majority of clients who were not in a controlled environment all 30 days met or surpassed the Addiction Severity Index (ASI) composite score cutoff for alcohol and/or drug severe SUD with no difference by follow-up status. The average score for the drug severity composite score was 0.36 for clients who did not complete a follow-up interview and 0.33 for clients who did complete a follow-up (see Table AC.5).

TABLE AC.5. SUBSTANCE ABUSE AND DEPENDENCE PROBLEMS AT INTAKE

Recent substance use problems among clients who were....	Not in a controlled environment all 30 days before entering treatment	
	NO n = 158	YES n = 34
Percentage of clients with ASI composite score equal to or greater than cutoff score for		
Severe alcohol or drug use disorder.....	94.9%	94.1%
Severe alcohol use disorder .....	7.7%	17.6%
Severe drug use disorder .....	94.9%	94.1%
Average composite score for alcohol use <sup>a</sup> .....	.04	.06
Average composite score for drug use <sup>b</sup> .....	.36	.33

a Score equal to or greater than .17 is indicative of severe alcohol use disorder.

b Score equal to or greater than .16 is indicative of severe drug use disorder.

There were no significant differences between the groups for having ever been in substance abuse treatment (see Table AC.6). Among clients who reported a history of substance abuse treatment, the average number of lifetime treatment episodes was 3.4 for those who did not complete a follow-up and 3.5 for those who did.

<sup>102</sup> Clients who were in a controlled environment all 30 days before intake were not included in this analysis because being in a controlled environment limits one’s access to substances.

TABLE AC.6. HISTORY OF SUBSTANCE ABUSE TREATMENT IN LIFETIME

	FOLLOWED UP	
	NO n = 158	YES n = 34
Ever been in substance abuse treatment in lifetime .....	73.4%	61.8%
Among those who had ever been in substance abuse treatment in lifetime,	(n = 116)	(n = 21)
Mean number of times in treatment .....	3.4	3.5

## MENTAL HEALTH AT INTAKE

The mental health questions included in the KORTOS intake and follow-up interviews are not clinical measures, but instead are research measures. A total of 9 questions were asked to determine if they met study criteria for depression, including at least one of the two leading questions: (1) “Did you have a two-week period when you were consistently depressed or down, most of the day, nearly every day?” and (2) “Did you have a two-week period when you were much less interested in most things or much less able to enjoy the things you used to enjoy most of the time?”. There was not a significant difference between clients who completed a follow-up interview and clients who did not complete a follow-up interview for the percent of clients who met criteria for depression: 58.8% vs. 45.6% (see Table AC.7).

A total of 7 questions were asked to determine if clients met study criteria for generalized anxiety, including the leading question: “In the 6 months before entering this program, did you worry excessively or were you anxious about multiple things on more days than not for all 6 months (like family, health, finances, school, or work difficulties)?” There was not a significant difference between clients who completed a follow-up interview and clients who did not complete a follow-up interview for the percent of clients who met criteria for generalized anxiety: 70.6% vs. 57.0%.

Two questions were included in the intake interview that asked about thoughts of suicide and attempted suicide in the 6 months before clients entered treatment. There were no significant differences between clients who were followed up and not followed up for thoughts of suicide or suicide attempts.

TABLE AC.7. PERCENT OF CLIENTS REPORTING MENTAL HEALTH PROBLEMS IN THE 6 MONTHS BEFORE ENTERING THE PROGRAM

	FOLLOWED UP	
	NO n = 158	YES n = 34
Depression .....	45.6%	58.8%
Generalized anxiety .....	57.0%	70.6%
Suicidality (e.g., thoughts of suicide or suicide attempts) .....	9.5%	14.7%

## PHYSICAL HEALTH AT INTAKE

To give an idea of the physical health of clients when they entered treatment, Table AC.8 presents the percent of clients that reported health problems at intake. There was not a significant difference between clients who completed a follow-up reported experiencing chronic pain (i.e., pain that lasted more than 3 months) at intake compared to clients who were and were not followed up (44.1% vs. 46.8%, respectively). Clients were asked at intake if a doctor had ever told them they had any of the 16 chronic medical problems listed (e.g., asthma, arthritis, cardiovascular disease, diabetes, chronic obstructive pulmonary disease [COPD], tuberculosis, severe dental disease, cancer, Hepatitis B, Hepatitis C, HIV, and other sexually transmitted diseases). The most commonly reported chronic medical problems are presented in Table AC.8: Hepatitis C, arthritis, severe dental disease, cardiovascular disease, and asthma.

TABLE AC.8. PHYSICAL HEALTH STATUS AT INTAKE

	FOLLOWED UP	
	NO n = 158	YES n = 34
Chronic pain (lasting at least 3 months).....	46.8%	44.1%
Ever told by a doctor that client had one of the 12 chronic medical problems listed.....	53.8%	70.6%
Hepatitis C.....	22.8%	20.6%
Arthritis.....	13.9%	23.5%
Severe dental disease.....	11.4%	17.6%
Cardiovascular disease.....	12.7%	14.7%
Asthma.....	12.7%	2.9%

## SOCIOECONOMIC INDICATORS

There were no significant differences between the groups on clients' level of education when entering treatment. Less than one-quarter of clients reported less than a high school diploma or GED at intake. Around 44% of clients who were not followed up and 35.3% of clients who were followed up reported having a GED or high school diploma. Thirty-eight percent of clients who did not complete a follow-up and 52.9% of clients who did complete a follow-up attended vocational school or higher.

TABLE AC.9. CLIENTS' HIGHEST LEVEL OF EDUCATION COMPLETED AT INTAKE

	FOLLOWED UP	
	NO n = 158	YES n = 34
<b>Highest level of education completed</b>		
Less than GED or high school diploma.....	17.7%	11.8%
GED or high school diploma.....	44.3%	35.3%
Vocational school to graduate school.....	38.0%	52.9%

There were no differences between clients who were followed up and not followed up on employment in the 30 days before entering treatment (see Table AC.10).

TABLE AC.10. EMPLOYMENT IN THE 30 DAYS BEFORE ENTERING TREATMENT

	FOLLOWED UP	
	NO n = 158	YES n = 34
<b>Employment</b>		
Not currently employed .....	46.2%	55.9%
Full-time.....	40.5%	38.2%
Part-time.....	7.6%	2.9%
Occasional .....	5.7%	2.9%

The majority of clients in both groups reported that their usual living arrangement in the 6 months before entering the program was living in their own home or apartment (see Table AC.11). There were no differences between the groups on clients who considered themselves homeless.

TABLE AC.11. LIVING SITUATION OF CLIENTS BEFORE ENTERING TREATMENT

	FOLLOWED UP	
	NO n = 158	YES n = 34
<b>Usual living arrangement in the 6 months before entering the program</b>		
Own home or apartment .....	64.6%	47.1%
Someone else’s home or apartment.....	29.1%	44.1%
Institutional facility, hotel or on the street .....	6.3%	8.8%
<b>Homelessness</b>		
Consider themselves to be currently homeless .....	15.2%	11.8%

Measures of economic hardship may be better indicators of the actual day-to-day stressors clients face than a measure of income. Therefore, the intake interview included several questions about clients’ ability to meet expenses for basic needs and food insecurity (SIPP). Clients were asked eight items, five of which asked about difficulty meeting basic needs such as food, shelter, utilities, and telephone, and three items asked about difficulty obtaining needed health care for financial reasons.

Table AC.12 shows that there were no significant difference between clients who were followed up and not followed up on difficulty meeting basic living needs such as shelter, utilities, phone, and food. About 44% of clients who were followed up reported difficulty meeting basic living needs at intake compared to 38.0% of clients who were not followed up. In addition, there was no difference between clients who were followed up and clients who were not followed up on being unable to receive needed health care for financial reasons (35.3% and 25.9%, respectively).

TABLE AC.12. DIFFICULTY MEETING BASIC NEEDS IN THE 6 MONTHS BEFORE ENTERING TREATMENT

	FOLLOWED UP	
	NO n = 158	YES n = 34
Had difficulty meeting basic living needs (e.g. shelter, utilities, phone, food).....	38.0%	44.1%
Had difficulty obtaining needed health care for financial reasons.....	25.9%	35.3%

## CRIMINAL JUSTICE SYSTEM INVOLVEMENT AT INTAKE

Close to one in ten clients in both groups were under supervision by the criminal justice system when they entered the program (e.g., probation, parole), with no significant difference by follow-up status (see Table AC.13).

There was no significant difference in the number of clients who were arrested for any charge in the 6 months before entering the program by follow up status (13.9% vs 5.9%). There was no significant difference between the groups for the number of clients who were incarcerated at least one night in the 6 months before entering the program. Of those who had been incarcerated, there were no significant differences in the average number of nights spent in jail with clients who completed a follow-up interview reporting an average of 31.5 nights and clients who did not complete a follow-up interview reporting an average of 41.7 nights.

TABLE AC.13. CRIMINAL JUSTICE SYSTEM INVOLVEMENT WHEN ENTERING TREATMENT

	FOLLOWED UP	
	NO n = 158	YES n = 34
Currently under supervision by the criminal justice system.....	9.5%	8.8%
<b>Arrested for any charge in the 6 months before entering treatment.....</b>	13.9%	5.9%
Of those arrested	n = 22	n = 2
<b>Incarcerated in the 6 months before the program.....</b>	13.3%	8.8%
Of those incarcerated	n = 21	n = 3
Average number of nights in jail.....	31.5	41.7

\*p < .05.




## APPENDIX D. TRENDS IN PRESCRIPTION OPIOID, METHADONE, BUP-NX, AND HEROIN USE AMONG KORTOS CLIENTS WITH A COMPLETED FOLLOW-UP INTERVIEW

Looking at trends over time for all clients with completed follow-up interviews, the percent of clients using prescription opioids peaked in calendar year 2009 and has steadily dropped. Similarly, the percent of clients who reported using non-prescribed methadone in the 30 days before entering treatment has declined since calendar year 2008. The percent of clients who reported using bup-nx slowly increased from 2007 through 2010, dipped slightly in 2011, and then dramatically increased in 2013 and has remained fairly stable since. The use of methamphetamine was relatively steady from CY 2007 to CY2011, but has gradually increased since.

The most notable change in substance use among KORTOS clients is for heroin. Small percentages of KORTOS clients reported using heroin in the 30 days before entering treatment from 2007 through 2011. The percentage tripled from CY 2011 (7.8%) to CY 2012 (26.7%) and then the percentage doubled to 53.7% in CY 2013. In CY 2015 the percent increased again with two-thirds of KORTOS clients (66.7%) reporting heroin use in the 30 days before intake and has increased further to 71.4% in CY 2020.

FIGURE AD.1. PERCENT OF FOLLOWED-UP CLIENTS REPORTING NON-PRESCRIBED USE OF PRESCRIPTION OPIOIDS, METHADONE, BUP-NX, HEROIN, AND METHAMPHETAMINES IN THE 30 DAYS BEFORE ENTERING TREATMENT (n = 2,177)<sup>103</sup>



<sup>103</sup> Clients who reported being in a controlled environment all 30 days before entering treatment are not included in this analysis.

## APPENDIX E. COMPARISON OF CLIENTS WHO WERE CURRENTLY INVOLVED WITH THE CLINIC AT FOLLOW-UP AND THOSE WHO WERE NOT INVOLVED ON TARGETED FACTORS

Clients who completed a follow-up interview but were no longer involved with the clinic ( $n = 13$ ) are compared in this section with clients who did completed a follow-up interview and were still involved with the clinic ( $n = 21$ ) on targeted factors.

### DEMOGRAPHIC FACTORS AT INTAKE

There were a few significant differences between clients who were still involved with the clinic at follow-up and clients who were not. More clients who were still involved in the clinic at follow-up were White and more clients who were not involved in the clinic at follow-up were married at intake.

TABLE AE.1. COMPARISON OF DEMOGRAPHICS FOR CLIENTS WHO WERE INVOLVED IN THE CLINIC AT FOLLOW-UP AND CLIENTS WHO WERE NOT INVOLVED IN THE CLINIC AT FOLLOW-UP

	INVOLVED IN THE CLINIC AT FOLLOW-UP	
	NO n = 13	YES n = 21
<b>Age</b> .....	43.4 years	37.6 years
<b>Gender*</b>		
Male.....	53.8%	33.3%
Female.....	46.2%	66.7%
<b>Race*</b>		
White.....	84.6%	100.0%
African American.....	15.4%	0.0%
<b>Marital status*</b>		
Never married.....	23.1%	28.6%
Married.....	53.8%	9.5%
Separated or divorced	7.7%	19.0%
Widowed.....	0.0%	4.8%
Cohabiting.....	15.4%	38.1%

\* $p < .10$ .

## SUBSTANCE USE AT FOLLOW-UP

Use of illegal drugs in past the 6 months at follow-up is presented by follow-up status in Table AE.2. Significantly more clients who were still involved in the clinic at follow-up reported using any illegal drugs in the past 6 months at follow-up.

TABLE AE.2. PERCENT OF CLIENTS REPORTING ILLEGAL DRUG USE IN THE PAST 6 MONTHS AT FOLLOW-UP

	INVOLVED IN THE CLINIC AT FOLLOW-UP	
	NO n = 13	YES n = 21
Any illegal drug* .....	30.8%	61.9%
Prescription opioid/opiate (illegal use).....	15.4%	4.8%
Heroin .....	15.4%	38.1%
Marijuana .....	15.4%	28.6%
CNS depressants .....	0.0%	4.8%
Cocaine .....	0.0%	4.8%
Stimulants (amphetamines, methamphetamine, prescription stimulants).....	0.0%	14.3%
Non-prescribed bup-nx .....	0.0%	0.0%
Non-prescribed methadone.....	0.0%	0.0%
Other illicit drugs (hallucinogens, inhalants, synthetic drugs) .....	0.0%	4.8%

\*p < .10.

There were no significant differences between clients who were involved with the clinic at follow-up and clients who were not for alcohol use at follow-up (see Table AE.3).

TABLE AE.3. PERCENT OF CLIENTS REPORTING ALCOHOL USE IN THE PAST 6 MONTHS AT FOLLOW-UP

	INVOLVED IN THE CLINIC AT FOLLOW-UP	
	NO n = 13	YES n = 21
Alcohol .....	7.7%	14.3%
Alcohol to intoxication.....	7.7%	0.0%
Binge drank alcohol (i.e., drank 5 or more [4 or more for women] drinks in 2 hours.....	7.7%	0.0%

In the past 6 months at follow-up, the majority of clients reported smoking tobacco products, with no difference between clients who were involved in the clinic at follow-up and those who were not (see Table AE.4). Significantly more clients who were not involved in the clinic at follow-up reported using e-cigarettes in the past 6 months at follow-up compared to clients who were still involved in the clinic.

TABLE AE.4. PERCENT OF CLIENTS REPORTING TOBACCO USE IN THE PAST 6 MONTHS AT FOLLOW-UP

	INVOLVED IN THE CLINIC AT FOLLOW-UP	
	NO n = 13	YES n = 21
Smoked tobacco .....	69.2%	71.4%
Smokeless tobacco.....	7.7%	4.8%
E-cigarettes* .....	0.0%	23.8%

\*p < .10.

Self-reported severity of alcohol and drug use was measured with Addiction Severity Index (ASI) alcohol and drug composite scores. Alcohol and drug composite scores are presented in Table AE.5 for those clients who were not in a controlled environment all 30 days before follow-up.<sup>104</sup> The highest composite score is 1.0 for each of the two substance categories.

Close to 15% of clients who were not involved in the clinic at follow-up and 23.8% of clients who were still involved in the clinic at follow-up met or surpassed the Addiction Severity Index (ASI) composite score cutoff for alcohol and/or drug severe SUD at follow-up with no difference by group. The average score for the drug severity composite score was 0.10 for clients who were not involved in the clinic at follow-up and 0.14 for clients who were still involved in the clinic, which was not a significant difference (see Table AE.5).

TABLE AE.5. SUBSTANCE ABUSE AND DEPENDENCE PROBLEMS AT FOLLOW-UP

Recent substance use problems among clients who were....	Not in a controlled environment all 30 days before follow-up	
	INVOLVED IN THE CLINIC AT FOLLOW-UP	
	NO n = 13	YES n = 21
Percentage of clients with ASI composite score equal to or greater than cutoff score for		
Severe alcohol or drug use disorder.....	15.4%	23.8%
Severe alcohol use disorder .....	7.7%	19.0%
Severe drug use disorder .....	7.7%	19.0%
Average composite score for alcohol use <sup>a</sup> .....	.04	.13
Average composite score for drug use <sup>b</sup> .....	.10	.14

<sup>a</sup> Score equal to or greater than .17 is indicative of severe alcohol use disorder.

<sup>b</sup> Score equal to or greater than .16 is indicative of severe drug use disorder.

<sup>104</sup> Clients who were in a controlled environment all 30 days before follow-up were not included in this analysis because being in a controlled environment limits one’s access to substances.

## MENTAL HEALTH AT FOLLOW-UP

The mental health questions included in the KORTOS follow-up interview are not clinical measures, but instead are research measures. A total of 9 questions were asked to determine if they met study criteria for depression, including at least one of the two leading questions: (1) “Did you have a two-week period when you were consistently depressed or down, most of the day, nearly every day?” and (2) “Did you have a two-week period when you were much less interested in most things or much less able to enjoy the things you used to enjoy most of the time?”. There was no significant difference between clients who were not involved in the clinic at follow-up and clients who were involved in the clinic for the percent of clients who met criteria for depression: 7.7% vs. 14.3% (see Table AE.6).

A total of 7 questions were asked to determine if clients met study criteria for generalized anxiety, including the leading question: “In the past 6 months, did you worry excessively or were you anxious about multiple things on more days than not for all 6 months (like family, health, finances, school, or work difficulties)?”. There was no significant difference between clients who were not involved in the clinic at follow-up and clients who were involved in the clinic for the percent of clients who met criteria for generalized anxiety: 7.7% vs. 19.0%.

Two questions were included in the intake interview that asked about thoughts of suicide and attempted suicide in the past 6 months at follow-up. There was no significant difference between clients who were not involved in the clinic at follow-up and clients who were involved in the clinic for thoughts of suicide or suicide attempts.

TABLE AE.6. PERCENT OF CLIENTS REPORTING MENTAL HEALTH PROBLEMS IN THE PAST 6 MONTHS AT FOLLOW-UP

	INVOLVED IN THE CLINIC AT FOLLOW-UP	
	NO n = 13	YES n = 21
Depression.....	7.7%	14.3%
Generalized anxiety.....	7.7%	19.0%
Suicidality (e.g., thoughts of suicide or suicide attempts).....	0.0%	0.0%

## PHYSICAL HEALTH AT FOLLOW-UP

To give an idea of the physical health of clients at follow-up, Table AE.7 presents the percent of clients that reported health problems at follow-up. There was not a significant difference for experiencing chronic pain (i.e., pain that lasted more than 3 months) at follow-up between clients who were not involved in the clinic at follow-up and clients who were. There were no significant differences for the average number of days physical health or mental health was poor in the past 30 days at follow-up and type of medical insurance between the two groups.

TABLE AE.7. PHYSICAL HEALTH STATUS AT FOLLOW-UP

	INVOLVED IN THE CLINIC AT FOLLOW-UP	
	NO n = 13	YES n = 21
Chronic pain (lasting at least 3 months).....	46.8%	44.1%
Average number of days physical health was poor in the past 30 days at follow-up.....	3.9	1.7
Average number of days mental health was poor in the past 30 days at follow-up .....	2.5	4.3
<b>Type of medical insurance</b>		
No medical insurance .....	0.0%	4.8%
Insurance through employer.....	46.2%	28.6%
Medicaid .....	46.2%	61.9%
Medicare .....	0.0%	4.8%
VA/Champus/Tricare.....	7.7%	0.0%

## QUALITY OF LIFE

There was no significant difference between clients who were still involved in the clinic and follow-up and those who were not for average quality of life rating.

TABLE AE.8. CLIENT RATING OF QUALITY OF LIFE AND WELL-BEING AT FOLLOW-UP

	INVOLVED IN THE CLINIC AT FOLLOW-UP	
	NO n = 13	YES n = 21
Quality of life rating (Ratings were from 1 = 'Worst imaginable' to 5 = 'Good and bad parts were about equal' to 10 = 'Best imaginable') .....	8.1	8.3

There were no differences between clients who were not involved in the clinic at follow-up and clients who were still involved at follow-up on employment in the past 30 days at follow-up (see Table AE.9).

TABLE AE.9. EMPLOYMENT IN THE 30 DAYS AT FOLLOW-UP

	INVOLVED IN THE CLINIC AT FOLLOW-UP	
	NO n = 13	YES n = 21
<b>Employment</b>		
Not currently employed .....	30.8%	47.6%
Full-time.....	69.2%	42.9%
Part-time.....	0.0%	9.5%
Occasional .....	0.0%	0.0%

The majority of clients in both groups reported that their usual living arrangement in the past 6 months at follow-up was living in their own home or apartment (see Table AE.10). None of the clients in either group considered themselves homeless at follow-up.

TABLE AE.10. LIVING SITUATION OF CLIENTS AT FOLLOW-UP

	INVOLVED IN THE CLINIC AT FOLLOW-UP	
	NO n = 13	YES n = 21
<b>Usual living arrangement in the 6 months before entering the program</b>		
Own home or apartment .....	84.6%	71.4%
Someone else's home or apartment.....	15.4%	28.6%
Institutional facility, hotel or on the street .....	0.0%	0.0%
<b>Homelessness</b>		
Consider themselves to be currently homeless .....	0.0%	0.0%

Measures of economic hardship may be better indicators of the actual day-to-day stressors clients face than a measure of income. Therefore, the intake interview included several questions about clients' ability to meet expenses for basic needs and food insecurity (SIPP). Clients were asked eight items, five of which asked about difficulty meeting basic needs such as food, shelter, utilities, and telephone, and three items asked about difficulty obtaining needed health care for financial reasons.

Table AE.11 shows that there was no significant difference between clients who were not involved in the clinic at follow-up and clients who were on difficulty meeting basic living needs such as shelter, utilities, phone, and food. In addition, there was no significant difference between clients who were not involved in the clinic at follow-up and clients who were for being unable to receive needed health care for financial reasons.

TABLE AE.11. DIFFICULTY MEETING BASIC NEEDS IN THE PAST 6 MONTHS AT FOLLOW-UP

	INVOLVED IN THE CLINIC AT FOLLOW-UP	
	NO n = 13	YES n = 21
Had difficulty meeting basic living needs (e.g. shelter, utilities, phone, food).....	23.1%	47.6%
Had difficulty obtaining needed health care for financial reasons.....	7.7%	28.6%

## CRIMINAL JUSTICE SYSTEM INVOLVEMENT AT FOLLOW-UP

Around 10% of clients who were still involved in the clinic at follow-up and none of the clients who were not involved in the clinic were under supervision by the criminal justice system at follow-up (e.g., probation, parole), with no significant difference between groups (see Table AE.12).

There was also no difference between the groups for the number of clients who were arrested for any charge or incarcerated for at least one night in the past 6 months at follow-up.

TABLE AE.12. CRIMINAL JUSTICE SYSTEM INVOLVEMENT IN THE PAST 6 MONTHS AT FOLLOW-UP

	INVOLVED IN THE CLINIC AT FOLLOW-UP	
	NO n = 13	YES n = 21
Currently under supervision by the criminal justice system .....	0.0%	9.5%
Arrested for any charge in the 6 months before entering treatment .....	0.0%	9.5%
Incarcerated in the 6 months before the program .....	0.0%	4.8%

## RECOVERY AT FOLLOW-UP

Significantly more clients who were no longer involved in the clinic than clients who were involved in the clinic at follow-up reported attending mutual help recovery meetings in the past 30 days at follow-up (61.5% vs. 14.3%, respectively). There was no significant difference between the two groups for the average number of people clients could count on for support at follow-up. There was also not a significant difference for how good they believed their chances were of getting off and staying off drugs/alcohol using a scale from 1 (Very poor) to 5 (Very good).



TABLE AE.12. CRIMINAL JUSTICE SYSTEM INVOLVEMENT IN THE PAST 6 MONTHS AT FOLLOW-UP

	INVOLVED IN THE CLINIC AT FOLLOW-UP	
	NO n = 13	YES n = 21
Attended AA/NA mutual help recovery meetings in the past 30 days at follow-up*** .....	61.5%	14.3%
Average number of people you can count on for recovery support.....	12.3	8.0
How good are chances that you can get off and stay off drugs (On a scale of 1 = 'Very poor' to 5 'Very good') .....	4.5	4.6

\*\*\*p < .01.

There were no significant differences between clients who were still involved in the clinic and clients who were not involved in the clinic at follow-up for multidimensional recovery. Almost half of clients who were still involved in the clinic at follow-up (47.6%) and 69.2% of clients who were not still involved in the clinic at follow-up were classified as having all positive dimensions of recovery (not presented in a table).