



Center on Drug and
Alcohol Research

University of Kentucky Center on Drug and Alcohol Research Behavioral Health Outcome Studies

Evidence Base for the Kentucky Treatment Outcome Study (KTOS) Assessment and Methods

September 2016

Executive Summary

The Kentucky Treatment Outcome Study (KTOS)

The Kentucky Treatment Outcome Study (KTOS) is a statewide data collection system designed to examine substance abuse treatment outcomes over time. This study started in 1996 as a result of a statute passed by the General Assembly and continues today. When initiated, the Kentucky Department of Behavioral Health, Developmental and Intellectual Disabilities (DBHDID), which was charged with carrying out the study, contracted with the University of Kentucky Center on Drug and Alcohol Research (UK CDAR) to develop and implement the study. DBHDID directed that KTOS be a statewide outcome study that documents the ongoing need for services and provide up-to-date regional and state data on substance use trends and treatment outcomes for Kentucky. **Although Kentucky is represented in a few national datasets, those national studies do not provide state, county- and regional-level data, nor do those national surveys consider Kentucky's unique cultural context.**

What Makes Kentucky Unique?

Kentucky's unique cultural context includes the fact that Kentucky has some of the highest rates in the nation for drug overdose fatalities, smoking, and serious health conditions (cancer deaths, cardiovascular related deaths, premature deaths, diabetes, obesity), along with the highest number of preventable hospitalizations and the second highest proportion of the population on disability in the nation. Other indicators show Kentucky ranks among the highest in number of self-reported poor days of physical health and mental health. Further, Kentucky ranks low in financial opportunity, financial well-being, and the percent of children living in poverty (Gallup Polls, 2014, 2015; Hess et al., 2015; Social Security Administration, 2011; United Health Foundation, 2015). **Given this context, the KTOS assessment is designed to identify drug use trends, substance use-related co-morbidities, and treatment outcomes in the context of Kentucky specific economic and health-related concerns.**

What is Evidence-Based Assessment?

Evidence-based assessment is a critical component of evidence-based practice but has **received limited research attention. Information obtained from evidence-based assessments can be used to help determine areas to target in treatment, to develop a case conceptualization, to increase client engagement, and to objectively monitor treatment.** The scope of evidence-based assessment includes both the process through which the assessment is conducted and the instruments utilized for evaluation.

The evidence base for the KTOS assessment conforms to the recommendations for evidence-based assessments for treatment providers in public agencies. The KTOS assessment:

1

Is based on theory and research about substance use-related comorbidities such as depression, anxiety, suicidality, criminal justice system involvement, quality of life, difficulties in employment, medical problems, housing instability, and recovery supports.

2

Is appropriate for the context of Kentucky substance abuse treatment programs and includes measures that consider the unique features of Kentucky.

3

Is face-valid and user-friendly, in part because of almost 20 years of experience, but also because it targets areas identified in theory and research as related to substance use, relapse, and treatment outcomes. KTOS is also relatively short, easy to use, and is provided to treatment centers at no cost. Further, once the intake assessment is completed, clinical providers can download a client-specific narrative report, which incorporates the information provided by the client during the assessment and provides the ASAM III level of care recommendations. A statewide survey of substance abuse directors and clinicians with the Community Mental Health Centers (CMHC) that participate in KTOS found that overall, clinicians and directors reported positive experiences with each of the components of KTOS including the KTOS assessment, KTOS annual outcome report, the Client Information System, the client narratives, and the ASAM Level of care recommendation.

4

Is made up of five core components (substance use, mental health, victimization and trauma, criminal justice system involvement, and quality of life) each with **strong reliability and validity research support** and three supplemental components (health and stress-related health consequences, economic and living circumstances, and recovery supports), most of which have strong reliability and validity research support.

5

Is focused primarily on dynamic or changeable factors rather than static factors by including measures such as mental health symptoms, quality of life, and recovery supports which can be changed within the treatment context rather than more static constructs generally thought to be less amenable to change through substance abuse treatment (e.g., antisocial personality disorder).

6

Has been used for 20 years with no reports of adverse reactions or consequences due to the assessment or the research procedures. In addition, KTOS assessment data are entered into an online, secure Client Information System (CIS) developed and maintained by UK CDAR. This server uses HTTPS for secure data transmission, data encryption for all identifying data elements which are also stored separately from assessment responses, secure server infrastructure that is in a locked-down facility with 24/7 monitoring, and user authentication. KTOS is reviewed annually by the University of Kentucky Medical Institutional Review Board (IRB) and has a Certificate of Confidentiality issued by the Federal Department of Health and Human Services to provide the highest protection for data privacy and security.

7

Is sensitive to individual-level change so that outcomes can be measured. Results from past KTOS outcomes indicate that a significant proportion of clients benefit from substance abuse treatment as substance use and substance abuse severity declines, quality of life is vastly improved, and criminal justice system involvement is significantly reduced. The 12-month follow-up uses the same KTOS evidence-based assessment that is conducted at intake in order to examine change over time. The study has a follow-up rate with over 75% of randomly selected clients with over 1,250 follow-up assessments completed each year.

Additional benefit

Provides data analysis and dissemination. An additional benefit of the Kentucky Treatment Outcome Study is that state-level trends in substance use along with the co-occurring anxiety and depression, criminal justice system involvement, employment and economic status, and quality of life trends for clients entering publicly funded treatment are provided each year. An important benefit of state-level outcome studies is that funders and legislators can see up-to-date state specific data to provide evidence of need for new programs, continuation of current programs, and changes in programmatic policies. Key trends in substance use and policy needs fluctuate annually depending on economic and other state-specific sociopolitical issues, each year's analytical findings, the latest research, and legislative research commission requests, making the need for easily-modifiable annual data collection even more important. In addition to annual statewide reports, the KTOS data is used for community-level reports on client characteristics and outcomes for communities applying for Federal or other grants. Specifically,

1. UK CDAR BHOS has produced **18 annual reports** using intake data and follow-up data from 1997 through 2016.
2. UK CDAR BHOS has produced **over 95 regional and other ad hoc reports using KTOS data along with over 25 different translational research products.**
3. The KTOS data has also been used in **numerous presentations and meetings** with clinical providers, agency boards of directors, and other state planning agencies that work closely with DBHDID.
4. **Nine peer reviewed, scholarly articles** using KTOS data have also been published.

The evidence base for the KTOS assessment suggests it is a robust, pragmatic, reliable, and valid assessment, which provides statewide and regional data about Kentucky drug use trends, substance use-related comorbidities, and substance abuse treatment outcomes.

Table of Contents

Executive Summary.....	2
Introduction	6
What Is Evidence-Based Assessment?.....	7
KTOS Intake and Follow-up: Evidence-Based Assessments.....	13
KTOS Core Assessment Components.....	13
1. Substance Use.....	13
2. Mental Health.....	16
3. Victimization and Trauma.....	19
4. Criminal Justice System Involvement.....	21
5. Quality of Life.....	22
KTOS Supplementary Assessment Components.....	23
1. Health and Stress-Related Health Consequences.....	23
2. Economic and Living Circumstances.....	25
3. Recovery Supports	27
KTOS demographic Information	27
Conclusion.....	28
Appendix A: References.....	30
Appendix B: Reports Generated Using KTOS Data	39
Appendix C: KTOS Publications.....	50

Evidence Base for the Kentucky Treatment Outcome Study (KTOS) Assessment and Methods

September 2016

University of Kentucky
Center on Drug and Alcohol
Research
Behavioral Health Outcome
Studies

TK Logan, Jennifer Cole, Jaime
Miller, & Allison Scrivner

Introduction

As Federal government and other funding agencies continue to merge and decrease funding for substance abuse, prevention, and mental health services, it is critical to have statewide outcome studies that document the ongoing need for services and provide up-to-date regional and state data on substance use trends and treatment outcomes for Kentucky. Although Kentucky is represented in a few national datasets, those national studies do not provide the state, county- and regional-level data and those national surveys do not consider or account for Kentucky's unique cultural context.

The Kentucky Treatment Outcome Study (KTOS) is a statewide substance abuse treatment evaluation study initiated in 1996 by the Department of Behavioral Health, Developmental, and Intellectual Disabilities (DBHDID) in collaboration with University of Kentucky Center on Drug and Alcohol Research (UK CDAR) to serve as a means of uniformly collecting and analyzing annual outcome information to meet the legislative requirement for KRS 222.465.¹ Data collection is required of all state-funded treatment providers including outpatient, residential, or inpatient treatment programs licensed as a chemical dependency treatment service. The first version of the KTOS assessment was implemented statewide in 1997 and has gone through numerous revisions based on stakeholder feedback, stakeholder needs, and analytic findings. The core of KTOS is: (1) the comprehensive web-based intake assessment; (2) CDAR conducted follow-up assessment; and, (3) data analysis and dissemination.

Although Kentucky is represented in a few national datasets, those national studies do not provide the state, county- and regional-level data and those national surveys do not consider or account for Kentucky's unique cultural context.

Treatment intake data are collected by community mental health center staff as clients enter treatment (including outpatient, outpatient intensive, and inpatient) using the evidence based KTOS intake assessment. Client responses are entered into an online secure Client Information System (CIS) developed and maintained by UK CDAR. Once the intake assessment is completed, clinical providers can download a client-specific narrative report, which incorporates the information provided by the client during the assessment and provides the ASAM III level of care recommendations. UK CDAR also conducts telephone follow-up interviews 12-months after completion of the intake using the evidence-based KTOS follow-up assessment with a randomly selected sample of clients who consent to participate in the follow-up at the intake and again when they are re-contacted. The study has a high follow-up rate of over 75% and completes over 1,250 follow-up assessments each year.

¹ A description of KRS 222.465 can be found at <http://www.lrc.ky.gov/statutes/statute.aspx?id=9953>.

What Is Evidence-Based Assessment?

Evidence-based assessment is an essential part of evidence-based practice but has received limited research attention (Beidas, Stewart, & Walsh, 2015; Jensen-Doss, 2015). Information obtained from evidence-based assessments can be used to help determine what to target in treatment, to develop a case conceptualization, to increase client engagement, and to objectively monitor treatment progress (Christon, McLeod, & Jensen-Doss, 2015; Hunsley, 2015; Jensen-Doss, 2015). The scope of evidence-based assessment includes both the process through which the assessment is conducted and the instruments utilized for evaluation.

Standardized assessments are generally recommended to help determine what treatment(s) to use with clients especially when a comprehensive approach is taken rather than a narrow approach (Basco et al., 2000; Jensen-Doss, 2015; Jensen-Doss, Youngstrom, E., Youngstrom, J., Feeny, & Findling, 2014; Jewell, Handwerk, Almquist, & Lucas, 2004; Tenney, Schotte, Denys, van Megen, & Westenberg, 2003). Fully accounting for clients' concerns has been linked to better treatment engagement and outcomes (Jensen-Doss & Weisz, 2008; Kramer, Robbins, Phillips, Miller, & Burns, 2003; Pogge et al., 2001). Standardized assessments can also provide valuable information about treatment outcomes, and understanding treatment outcomes is a critical component of documenting the effectiveness of evidence-based practice (Beidas et al., 2015).

In general, recommendations for evidence-based assessments for treatment providers in public agencies, who tend to have more limited resources, higher workloads, and more limited time (Glasgow, 2013; Nunno, 2006; Scott & Lewis, 2015) include: (1) the use of theory and research to determine the selection of assessment targets or components most relevant to the client's situation (Hunsley & Mash, 2007); (2) contextual appropriateness for the specific setting in which the measures will be used; in other words that the assessment is appropriate for the target population, local context, and targets the relevant constructs of interest (Glasgow, 2013); (3) having face validity (i.e., measuring what people think it ought to measure) and being user-friendly (including not overburdening staff or clients); (4) having established reliability and validity; (5) measuring dynamic rather than static constructs (amenable to change); (6) not producing adverse reactions or consequences; and (7) being sensitive to change so that outcomes can be measured (Beidas et al., 2015; Glasgow, 2013; Hunsley, 2015; Hunsley & Mash, 2007).

Evidence-based measures are intended to be used in conjunction with clinician decision-making (Hunsley, 2015). The KTOS assessment is not meant to replace clinician decision-making but rather to assist in the assessment process by examining a range of potential co-occurring

The KTOS assessment is not meant to replace clinician decision-making but rather to assist in the assessment process by examining a range of potential co-occurring problems and to provide information about treatment outcomes. The KTOS assessment can be used to inform treatment(s), engage clients through self-report, and monitor outcomes.

problems and to provide information about treatment outcomes. The KTOS assessment can be used to inform treatment(s), engage clients through self-report, and monitor outcomes.

The following bulleted points highlight how the KTOS assessment meets each of the evidence-based criteria listed above and one additional benefit is described:

1. **Use of theory and research.** The KTOS assessment includes a set of instruments developed to provide screening and assessment of psychosocial issues identified in theory and research as related to substance use including difficulties in employment, medical problems, housing instability, depression, anxiety, suicidality, criminal justice system involvement, and recovery supports (or engagement in the treatment process; Peters, Sherman, & Osher, 2008). Specific research support is outlined with each assessment component described in the next section.
2. **Contextual appropriateness.** The KTOS assessment was originally developed to consider the unique sociocultural context of Kentucky. Revisions were made as needed after data analysis and feedback from users and other stakeholders (see Figure 1 on the next page).
 - More specifically, the Kentucky context includes being 3rd in the nation for highest drug overdose-related deaths and 2nd in the nation for highest smoking rates (United Health Foundation, 2015). In addition, Kentucky is in the bottom five worst states for overall well-being (which considers social, financial, and physical indicators; Gallup Polls 2014; 2015), preventable hospitalizations (50th), cancer deaths (50th), premature deaths (47th), diabetes (45th), obesity (44th), and is in the bottom 10 for cardiovascular deaths (43rd). Kentucky was ranked 2nd in the nation for the highest number of self-reported poor physical health days in the past 30 days and 4th in the nation for the overall number of self-reported poor mental health days in the past 30 days.
 - Further, the Social Security Administration (2011) indicates 8.1% of the Kentucky population between 18 and 64 are on disability which is the 2nd highest in the nation. Another study found Kentucky had the 4th highest prevalence rate (16.1%) for disability among non-institutionalized working age individuals (ages 21 – 64) in the U.S. 50 states and territory of Puerto Rico (Erickson, Lee, & von Schrader, 2014).
 - Another report on poverty and economic opportunity ranks Kentucky as 48th in the nation for economic opportunity (Hess et al., 2015) while Gallup Polls (2014) ranked Kentucky as 46th in the nation for financial well-being (which considers having enough money for food, health care, and people’s perceived standard of living). Kentucky also was ranked 2nd in the nation for highest percentage of children living in poverty (United Health Foundation, 2015).

Figure 1.

Kentucky in Context

The KTOS assessment was originally developed to consider the unique features of Kentucky and has been revised frequently after data analysis and feedback from users and other stakeholders to consider the unique context of Kentucky.

Kentucky ranks among the highest in the nation for drug overdose deaths and smoking:



Source: United Health Foundation, 2015

3rd highest
in the nation for
DRUG OVERDOSE DEATHS



2nd highest
in the nation for
SMOKING RATES

Kentucky ranks as one of the unhealthiest states in the nation:



Source: Gallup Poll, 2014, 2015; United Health Foundation, 2015

50th
in the nation for
PREVENTABLE HOSPITALIZATIONS

45th
in the nation for
DIABETES

50th
in the nation for
CANCER DEATHS

44th
in the nation for
OBESITY

47th
in the nation for
PREMATURE DEATHS

43rd
in the nation for
CARDIOVASCULAR DEATHS

Kentucky also ranks as one of the highest in the nation for the number of disability recipients as well as the number of children in poverty:



Source: Gallup Polls, 2014; Hess et al., 2015; Social Security Administration, 2011; United Health Foundation, 2015

2nd highest
in the nation for
POPULATION ON DISABILITY

2nd highest
in the nation for
CHILDREN IN POVERTY

Kentucky ranks as one of the states with the lowest financial well-being (which considers having enough money for food, health care, and people's perceived standard of living) and economic opportunity.

46th
in the nation for
FINANCIAL WELL-BEING

48th
in the nation for
ECONOMIC OPPORTUNITY

3. **Face valid and user-friendly.** The KTOS assessment is face valid as it focuses on components identified in theory and research as related to substance use, relapse, and treatment outcomes. Further, many standardized assessments are extremely time consuming, labor intensive, and/or costly (Beidas et al., 2015; Bumbarger & Campbell, 2012; Connors, Arora, Curtis, & Stephan, 2015; Jensen-Doss & Hawley, 2010; Peters et al., 2008). The KTOS assessment is a brief instrument (35 minutes on average) which can be used to document symptoms and patterns of substance abuse and related psychosocial problems as well as to engage clients in the treatment process by allowing clients to report their concerns and problems (Christon et al., 2015; Jensen-Doss, 2015; Peters et al., 2008; Scott & Lewis, 2015). A brief satisfaction survey with substance abuse directors and clinicians with the Community Mental Health Centers (CMHC) that participate in KTOS to gather feedback on the study was conducted in 2013.² Overall, clinicians and directors reported positive experiences with each of the components of KTOS including the KTOS assessment, KTOS annual outcome report, the Client Information System, the client narratives, and the ASAM Level of care recommendation.³ In addition, respondents reported the KTOS findings publications (e.g., annual report, findings at a glance, and fact sheet) were useful.

4. **Established reliability and validity.** The KTOS assessment has five core components and three supplemental components. The five core assessment components include: (1) substance use, (2) mental health, (3) victimization and trauma, (4) criminal justice system involvement, and (5) quality of life. The three supplemental assessment components that have been associated with substance abuse and relapse include: (1) health and stress-related health consequences, (2) economic and living circumstances, and (3) recovery supports. Each of the core assessment components and most of the supplementary components of the KTOS assessment show excellent reliability and validity. Specific reliability and validity information for each assessment component is outlined in the following section.

Each of the core assessment components and most of the supplementary components of the KTOS assessment show excellent reliability and validity.

5. **Measuring dynamic rather than static constructs.** Although KTOS does include key demographic indicators the majority of the assessment components focus on current status, symptoms, and constructs that change over time. For example, mental health symptoms, quality of life, and recovery supports are all changeable within the context of substance abuse treatment whereas measures of personality or criminal histories are considered more static or less amenable to change.

² Twelve of the fourteen substance abuse directors were contacted; two could not be reached to complete a survey. Two of the twelve directors contacted did not complete the entire survey because of either their inexperience with KTOS or because they chose not to continue the survey. Twenty-seven clinicians from twelve of the fourteen CMHCs were contacted; clinicians from two CMHCs could not be reached.

³ The *KTOS and KTOS Client Information System Satisfaction Survey Summary* can be found in Appendix C of Logan, T., Cole, J., Scrivner, A., Messer, J., Emmick, C., Spencer, M., Miller, J., & Hunt, T. (2014). *KTOS, AKTOS, KORTOS Projects FY14 Annual Progress Report*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

6. **Not producing adverse reactions or consequences.** In the 20 years of conducting KTOS no adverse reactions or consequences due to the assessment or the research procedures have been reported. Client responses are entered into an online, secure Client Information System (CIS) developed and maintained by UK CDAR. The web-based intake data collection system uses extremely robust security protocols and state-of-the art technology to provide a secure, user-friendly interface for data collection and management. This server uses HTTPS for secure data transmission, data encryption for all identifying data elements which are also stored separately from assessment responses, secure server infrastructure that is in a locked-down facility with 24/7 monitoring, and user authentication. The KTOS assessment and the research methods are reviewed annually by the CDAR team in collaboration with the state and community substance abuse and mental health treatment programs. The KTOS assessment and the research methods are also reviewed annually by the University of Kentucky Institutional Review Board (IRB) and has a Certificate of Confidentiality from the Federal Department of Health and Human Services.

7. **Sensitive to change so that outcomes can be measured.** KTOS outcome results generally suggest that clients of publicly-funded substance abuse treatment, including a variety of treatment modalities, make significant strides in all of the targeted outcomes. Specifically, there are significant decreases in use of alcohol and all drugs (except tobacco), a significant increase in full-time employment, decreases in depression, anxiety and suicidality, decreases in arrests and incarceration, and increases in recovery supports. Moreover, an estimate of the cost to Kentucky for alcohol and drug dependence in the year before treatment compared to the cost to the state for alcohol and drug dependence in the year after treatment intake, while taking into account the cost of publicly-funded treatment, show significant cost savings. However, each year there remains a significant minority that still seem to be struggling with their addiction and KTOS results can provide more detailed information about those clients.

8. **Data Analysis and Dissemination.** An added benefit of this Kentucky Treatment Outcome Study is that state-level trends in substance use along with the co-occurring anxiety and depression, criminal justice system involvement, employment and economic status, and quality of life trends for clients entering publicly funded treatment are provided each year. This data system also provides state-level trends in recovery and recovery correlates over time. An important benefit of state-level outcome studies is that funders and legislators can see up-to-date state specific data to provide evidence of need for new programs, continuation of current programs, and changes in programmatic policies. Key trends in substance use and policy needs fluctuate annually depending on economic and other state-specific sociopolitical issues, each year's analytical findings, the latest research, and legislative research commission requests, making the need for easily-modifiable annual data collection even more important. In addition to annual statewide reports, the KTOS data is used for community-level reports on client characteristics and outcomes for communities applying for Federal or other grants (see Appendix B). Specifically,
 1. UK CDAR BHOS has produced **18 annual reports** using intake data and follow-up data from 1997 through 2016.
 2. UK CDAR BHOS has produced **over 95 regional and other ad hoc reports using KTOS data along with over 25 different translational research products.**

3. The KTOS data has also been used in **numerous presentations and meetings** with clinical providers, agency boards of directors, and other state planning agencies that work closely with DBHDID.
4. **Nine peer reviewed, scholarly articles** using KTOS data have also been published (see Appendix C).

KTOS Intake and Follow-up: Evidence-Based Assessments

The following paragraphs describe the KTOS evidence base including the reliability and validity information specific to each KTOS assessment component, the relevant research related to supplementary assessment components, and assessment adaptations or additions in consideration of the Kentucky context. The KTOS assessment has demonstrated evidence that each component is sensitive to change and KTOS provides critical information about treatment outcomes and factors related to relapse.

The KTOS assessment has five core components and three supplemental components. The five core assessment components include: (1) substance use, (2) mental health, (3) victimization and trauma; (4) criminal justice system involvement, and (5) quality of life. The three supplemental assessment components that have been associated with substance abuse and relapse include: (1) health and stress-related health consequences, (2) economic and living circumstances, and (3) recovery supports. Specific demographic information is collected in the last section of the assessment.

KTOS Core Assessment Components

1. Substance Use

Substance use is the key construct to examine in a substance abuse treatment outcome study. The substance use measures include: (1) The Addiction Severity Index (ASI) substance use questions including alcohol and drug use along with the ASI composite score questions; (2) DSM-V criteria for substance use disorder; and (3) targeted questions about smoking, smokeless tobacco, and e-cigarette use, needle use and needle exchange programs per the request of community and state partners. Data from the KTOS substance use assessment component has been analyzed and included in over 95 regional and ad hoc reports describing substance abuse trends and treatment outcome trends across the state. KTOS data has also been used in several peer review publications including one that found women in rural Appalachia had disproportionately high rates of opioid and sedative/tranquilizer use compared to women from non-Appalachian areas who had higher rates of methamphetamine, cocaine, marijuana and alcohol use (Shannon, Havens, Mateyoke-Scriver, & Walker, 2009).

SUBSTANCE USE MEASURES

The KTOS substance use assessment section includes items from the alcohol and drug use sections of the *Addiction Severity Index (ASI)* (5th edition) including questions about use and questions used to compute the ASI drug and alcohol use composite scores, which are recommended for measuring substance abuse treatment outcomes (McLellan et al., 1985). The ASI was developed as a clinical/research assessment of substance use and multiple related problems found in alcohol and drug-dependent individuals. Further, the ASI is a commonly used public domain assessment (McLellan et al., 1985).

The ASI substance use measure has shown very good validity and reliability in measuring substance use. The ASI, like the KTOS, assesses several main and supplementary areas. The KTOS assessments use only the substance use domain of the ASI because of the good validity and reliability of this section and because the other KTOS components were not addressed in-depth in the ASI.⁴

The KTOS assessment primarily uses the substance use domain of the ASI because of the good validity and reliability of this section and because the other KTOS core components were not addressed in-depth in the ASI.

Several studies have examined the construct validity (i.e., the extent the measure actually measures the construct of interest) of the ASI and with different populations such as veterans, homeless individuals, and individuals with comorbid psychiatric disorders (Cronbach & Meehl, 1955). Construct validity has multiple components including: (1) criterion-related validity, which is the degree to which a measure is related to an external criterion or outcome (e.g., self-reported substance use with urinalysis); (2) convergent validity, which is the degree to which two measures of constructs that are posited by a theory to be related are actually related. For instance, if one has developed a new measure (i.e., series of related questions) of problematic substance use, one would want to examine the relationship of the scores on the new measure along with scores on other similar measures, such as the Alcohol Use Disorders Identification Test (AUDIT), CAGE, and Drug Abuse Screen Test (DAST). And (3) discriminant validity, which refers to whether constructs that are supposed to be unrelated are in fact not related (Campbell, 1959). For example, one would want to demonstrate that scores on a newly developed measure of problematic substance use were not closely correlated with measures of other constructs such as impulsivity or antisocial personality disorder.

The ASI substance use scores show high correlation with other measures of substance use. For example, the ASI was examined with other validated comparison instruments including the Michigan Alcoholism Screening Test (MAST; Selzer, 1971), Cohen and Klein Drug Use Scale (Cohen & Klein, 1971), and the Gunderson Drug Scale (Gunderson, Russell, & Nail, 1973) to determine the convergent and discriminant validity of the ASI multidimensional scores and the results showed good convergent and discriminate validity (McLellan et al., 1985). In addition, the ASI drug and alcohol composite scores correlated well with other instruments such as the Structured Clinical Interview for DSM-IV Axis I Disorders (SCID-I; First, Spitzer, Gibbon, & Williams, 1995) and with DSM-III-R

⁴ The KTOS assessment includes core components that are not addressed in-depth in the ASI. Also, not all dimensions of the ASI have equal support for their reliability and validity. For example, at least one study with a sample of individuals receiving psychiatric care did not find support for discriminant validity of the alcohol, psychiatric, legal, or medical scales (Carey, Cocco, & Correia, 1997). Second, studies conducted with special populations, such as homeless clients and individuals with severe psychiatric disorders have found low test-retest reliability in some of the ASI composite scores including the medical, legal, drug use (Corse, Zanis, & Hirschinger, 1995; Zanis, McLellan, & Corse, 1997), and family/social (Hodgins & El-Guebaly, 1992). Third, there are some problems with the reliability and validity of ASI severity ratings, which are based on subjective judgment of interviewers (Stöffelmayr, Mavis, & Kasim, 1994; Wertz, Cleaveland, & Stephens, 1995). The severity ratings are not intended to be used as outcome measures (McLellan et al., 1992). Thus, ASI severity items were not included in the KTOS interview instruments. Fourth, the authors acknowledge that the family/social dimension of the ASI concentrates on individuals' conflicts with family and other persons; however, other critical dimensions of family and social functioning are not included in the ASI (McLellan et al., 1992).

diagnoses (Appleby, Dyson, Altman, & Luchins, 1997) and DSM-IV-TR diagnoses (Rikoon, Cacciola, Carise, Alterman, & McLellan, 2006). In the Rikoon et al. (2006) study ASI composite scores for alcohol use and drug use identified dependent clients with 85% sensitivity and 80% specificity when ASI composite scores were matched to independent clinical diagnoses.

Other studies have examined how well the ASI self-reported substance use questions correlate with urinalysis results (Chermack et al., 2000; Zanis, McLellan, & Randall, 1994). In a sample of 563 clients admitted for treatment in substance abuse treatment outpatient clinics, conditional kappa values were good and indicated high levels of agreement between self-reported substance use and urinalysis: highest for cannabis (0.93) and lowest for opioids (0.84; Denis et al., 2012). Contrary to what may be expected, most of the discordance between self-reported substance use and urinalysis was because clients reported use of a particular substance but had a negative urinalysis for that substance. Part of the discrepancy is due to the fact that the 30-day self-report period is longer than the time frame captured in urinalysis results.

In general, examinations of various facets of the reliability of multiple dimensions of the ASI have found good interrater reliability and good test-retest reliability for the substance use composite scores (Calsyn et al., 2004; Mäkelä, 2004; McLellan et al., 1985; Wertz et al., 1995). Test-retest reliability, which is a measure of consistency of responses to the same set of questions at two periods, has been examined by administering the ASI interview to the same persons typically 3 days to 10 days apart (Mäkelä, 2004). Interrater reliability, which is the estimate of the equivalence of the responses between more than one rater, has been examined in these studies by having the rater observe the interview being conducted by the primary reviewer through a one-way mirror or via a videotaped recording and recording the interviewees' responses (Stöffelmayr et al., 1994). The degree of agreement between the primary interviewer's recorded responses and the observer's recorded responses is interrater reliability.

A third type of reliability that has been examined in studies is internal consistency reliability, which is a measure of the correlation between several items that purportedly measure the same construct. In other words, low correlations between items that purportedly measure the same construct indicate that the items are likely not measuring the same construct. In a review of studies that examined the reliability and validity of the ASI, Mäkelä (2004) discussed how three of the seven composite scores had consistently been found to have high internal consistency reliability: alcohol use, medical status, and psychiatric status.

DSM-V MEASURE

The DSM-V diagnostic criteria for substance use disorders included in the KTOS assessment⁵ are similar to the criteria for DSM-IV, which has evidence of excellent test-retest reliability (Hasin et al.,

⁵ The difference in diagnostic criteria of the DSM-V from the DSM-IV are the deletion of the legal problems criterion, addition of the cravings criterion, and lack of distinguishing between abuse and dependence in the DSM-V. Instead the threshold of two or more criteria is used to diagnose substance use disorder in the DSM-V. Because the DSM-V is a relatively recent revision, no reliability and validity studies have been conducted using the DSM-V criteria for diagnosing substance use disorder. Nonetheless, the slight differences between the DSM-IV and DSM-V diagnostic criteria suggest the DSM-V diagnostic criteria for substance use disorders will also have good reliability and validity once the body of research is conducted.

1996) and validity. For example, Horton, Compton, and Cottler (2000) found excellent test-retest reliability in a sample of African American and Caucasian individuals with alcohol dependence ($k = 0.78$, $k = 0.80$, respectively) and opiate dependence ($k = 0.77$, $k = 0.71$, respectively). Evidence of criterion-related validity is provided by genetics research that some genetic variants lower the threshold for the induction of nicotine dependence, which is summarized by Hogg and Bertrand (2004). In a national probability sample, the 1992 National Longitudinal Alcohol Epidemiologic Survey (NLAES), diagnosis of alcohol abuse and dependence made with the DSM-IV was compared with criterion measured with the Alcohol Use Disorders and Associated Disabilities Interview Schedule (AUDADIS; Hasin & Paykin, 1999). The odds ratios for diagnosis of dependence vs. no diagnosis, abuse vs. no diagnosis, and dependence vs. abuse were all statistically significant with the criterion variables: alcohol consumption, treatment seeking, suicidal ideation/attempts, and alcohol-induced blackouts (Hasin & Paykin, 1999). However, the DSM-V does away with the distinction between substance abuse and dependence, substituting severity ranking instead.

TARGETED SUBSTANCE USE MEASURES

The question regarding the use of needles to inject drugs in the KTOS assessment is from the ASI. However, the specific questions targeting needle exchange programs were requested to be added by key stakeholders in the community and DBHDID to assess use of these new programs. Furthermore, due to the significant issue with smoking in Kentucky (26.2% of the population, which is the second highest rate in the nation) along with e-cigarette use which is growing each year (Barrington-Trimis et al., 2016; Singh et al., 2016), use of smoking tobacco, smokeless tobacco, and e-cigarettes are assessed with items that are worded to be consistent with the alcohol and drug use questions. The age of first use for smoking, using smokeless tobacco, first alcoholic drink (other than a few sips), and first used illicit drugs is also included in the KTOS assessment.

2. Mental Health

The goal of administering mental health symptom measures is to characterize severity and change over the course of treatment (Scott & Lewis, 2015). The KTOS mental health section focuses on depression, anxiety, and suicidal ideation and suicide attempts.⁶ Items for the depression measure were adapted from the *Patient Health Questionnaire-9 (PHQ-9)* and items for the anxiety measure were adapted from the *Generalized Anxiety Disorder (GAD-7)*.

Both the PHQ-9 and the GAD-7 have been shown to be valid and reliable measures of depression and anxiety respectively.

The Patient Health Questionnaire is an instrument for making criteria-based diagnoses of eight DSM-IV mental health disorders, one of which is major depressive disorder (Kroenke, Spitzer, & Williams, 2001). The GAD-7 was developed to identify probable cases of generalized anxiety disorder and to assess symptom severity for the criteria

⁶ Different measures of depression and anxiety were incorporated into the KTOS a few years ago because the ASI mental health measures were not found to be sensitive to change over time in the Kentucky target population.

symptoms in the DSM-IV (Spitzer, Kroenke, Williams, & Löwe, 2006). Both the PHQ-9 and the GAD-7 have been shown to be valid and reliable measures of depression and anxiety respectively.

DEPRESSION

The Patient Health Questionnaire-9 (PHQ-9) includes 9 items that comprise the PHQ depression scale, which ask about the 9 symptoms listed as criteria in the DSM-IV for diagnosis of major depressive disorder (Kroenke et al., 2001). The response options range from 0 (*Not at all*) to 3 (*Nearly every day*). Thus, as a severity measure, the PHQ-9 can range from 0 to 27. A diagnosis of major depression is indicated if 5 or more of the criteria have been present at least “more than half the days” in the past 2 weeks, and 1 of the symptoms is depressed mood or anhedonia (Kroenke et al., 2001).

Data from two studies with 6,000 patients in primary care and obstetrics/gynecology clinics provide evidence that the PHQ-9 has good internal consistency reliability (Cronbach’s $\alpha = 0.89$) and excellent test-retest reliability (0.84) between the original administration of the PHQ-9 in the clinics and then 48 hours later by telephone.

Data from this same study also found evidence for good criterion-related and convergent validity of the PHQ-9 (Kroenke et al., 2001). Specifically, criterion-related validity was examined with the correlation between the PHQ-9 scores and depression diagnosis by a mental health professional who was blinded to the PHQ-9 score for 580 patients who agreed to be contacted after the initial interview. The PHQ-9 score greater than or equal to 10 had a sensitivity of 88% and a specificity of 88% for major depressive disorder. Sensitivity is a measure of how many of the individuals diagnosed with depression by a mental health professional were also identified by the PHQ-9 as having moderate to severe depression, whereas specificity is a measure of how many of the individuals who were not diagnosed with depression by a mental health professional were identified by the PHQ-9 as having minimal or mild depression (i.e., scores of less than 10). Furthermore, in the same study, several validated measures were included to examine the relationship between scores on the PHQ-9 and constructs that are hypothesized to be related to depression such as lower functioning and quality of life. The highest correlations were found between PHQ-9 scores and the functioning scales that previous studies have demonstrated would be most strongly related to depression: overall mental health, social functioning, overall functioning, and role functioning.

In the KTOS assessment the items were changed to ask if the client experienced the 9 symptom criteria nearly every day in the same two-week period and the response options were changed to 0 (*No/Absent*) to 1 (*Yes/Present*). Thus, unlike the original PHQ-9 the maximum value is 9. Individuals who responded “Yes” to the depressed mood or anhedonia items and responded “Yes” to at least 5 of the 9 criteria were classified as having met criteria for depression in the KTOS study. Excellent internal consistency reliability was found in the sample of KTOS clients who completed an intake interview in FY 2014 and were included in the 2016 Report ($n = 5,273$): Cronbach’s $\alpha = 0.954$.

ANXIETY

The Generalized Anxiety Disorder (GAD-7) was developed to identify probable cases of generalized anxiety disorder and to assess symptom severity for the criteria symptoms in the DSM-IV (Spitzer et

al., 2006). The original scale is a 7 item measure that asks about the frequency of anxiety symptoms over the last two weeks. Response options range from 0 (*Not at all*) to 3 (*Nearly every day*). Total scores range from 0 to 21 with higher scores indicating greater severity/frequency of anxiety.

Internal consistency reliability is excellent for the GAD-7, with Cronbach α ranging from 0.89 - 0.92 (Delgado et al., 2012; Löwe et al., 2008; Spitzer et al., 2006). In a second study with adults in a drug treatment facility in England, for the sample of 60 individuals who completed a retest 4 - 6 weeks later, test-retest reliability was good (Intraclass coefficient [ICC] = .85; Delgado et al., 2012).

A validation study of the GAD-7 performed in 15 primary care clinics (n = 2,740) found good criterion-related validity for the GAD-7. Specifically, the study found that a cut-off score of 10 was the ideal score to maximizing sensitivity (89%) and specificity (82%) for a diagnosis of generalized anxiety disorder (GAD) made by a mental health professional (Spitzer et al., 2006). In other words, most patients who were diagnosed with GAD by a mental health professional (89%) had GAD-7 scores of 10 or higher, whereas most patients who were not diagnosed with GAD by a mental health professional (82%) had GAD-7 scores lower than 10. Another study also examined the diagnostic accuracy of the GAD-7 in comparison with ICD-10 psychiatric diagnoses that were assessed using the Revised Clinical Interview Schedule (CIS-R)—a well-validated structured diagnostic interview (Delgado et al., 2012). A GAD-7 score of 9 or higher had a sensitivity of 80% and specificity of 86% for any anxiety disorder. In other words, 80% of individuals who were diagnosed with an anxiety disorder using the CIS-R had scores of 9 or higher on the GAD-7 and 86% of individuals who were not diagnosed with an anxiety disorder using the CIS-R had scores of 8 or lower on the GAD-7.

Moreover, convergent validity was found for the GAD-7 which was correlated with two anxiety scales: Beck Anxiety Inventory ($r = 0.72$) and the anxiety subscale of the Symptom Checklist-90 ($r = 0.74$) (Spitzer et al., 2006). More evidence of good convergent validity was found in the large effect sizes of GAD-7 severity score classification (i.e., minimal, mild, moderate, and severe) with the Medical Outcomes Study Short-Form General Health Survey (SF-20) functioning subscale scores because as anxiety symptoms increase, functioning is hypothesized to decrease (Spitzer et al., 2006).

In the KTOS assessments, the response options were changed to 0 (*No/Absent*) to 1 (*Yes/Present*). Thus, unlike the original GAD-7 the maximum value is a 7. Individuals who responded “Yes” to the item about worrying excessively or being anxious about multiple things on more days than not and “Yes” to at least 3 of the 7 symptoms were classified as having met criteria for generalized anxiety in the KTOS study. Excellent internal consistency reliability was found in the sample of KTOS clients who completed an intake interview in FY 2014 and were included in the 2016 Report (n = 5,273): Cronbach’s $\alpha = 0.975$.

SUICIDE IDEATION AND ATTEMPTS

These two items were adapted from the ASI psychiatric domain. There is no validity information for these two items; however, there is good test-retest reliability and inter-rater reliability for the ASI generally (Mäkelä, 2004; Stöffelmayr et al., 1994).

3. Victimization and Trauma

Trauma and victimization measures are included in KTOS because these experiences have been found to be linked to substance abuse in treatment populations, prison populations, and in the general public. More specifically, victimization and trauma history increases the risk for drug and alcohol use (Brady, Back, & Coffey, 2004; Kessler, Sonnega, Bromet, Hughes, & Nelson, 1995; Logan, Walker, Cole & Leukefeld, 2002; Logan, Walker, Jordan, & Leukefeld, 2006; Regier et al., 1990) and those who have a substance use disorder report more victimization and traumatic events (Cottler, Compton, Mager, Spitznagel, & Janca, 1992; Farley, Golding, Young, Mulligan, & Minkoff, 2004; Logan et al., 2002; Logan et al., 2006; Najavits et al., 2003; Najavits, Sonn, Walsh, & Weiss, 2004; Shane, Diamond, Mensinger, Shera, & Wintersteen, 2006). High rates of victimization exposure are also found in individuals involved in the criminal justice system (Goff, E. Rose, S. Rose, & Purves, 2007; Wolff & Shi, 2012) and arrests, incarceration, and violent criminal charges are associated with a history of trauma (Donley et al., 2012; Sadeh & McNeil, 2015). Additionally, the risk of relapse increases as the number of trauma events increase and those who report a relapse also report experiencing interpersonal violence more often than those who had no history of relapse (Farley et al., 2004). The KTOS assessment has three main measures of victimization and trauma: (1) the Adverse Childhood Experiences; (2) a victimization screen; and (3) a measure of Post-Traumatic Stress Disorder (PTSD).

ADVERSE CHILDHOOD EXPERIENCES

Adverse childhood experiences, defined as abuse and household dysfunction, are common. In the Adverse Childhood Experiences Study (ACES), which surveyed over 17,000 adults who were members of a health maintenance organization (HMO), the questionnaire asked about 10 major categories of childhood trauma: three types of abuse (emotional, physical, and sexual), two types of neglect (emotional and physical), and five types of family dysfunction (having a mother who experienced intimate partner violence, having a household member who was an alcoholic, having a household member who was a drug user, a household member who was incarcerated, a household member diagnosed with a mental disorder or committed suicide, or parents who were separated or divorced; Felitti et al., 1998). Almost two-thirds of HMO adult members who participated in the ACES reported at least one adverse childhood experience, and more than 1 in 5 reported 3 or more (Dong et al., 2004). As the number of adverse experiences increase the risk of many health, mental health, and social problems also increases (Edwards et al., 2005; Felitti et al., 1998). For example, increases in ACE scores is associated with a greater likelihood of depressed mood (Anda et al., 2006; Dube, Felitti, Dong, Giles, & Anda, 2003), suicide attempts (Dube et al., 2001), and panic/anxiety (Anda et al., 2006).

Of particular importance, is that the risk of alcohol or drug use increases as the number of adverse childhood experiences increases (Anda et al., 2006; Dube et al., 2003a,b; Felitti et al., 1998). Higher ACE scores are associated with initiating alcohol abuse and smoking in adolescence (Anda et al., 1999; Dube et al., 2006). Additionally, experiencing more types of childhood abuse is associated with greater likelihood of experiencing an unintended first pregnancy among women (Dietz et al., 1999). Poor self-rated health as well as health problems such as ischemic heart disease, cancer, and liver disease were more prevalent in those who reported a higher number of ACEs (Felitti et al., 1998). Poor sleep, severe obesity, and multiple somatic symptoms were increased for those with ACE scores over

4 (Anda et al., 2006). Higher ACE scores have been linked to having a higher number of health risk factors for leading causes of death in adults (Felitti et al., 1998) and a higher rate of mortality in women (Chen, Turiano, Mroczek, & Miller, 2016).

The only report of internal consistency reliability for the ACES survey was conducted with a sample of 75 urban women in a clinical and community sample (Murphy et al., 2014). In this study, internal consistency reliability was excellent (Cronbach's $\alpha = 0.88$). Test-retest reliability was examined for 658 individuals who filled out the questionnaire in two waves of the study (Dube, Williamson, Thompson, Felitti, & Anda, 2004). Kappa coefficients were in the good to excellent range as noted by Fleiss (1981) for abuse categories (0.51 – 0.69) and the household dysfunction categories (0.51 – 0.86) with the exception of having an incarcerated household member (0.46). and test-retest reliability was good for emotional abuse (94%), physical abuse (83%), sexual abuse (90%), and overall ACE score (weighted kappa = .64; Dube et al., 2004).

VICTIMIZATION SCREEN

This screen examines a wide variety of harassment and threatening situations including street harassment, sexual harassment, home invasions, robbery, burglary, assault, rape, stalking, and partner violence. The majority of the threatening situations assessed are included in national surveys (Breiding et al., 2014; Logan et al., 2006; Logan, 2016; Office for Victims of Crime, 2015; Perreault, 2015; Truman & Langton, 2015). A few modifications were made to more clearly assess exposure to specific threats including: a) assault was assessed by asking about assault with and without a weapon as well as assault by a neighbor, coworker, or schoolmate; b) firearms violence was separated into three categories including directly or indirectly threatened with a gun, being held at gunpoint, and experiencing a public or mass shooting; and c) burglary and robbery were specifically assessed by asking whether they had experienced a robbery or mugging; experienced a home break-in while not at home; and a home invasion while home. The victimization screen also assesses harassment including verbal street harassment and street sexual harassment (Kearl, 2014) as well as road rage (AAA Foundation for Traffic Safety, 2016; Sansone & Sansone, 2010; Smart, Mann, & Stoduto, 2003). The screen also assesses being kidnapped or held hostage (Blumenstein, 2015) and repeated sexual harassment at work, school, or some other place by the same individual or group of individuals (other than an [ex] partner) (Ilies, Hauserman, Schwochau, & Stibal, 2003; Stockdale, Logan, Sliter, & Berry, 2014). Additionally, this screen asks about two indirect victimization experiences including whether someone close to the participant experienced a violent victimization and if someone close to the participant or someone in their family had been murdered (Hale, 1996).

POST-TRAUMATIC STRESS DISORDER (PTSD)

One significant possible consequence of victimization is PTSD. About 1 in 10 of individuals with exposure to traumatic events developed PTSD at some point, with the highest risk of PTSD associated with assaultive violence (20.9%; Breslau et al., 1998). Individuals with PTSD have a high rate of alcohol/drug abuse or dependence in their lifetime (Kessler et al., 1995; Regier et al., 1990) and the overall prevalence of PTSD is high among substance users (Cottler et al., 1992; Najavits et al., 2003). The KTOS assessment includes a 4-item PTSD Checklist (PCL-5; Weathers et al., 2013). This 4-item PTSD checklist was derived from a 20-item self-report measure of posttraumatic stress symptoms that

is designed to reflect the changes to the diagnostic criteria for posttraumatic stress disorder (PTSD) in the fifth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5; American Psychiatric Association, 2013). The full PCL-5 displays strong internal consistency (α ranging from .83 to .98), test-retest reliability (.66 to .96), convergent (.62 to .93) and discriminant validity (.87), and sensitivity to change during treatment (Blevins, Weathers, Davis, Witte, & Domino, 2015; Bovin et al., in press; Wortmann et al., in press).

Two abbreviated scales have been developed from this 20-item measure: an 8-item scale and a 4-item scale (Price, Szafranski, van Stolk-Cooke & Gros, 2016). These abbreviated scales, particularly the 4-item scale, have performed as well or better than the 20-item PCL-5 as a screening measure for PTSD (Price et al., 2016). In a sample of veterans receiving treatment at a psychotherapy clinic, the abbreviated 4-item version of the PCL-5 demonstrated high correlations with the full PCL-5 (86%) and good internal consistency ($\alpha = 0.82$; Price et al., 2016). The 4-item scale was found to have a significantly higher specificity (0.52; $p < .01$) than the full PCL-5 (0.35) and 8-item scale (0.39) and using a cut score of 10 on the measure resulted in a sensitivity of .76 (Price et al., 2016). The 4-item scale was also just as good as the full PCL-5 at discriminating between those with PTSD and those without PTSD (AUC = .72; Price et al., 2016). These results suggest that the 4-item measure may be a better screening tool for PTSD (Price et al., 2016).

4. Criminal Justice System Involvement

The KTOS criminal justice system section asks five main questions which were adapted from the ASI: (1) nights incarcerated in the past 12 months; (2) times arrested and charged with an offense in the past 12 months; (3) misdemeanor and felony convictions in the past 12 months; (4) whether they are currently on probation; and (5) whether they are currently on parole.

In general, research suggests that self-reported criminal justice system involvement is reliable such that self-reported arrests correspond well to arrests noted in official datasets with one study finding self-reported arrests equal to or greater than arrests in the official dataset (Marquis, 1981). Another study that found 73% of those with an official arrest had also self-reported an arrest and 21% had reported an arrest although there was no official history of arrest (Maxfield, Weiler, & Widom, 2000).

Consistent with other research the KTOS criminal justice system self-reported information was found to be valid when compared with an independent database. Specifically, a sub-study to examine the concordance between self-reported criminal justice system involvement in the KTOS assessment with official records was conducted for a 20% random sample of KTOS clients with an intake during FY 2014 and a follow-up during FY 2015 ($n = 260$). Self-reported criminal justice system status was compared with the Kentucky Offender Monitoring System (KOMS) database. Half of the clients were found in KOMS ($n = 130$). Of those individuals with information in KOMS, there was a 96.2% agreement for any incarceration, or incarceration was reported on KTOS but was not

Of those individuals with information in KOMS, there was a 96.2% agreement for any incarceration, or incarceration was reported on KTOS but was not in the KOMS data at intake.

in the KOMS data at intake (KOMS does not include local jail data). There was 100% agreement with the information in KOMS at the follow-up assessment for incarceration. Additionally, there was an 83.1% agreement for probation at intake and at follow-up and 96.9% agreement for parole at intake and 91.5% agreement at follow-up.

5. Quality of Life

While symptom change often is the primary goal of treatment, quality of life assesses well-being rather than just the absence of a disorder. Quality of life is a commonly used metric for assessing the cost utility of treatment and is an important index in understanding treatment outcomes (Scott & Lewis, 2015). The KTOS Quality of Life measures have two components: (1) the *Satisfaction with Life Scale* (SWLS; Pavot & Diener, 1993), and (2) one global question asking the client to rate their quality of life today.

SATISFACTION WITH LIFE SCALE

The Satisfaction With Life Scale (SWLS) refers to a cognitive, evaluative process, in which individuals assess the quality of their lives on the basis of self-imposed standards (Pavot & Diener, 1993). In the Satisfaction With Life Scale (SWLS; Diener, Emmons, Larsen, & Griffin, 1985) the 5 items are global rather than specific.

Evidence of convergent validity has been demonstrated with moderately strong correlations of SWLS with 10 other subjective well-being scales, indicating the SWLS measures the same or a similar construct to the other subjective well-being scales (Pavot & Diener, 1993). Further, changes in life conditions are associated with changes in scores on the SWLS in the ways hypothesized; specifically, increases in caregiver burden were associated with decreases on the SWLS scores (Vitaliano, Russo, Young, Becker, & Maiuro, 1991). Moreover, consistent with theorized relationships between life satisfaction and psychological distress, several studies have found that greater distress (i.e., depression, negative affect, anxiety, and general psychological distress) is associated with lower life satisfaction, which provides evidence of convergent validity (Arrindell, Meeuwesen, & Huyse, 1991; Larsen, Diener, & Emmons, 1985). Finally, there is evidence from numerous studies that the SWLS has discriminant validity (Pavot & Diener, 1993). Specifically, scores on SWLS have not been correlated (positively or negatively) with affect intensity and impulsivity (Diener et al., 1985).

The SWLS has good internal consistency: Cronbach's $\alpha = 0.87$ (Diener et al., 1985) as well as good test-retest reliability with the correlation coefficient at 0.82 for 76 students who were re-administered the scale two months after the initial administration. In many studies using the SWLS both strong internal consistency reliability (ranging from Cronbach's $\alpha = 0.79 - 0.89$) and moderate test-retest reliability (ranging from 0.50 - 0.84) have been found (Pavot & Diener, 1993).

In the KTOS study, the response options were decreased to 5 options ranging from 1 (*Strongly disagree*) to 5 (*Strongly agree*). Internal consistency reliability was good in the most recent sample of intake interviews completed by KTOS clients ($n = 6,766$): Cronbach's $\alpha = 0.861$.

GLOBAL RATING OF QUALITY OF LIFE

One other question about the client's assessment of their quality of life is included in the KTOS assessment. Clients are asked to rate their quality of life, where 1 is approximately worst imaginable, 5 is good and bad parts are about equal, and 10 was best imaginable. Clients are allowed to select their rating anywhere along the scale.

KTOS Supplementary Assessment Components

1. Health and Stress-Related Health Consequences

The health and stress-related consequences section includes an assessment of: (1) general health status, (2) chronic pain, and (3) stress-related health consequences.

GENERAL HEALTH STATUS

General health status is included in the substance abuse treatment outcome studies because it has been found to be worse among individuals with substance abuse compared to the general population (Morgen, Astone-Twerell, Hernitche, Gunneson, & Santangelo, 2007; World Health Organization [WHO], 2014). In addition, Kentucky ranks high in the nation for a number of significant health conditions including cancer deaths, cardiovascular related deaths, premature deaths, diabetes, and obesity.

General health status is included in the substance abuse treatment outcome studies because it has been found to be worse among individuals with substance abuse compared to the general population.

The general health questions were adapted from the Behavioral Risk Factor Surveillance System (BRFSS) Health-Related Quality of Life (HRQOL; Centers for Disease Control and Prevention, 2000; Hennessy, Moriarty, Zach, Scherr, & Brackbill, 1994). Studies show that the health questions are correlated with each other such that those who self-reported fair/poor overall health also reported more days that their physical and mental health were not good (Centers for Disease Control and Prevention, 2000; Hennessy et al., 1994). Good test-retest reliability was also found for the Healthy Days questions ($r = 0.75$; Andresen, Catlin, Wyrwich, & Jackson-Thompson, 2003). In a validation study of the BRFSS HRQOL, the physical health not good, mental health not good, and days not good health limited activities items were correlated with the SF-36 HRQOL scales in expected ways, demonstrating good criterion-related validity (Newschaffer, 1998). Specifically, the number of days the respondent's physical health was not good and the number of days that not good health limited activities were significantly, negatively correlated with the SF-36 scales meaning that the higher days of reported poor physical or mental health the lower scores on general health, physical functioning, physical role, mental health, emotional role, social functioning, and vitality. Similarly, the item about the number of days respondents' mental health was not good was significantly, negatively correlated with SF-36 HRQOL scales, with the exception of the physical functioning scale (Newschaffer, 1998).

CHRONIC PAIN

There is a connection between chronic pain and prescription opioids, and a connection between chronic pain and relapse (Atkinson, Slater, Patterson, Grant, & Garfin, 1991; Edlund, Sullivan, Han, & Booth, 2013; Mertens, Lu, Parthasarathy, Moore, & Weisner, 2003; Sheu et al., 2008) and this association has also been found using KORTOS data (the KORTOS assessment is based on the KTOS assessment

There is a connection between chronic pain and prescription opioids, and a connection between chronic pain and relapse.

structure and components⁷; Stevenson, Cole, Walker, & Logan, 2014). Given the significant problem of nonprescription opioid use in Kentucky, it is critical to include an assessment of chronic pain. In general, nonprescription opioid use is a continuing health concern in Kentucky where 4.1% of adults report nonmedical use of pain relievers (Substance Abuse and Mental Health Services Administration, 2015). The two most frequently reported illicit drugs mentioned as clients' primary substance of abuse were prescription opioids and heroin in 2013 (Center for Behavioral Health Statistics and Quality, 2015). Since 2000, the rate of deaths from drug overdose involving opioids has increased 200% (Rudd, Aleshire, Zibbell, & Gladden, 2016). The United Health Foundation (2015) has Kentucky ranked as 3rd in the nation for highest drug overdose-related deaths with 24 deaths per 100,000 in 2011 - 2013. In 2014, Kentucky had the 4th highest age-adjusted drug overdose death rate in the United States, with 24.7 deaths per 100,000 people (Rudd et al., 2016), and prescription opioids was the primary drug class involved in drug overdose deaths (Slavova, Bunn, & Gao, 2015).

The chronic pain questions included in the KTOS assessment were adapted from the *Brief Pain Inventory (BPI)* which is one of the most widely used tools for assessing clinical pain and has been shown to appropriately measure pain caused by many different clinical conditions (Cleeland, 2009; Cleeland & Ryan, 1994). In an early study of validity and reliability, the BPI was given to cancer patients as well as rheumatoid arthritis patients and the correlation patterns among pain and interference measures were different for diseases with different pain mechanisms. Patients with rheumatoid arthritis were tested early in the day when their pain may be at its worst and showed a high correlation between worst pain scores and current pain scores (.71) while the same correlation among breast, colorectal, and gynecological cancer patients was much lower (.35, .27, and .42; Daut, Cleeland & Flanery, 1983). There are four severity items on the BPI that are rated 0-10 and can be averaged to get a composite score. The KTOS assessment uses only one of these items – rating the client's pain on average. Using this single question as a representation of pain severity is supported by the FDA Draft Guidance for Industry: Patient-Reported Outcome Measures (Cleeland, 2009).

⁷ Copies of Logan, T., Cole, J., Miller, J., Scrivner, A., & Walker, R. (2016). *Evidence Base for Kentucky Opiate Replacement Treatment Outcome Study Assessment and Methods*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research are available upon request.

STRESS-RELATED HEALTH CONSEQUENCES SCALE

Members of the UK CDAR BHOS research team developed a scale to measure recent stress-related health consequences (Logan & Walker, 2010). Chronic exposure to stress can tax the body by continuously activating the stress response, which alters the body's normal way of responding to external stimuli (McEwen, 2000). When this process interferes with the body's ability to maintain equilibrium, an individual's allostatic load increases (McEwen, 2000, 2004). High allostatic load over time is associated with physical health and mental health problems such as a weakened immune system, impaired memory, increased risk for heart disease, depression, and anxiety (McEwen, 2004). Further, individuals with a high allostatic load seek ways to return to equilibrium, and substance use may achieve this goal, at least initially (Cleck & Blendy, 2008; Wahler, 2012). However, over time addiction alters the way the way the body responds to stress, increasing allostatic load (Cleck & Blendy, 2008).

Individuals with a high allostatic load seek ways to return to equilibrium, and substance use may achieve this goal, at least initially.

The scale contains 15 symptoms and behaviors and asks clients to indicate how often they have experienced the symptoms/behaviors in the past 7 days. Examples of symptoms include: unexplained aches and pains, poor sleep, increased heart rate not related to exertion. Response options range from 0 (*None of the time*) to 3 (*All of the time*). The score is computed by summing the responses to all 15 items. Higher scores on the scale indicate greater physiological indicators of stress. The minimum score is 0 and the maximum score is 45. Internal consistency reliability was excellent in a sample of intake interviews completed by KTOS clients in FY 2014 ($n = 5,273$): Cronbach's $\alpha = 0.901$.

2. Economic and Living Circumstances

The economic and living circumstances examines; (1) living situation, (2) employment and disability status, and (3) economic hardship.

Prior research suggests that unemployment and lower socioeconomic status are important predictors of alcohol use relapse following treatment (Adamson, Sellman, & Frampton, 2009). In addition, one study found that individuals with higher resource needs (e.g., housing, employment, child care) were more likely to relapse 2 years after substance abuse treatment (Walton, Blow, Bingham, & Chermack, 2003). In FY 2012, using KTOS data,

Economic indicators and economic hardship are associated with higher stress as well as substance abuse treatment relapse.

a regression analysis showed that individuals who reported having difficulty meeting more basic needs were significantly more likely to report using alcohol and/or drugs at follow-up (Logan, Cole, Scrivner, & Spence, 2014). The high percentage of individuals who reported having trouble meeting basic needs

at both intake and follow-up shows that economic difficulties continue to be a problem for adults after they are in substance abuse treatment. Two other recent studies using KTOS data find that economic indicators and economic hardship are associated with higher stress as well as substance abuse treatment relapse (Wahler & Otis, 2014; Wahler, 2015). In another published study using KTOS data, economic hardship was associated with more stress (Cole, Logan, & Walker, 2011) and stress is associated with increased substance use and abuse and relapse (Sinha, 2008).

Assessing economic and living circumstances is important because Kentucky ranks as one of the highest states (48th in the nation) for poverty as well as the lowest for economic opportunity (Hess et al., 2015) while Gallup Polls (2014) ranked Kentucky as 46th in the nation for financial well-being (which considers having enough money for food, health care, and peoples perceived standard of living). Kentucky also was ranked 49th in the nation for children living in poverty (United Health Foundation, 2015).

LIVING SITUATION

This section assesses where the client has lived in the prior 12 months. The question and responses are adapted from the Government Performance and Reporting Act of 1993 (GPRA; Public Law 103-62) to ask about the past 12 months instead of the past 30 days (Mulvey, Atkinson, Avula, & Luckey, 2005) and whether they have been homeless or not.

EMPLOYMENT AND DISABILITY STATUS

The employment status questions were adapted from the ASI and the categories of type of work were adapted from the Standard Occupational Classification (U.S. Dept. of Labor, 2010). Disability status was included due to the high prevalence of disability in Kentucky. Using data from the 2013 American Community Survey (ACS) Kentucky had the 4th highest prevalence rate (16.1%) of disability among non-institutionalized working age individuals (ages 21 – 64) in the U.S. 50 states and territory of Puerto Rico (Erickson et al., 2014). Further, the Social Security Administration (2011) indicates 8.1% of the Kentucky population between 18 and 64 are on disability which is the 2nd highest in the nation.

ECONOMIC HARDSHIP

KTOS assessment includes a measure of economic hardship that was modified from the Survey of Income and Program Participation (SIPP), which is a multi-panel longitudinal nationally representative survey of the non-institutional population conducted by the U.S. Census Bureau. Information on economic hardship was collected as part of the eighth wave of data collection in the 1996 wave, which was in the field in 1998 (Beverly, 2001; Iceland & Bauman, 2004; She & Livermore, 2007). Economic hardship includes difficulty meeting basic needs including food, housing, clothing, and medical care (Beverly, 1999).

In the KTOS study, the telephone disconnection item was updated to consider difficulty with maintaining their cell phone cost given the prevalence of cell phones rather than landlines with many clients today. Additionally, the food insecurity item was adapted to a 12-month period. Finally, in the SIPP, the inability to obtain health care was measured with two items: needed to go to the doctor or

hospital but did not go, or needed to see a dentist but did not go. Because affordable access to prescription drugs is also an important dimension of health care that individuals may lack, an item was added to inquire about participants' inability to obtain a prescription drug because of financial problems.

Economic hardship was measured with two subscales in the KTOS study: 5 items measuring difficulty meeting basic living needs and 3 items measuring difficulty meeting health care needs. Good internal consistency reliability was found for the scale as a whole in the sample of KTOS clients who completed an intake interview in FY 2014 and were included in the 2016 Report (n = 5,273): Cronbach's $\alpha = 0.829$. Good internal consistency reliability was also found for the basic living needs (Cronbach's $\alpha = 0.756$) and health care needs (Cronbach's $\alpha = 0.826$) subscales.

3. Recovery Supports

The Recovery Supports section closes the KTOS assessment by asking about: (1) attending AA/NA/MA or other self-help group meetings and whether or not they have had contact with a sponsor recently; (2) how many people the client has they can count on to help them with their recovery and whether their friends or family were supportive of their recovery; and (3) what is most useful beside substance abuse treatment that helps them in their recovery and readiness to change (their perceived chances they can get off and stay off of drugs/alcohol).

Research has shown that recovery and positive social supports are linked to a lower risk of relapse. In addition, individuals in recovery cite their access to social and spiritual supports as an important key to their success.

The recovery supports questions were adapted from the GPRA (Mulvey et al., 2005) with feedback from discussions with state and community stakeholders. Research has shown that recovery and positive social supports are linked to a lower risk of relapse (Havassy, Hall, & Wasserman, 1991). In addition, individuals in recovery cite their access to social and spiritual supports as an important key to their success (Flynn, Joe, Broome, Simpson, & Brown, 2003). The last question in the KTOS assessment assesses readiness to change (Prochaska & DiClemente, 1983). However, measurement of readiness to change is often lengthy so this one-item was developed for KTOS assessments. One study using KTOS data found that client reported perceived chances they can get off and stay off drugs/alcohol (readiness to change) and 12-step program participation at follow-up was associated with positive treatment outcomes, while persistent depression was associated with negative treatment outcomes (Walker, Cole, & Logan, 2008).

KTOS demographic Information

The KTOS demographic information includes items that were taken or adapted slightly from the standardized Government Performance and Reporting Act of 1993 (GPRA; Public Law 103-62) monitoring tool, which is used by all Center for Substance Abuse Treatment (CSAT) and Substance abuse and Mental Health Services Administration (SAMHSA) funded grantees (Mulvey et al., 2005), or were included on KTOS as context specific questions: gender, race/ethnicity, age, marital status, education status, military experience, medical insurance type, and primary referral source.

Conclusion

The Kentucky Treatment Outcome Study (KTOS) is a statewide treatment outcome evaluation that is updated and enhanced annually. The KTOS assessment consists of three main components: (1) an evidence-based intake assessment administered by treatment staff using a secure, web-based instrument as clients enter publicly funded treatment programs; (2) an evidence-based follow-up assessment for a randomly selected sample of clients 12-months after intake. The follow-up rate is over 75% each year and over 1,250 clients are assessed at the 12-month follow-up each year in the past five years; and, (3) data analysis and dissemination.

The KTOS assessment is a brief self-report instrument that documents symptoms and patterns of substance abuse and related psychosocial problems. The KTOS is easy to use and takes about 30 minutes to complete. The KTOS assessment was developed in collaboration with key stakeholders and adapted to consider the Kentucky context as well as the unique substance abuse and related trends over time in Kentucky. The KTOS assessment has five core assessment components which all have strong reliability and validity research data including: (1) substance use, (2) mental health, (3) victimization and trauma; (4) criminal justice system involvement, and (5) quality of life. The three supplemental KTOS assessment components also have strong reliability and validity data for most of the assessment components and includes: (1) health and stress-related health consequences, (2) economic and living circumstances, and (3) recovery supports.

The evidence base for KTOS conforms to the 7 recommendations for evidence-based assessments for treatment providers in public agencies presented in the first section of this document.

- (1) *Use of Theory and Research.* The KTOS assessment includes a set of instruments developed to provide screening and assessment of psychosocial issues identified in theory and research as related to substance use including difficulties in employment, medical problems, housing instability, depression, anxiety, suicidality, criminal justice system involvement, and recovery supports (or engagement in the treatment process).
- (2) *Contextual Appropriateness.* The KTOS assessment was originally developed to consider the unique features of Kentucky and has been revised frequently after data analysis and feedback from users and other stakeholders to consider the unique context of Kentucky.
- (3) *Face Valid and User-friendly.* The KTOS assessment is face valid and focuses on components identified in theory and research as related to substance use, relapse, and treatment outcomes. Further, KTOS is easy to use and takes about 30 minutes to complete.
- (4) *Established Reliability and Validity.* The KTOS assessment has five core components (substance use, mental health, victimization and trauma, criminal justice system involvement, and quality of life) each with strong reliability and validity research support and three supplemental components (health and stress-related health consequences, economic and living circumstances, and recovery supports) many of which have strong reliability and validity research support.
- (5) *Measuring Dynamic Rather than Static Constructs.* Although KTOS does include key demographic indicators the majority of the assessment components focus on current

status, symptoms, and constructs that are amenable to change targeted in treatment over time.

- (6) *Not Producing Adverse Reactions or Consequences.* In the 20 years of conducting KTOS no adverse reactions or consequences due to the assessment or the research procedures have been reported.
- (7) *Sensitive to Change So That Outcomes Can Be Measured.* Results from past KTOS outcomes that a significant proportion of clients benefit from substance abuse treatment as substance use and substance abuse severity declines, quality of life is vastly improved, and criminal justice system involvement is significantly reduced.

Additional Benefit of Data Analysis and Dissemination. An added benefit of this Kentucky Treatment Outcome Study is that state-level trends in substance use along with the co-occurring anxiety and depression, criminal justice system involvement, employment and economic status, and quality of life trends for clients entering publicly funded treatment are provided each year. This data system also provides state-level trends in recovery and recovery correlates over time. An important benefit of state-level outcome studies is that funders and legislators can see up-to-date state specific data to provide evidence of need for new programs, continuation of current programs, and changes in programmatic policies. Key trends in substance use and policy needs fluctuate annually depending on economic and other state-specific sociopolitical issues, each year's analytical findings, the latest research, and legislative research commission requests, making the need for easily-modifiable annual data collection even more important. In addition to annual statewide reports, the KTOS data is used for community-level reports on client characteristics and outcomes for communities applying for Federal or other grants.

Key trends in substance use and policy needs fluctuate annually depending on economic and other state-specific sociopolitical issues, each year's analytical findings, the latest research, and legislative research commission requests, making the need for easily-modifiable annual data collection even more important.

The KTOS assessment is not meant to replace clinical decision-making or render diagnosis. The KTOS assessment can be used to inform treatment(s), engage clients through self-report, and monitor outcomes. The KTOS assessment, to minimize burden and cost, is not as lengthy, resource intensive, or as costly as other assessments. This may mean that if diagnosis information in general, or for a wider variety of conditions (e.g., personality disorder) is sought the KTOS assessment will need to be supplemented. Further, although the KTOS assessment is robust and pragmatic, it is relatively short (30 minutes) in order to reduce staff burden. That means that some of the substance use-related problems are not assessed while other components could be assessed more comprehensively.

The evidence base for the KTOS assessment suggests it is a robust, pragmatic, reliable, and valid assessment, which provides statewide and regional data about Kentucky drug use trends, substance use-related comorbidities, and substance abuse treatment outcomes.

Appendix A: References

- AAA Foundation for Traffic Safety (2016). *Prevalence of self-reported aggressive driving behavior: United States, 2014*. Washington, DC: AAA Foundation for Traffic Safety.
- Adamson, S., Sellman, J., & Frampton, C. (2009). Patient predictors of alcohol treatment outcome: A systematic review. *Journal of Substance Abuse Treatment, 36*, 75-86.
- American Psychiatric Association (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.
- Anda, R., Croft, J., Felitti, V., Nordenberg, D., Giles, W., Williamson, D., & Giovino, G. (1999). Adverse childhood experiences and smoking during adolescence and adulthood. *Journal of the American Medical Association, 282*, 1652-1658.
- Anda, R., Felitti, V., Walker, J., Whitfield, C., Bremner, J., Perry, B., Dube, S., & Giles, W. (2006). The enduring effects of abuse and related adverse experiences in childhood: a convergence of evidence from neurobiology and epidemiology. *European Archives of Psychiatry and Clinical Neurosciences, 56*(3), 174-86.
- Andresen, E., Catlin, T., Wyrwich, K., & Jackson-Thompson, J. (2003). Retest reliability of surveillance questions on health related quality of life. *Journal of Epidemiology and Community Health, 57*, 339-343.
- Appleby, L., Dyson, V., Altman, E., & Luchins, D. (1997). Assessing substance use in multiproblem patients: Reliability and validity of the Addiction Severity Index in a mental hospital population. *Journal of Nervous & Mental Disease, 185*(3), 159-165.
- Arrindell, W., Meeuwesen, L., & Huysse, F. (1991). The Satisfaction With Life Scale (SWLS): Psychometric properties in a non-psychiatric medical outpatients sample. *Personality and Individual Differences, 12*, 117-123.
- Atkinson, J., Slater, M., Patterson, T., Grant, I., & Garfin, S. (1991). Prevalence, onset, and risk of psychiatric disorders in men with chronic low back pain: A controlled study. *Pain, 45*(2), 111-121.
- Barrington-Trimis, J., Urman, R., Berhane, K., Unger, J., Cruz, T., Pentz, M., Samet, J., Leventhal, A., & McConnell, R. (2016). E-Cigarettes and future cigarette use. *Pediatrics, 138*(1), 1-8.
- Basco, M., Bostic, J., Davies, D., Rush, J., Witte, B., Hendrickse, W., & Barnett, V. (2000). Methods to improve diagnostic accuracy in a community mental health setting. *American Journal of Psychiatry, 157*, 1599-1605.
- Beidas, R., Stewart, R., & Walsh, L. (2015). Free, brief, and validated: Standardized instruments for low-resource mental health settings. *Cognitive and Behavioral Practice, 22*, 5-19.
- Beverly, S. (1999). *Economic poverty reconsidered: Material hardship and income-poverty in the United States*. Washington University, St. Louis, Missouri.
- Beverly, S. (2001). Material hardship in the United States: Evidence from the Survey of Income and Program Participation. *Social Work Research, 25*(3), 143-151.
- Blevins, C., Weathers, F., Davis, M., Witte, T., & Domino, J. (2015). The posttraumatic stress disorder checklist for DSM-5 (PCL-5). Development and initial psychometric evaluation. *Journal of Traumatic Stress, 28*, 489-498.
- Blumenstein, L. (2015). Exploratory study of intimate partner kidnapping: Is this a prevailing issue in intimate partner relationships? *Partner Abuse, 6*(4), 403-424.
- Bovin, M., Marx, B., Weathers, F., Gallagher, M., Rodriguez, P., Schnurr, P., & Keane, T. (in press). Psychometric properties of the PTSD Checklist for Diagnostic and Statistical Manual of Mental Disorders-Fifth Edition (PCL-5) in veterans. *Psychological Assessment*.

- Brady, K., Back, S., & Coffey, S. (2004). Substance abuse and posttraumatic stress disorder. *Current Directions in Psychological Science*, 13(5), 206-209.
- Breiding, M., Smith, S., Basile, K., Walters, M., Chen, J., & Merrick, M. (2014). Prevalence and characteristics of sexual violence, stalking, and intimate partner violence victimization—National intimate partner and sexual violence survey, United States, 2011. *Morbidity and Mortality Weekly Report*, 63(8), 1-18.
- Breslau, N., Kessler, R., Chilcoat, H., Schultz, L., Davis, G., & Andreski, P. (1998). Trauma and posttraumatic stress disorder in the community. The 1996 Detroit Area Survey of Trauma. *Archives of General Psychiatry*, 55(7), 626-632.
- Bumbarger, B., & Campbell, E. (2012). A state agency-university partnership for translational research and the dissemination of evidence-based prevention and intervention. *Administration and Policy in Mental Health Services Research*, 39(4), 268-277.
- Calsyn, D., Saxon, A., Bush, K., Howell, D., Baer, J., Sloan, K., Malte, C., & Kivlahan, D. (2004). The Addiction Severity Index medical and psychiatric composite scores measure similar domains as the SF-36 in substance-dependent veterans: Concurrent and discriminant validity. *Drug and Alcohol Dependence*, 76, 165-171.
- Campbell, D. (1959). Convergent and discriminant validation by the multitrait-multimethod matrix. *Psychological Bulletin*, 56, 81-105.
- Carey, K., Cocco, K., & Correia, C. (1997). Reliability and validity of the Addiction Severity Index among outpatients with severe mental illness. *Psychological Assessment*, 9(4), 422-428.
- Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration. *Treatment Episode Data Set (TEDS)*. Based on administrative data reported by states to TEDS through July 5, 2016. Retrieved from <http://www.dasis.samhsa.gov/webt/quicklink/KY13.htm>.
- Centers for Disease Control and Prevention (2000). *Measuring Healthy Days*. Atlanta, GA: CDC. Retrieved from: <https://www.cdc.gov/hrqol/pdfs/mhd.pdf>
- Chen, E., Turiano, N., Mroczek, D., & Miller, G. (2016). Association of reports of childhood abuse and all-cause mortality rates in women. *Journal of the American Medical Association*, 73(9), 920-927.
- Chermack, S., Roll, J., Reilly, M., Davis, L., Kilaru, U., & Gravowski, J. (2000). Comparison of patient self-reports and urinalysis results obtained under naturalist methadone treatment conditions. *Drug & Alcohol Dependence*, 59(1), 43-49.
- Christon, L., McLeod, B., & Jensen-Doss, A. (2015). Evidence-based assessment meets evidence-based treatment: An approach to science-informed case conceptualization. *Cognitive and Behavioral Practice*, 22, 36-48.
- Cleck, J., & Blendy, J. (2008). Making a bad thing worse: Adverse effects of stress on addiction. *Journal of Clinical Investigation*, 118(2), 454-461.
- Cleeland, C. (2009). The Brief Pain Inventory: A User's Guide. Retrieved from https://www.mdanderson.org/content/dam/mdanderson/documents/Departments-and-Divisions/Symptom-Research/BPI_UserGuide.pdf (accessed September 2, 2016).
- Cleeland, C., & Ryan, K. (1994). Pain assessment: Global use of the brief pain inventory. *Annals Academy of Medicine*, 23(2), 129-138.
- Cohen, M., & Klein, D. (1971). A measure of severity of multi-drug use among psychiatric patients. *International Journal of Pharmacopsychiatry*, 6, 83-89.

- Cole, J., Logan, T., & Walker, R. (2011). Social exclusion, self-regulation, and stress among substance abuse treatment clients. *Drug and Alcohol Dependence, 113*, 13-20.
- Connors, E., Arora, P., Curtis, L., & Stephan, S. (2015). Evidence-based assessment in school mental health. *Cognitive and Behavioral Practice, 22*(1), 60-73.
- Corse, S., Zanis, D., & Hirschinger, N. (1995). The use of the Addiction Severity Index with people with severe mental illness. *Psychiatric Rehabilitation Journal, 19*, 9-18.
- Cottler, L., Compton, W., Mager, D., Spitznagel, E., & Janca, A. (1992). Posttraumatic stress disorder among substance users from the general population. *American Journal of Psychiatry, 149*, 664-670.
- Cronbach, L., & Meehl, P. (1955). Construct validity in psychological tests. *Psychological Bulletin, 52*(4), 281-302.
- Daut, R., Cleeland, C., & Flanery, R. (1983). Development of the Wisconsin Brief Pain Questionnaire to assess pain in cancer and other diseases. *Pain, 17*, 197-210.
- Delgadillo, J., Payne, S., Gilbody, S., Godfrey, C., Gore, S., Jessop, D., & Dale, V. (2012). Brief case findings tools for anxiety disorders: Validation of GAD-7 and GAD-2 in addictions treatment. *Drug and Alcohol Dependence, 125*, 37-42.
- Denis, C., Fatséas, M., Beltran, V., Bonnet, C., Picard, S., Combourieu, I., Daulouède, J., & Auriacombe, M. (2012). Validity of the self-reported drug use section of the Addiction Severity Index and associated factors used under naturalistic conditions. *Substance use & Misuse, 47*, 356-363.
- Diener, E., Emmons, R., Larsen, R., & Griffin, S. (1985). The Satisfaction With Life Scale. *Journal of Personality Assessment, 49*, 71-75.
- Dietz, P., Spitz, A., Anda, R., Williamson, D., McMahon, P., Santelli, J., Nordenberg, D., Felitti, V., & Kendrick, J. (1999). Unintended pregnancy among adult women exposed to abuse or household dysfunction during their childhood. *Journal of the American Medical Association, 282*, 1359-1364.
- Dong, M., Anda, R., Felitti, V., Dube, S., Williamson, D., Thompson, T., Loo, C., & Giles, W. (2004). The interrelatedness of multiple forms of childhood abuse, neglect, and household dysfunction. *Child Abuse and Neglect, 28*, 771-784.
- Donley, S., Habib, L., Jovanovic, T., Kamkwala, A., Evces, M., Egan, G., Bradley, B., & Ressler, K. (2012). Civilian PTSD symptoms and risk for involvement in the criminal justice system. *Journal of the American Academy of Psychiatry and the Law, 40*(4), 522-529.
- Dube S., Anda, R., Felitti, V., Chapman, D., Williamson, D., & Giles, W. (2001). Childhood abuse, household dysfunction and the risk of attempted suicide throughout the life span: Findings from Adverse Childhood Experiences Study. *Journal of the American Medical Association, 286*, 3089-3096.
- Dube, S., Felitti, V., Dong, M., Chapman, D., Giles, W., & Anda, R. (2003a). Childhood abuse neglect and household dysfunction and the risk of illicit drug use: The Adverse Childhood Experiences Study. *Pediatrics, 111*, 564-572.
- Dube, S., Felitti, V., Dong, M., Giles, W., & Anda, R. (2003b). The impact of adverse childhood experiences on health problems: evidence from four birth cohorts dating back to 1900. *Preventative Medicine, 37*, 268-277.
- Dube, S., Miller, J., Brown, D., Giles, W., Felitti, V., Dong, M., & Anda, R. (2006). Adverse childhood experiences and the association with ever using alcohol and initiating alcohol use during adolescence. *Journal of Adolescent Health, 38*(4), 444.e1-10.
- Dube, S., Williamson, D., Thompson, T., Felitti, V., & Anda, R. (2004). Assessing the reliability of retrospective reports of adverse childhood experiences among adult HMO members attending a primary care clinic. *Child Abuse and Neglect, 28*(7), 729-737.

- Edlund, M., Sullivan, M., Han, X., & Booth, B. (2013). Days with pain and substance use disorders: Is there an association? *Clinical Journal of Pain, 29*(8), 689-695.
- Edwards, V., Anda, R., Dube, S., Dong, M., Chapman, D., & Felitti, V. (2005). The wide-ranging health consequences of adverse childhood experiences. In K. Kendall-Tackett, & S. Giacomoni (Eds.), *Child Victimization: Maltreatment, Bullying, and Dating Violence Prevention and Intervention*. Kingston, NJ: Civic Research Institute
- Erickson, W., Lee, C., & von Schrader, S. (2014). *2013 Disability Status report: Kentucky*. Ithaca, NY: Cornell University Employment and Disability Institute (EDI).
- Farley, M., Golding, J., Young, G., Mulligan, M., & Minkoff, J. (2004) Trauma history and relapse probability among patients seeking substance abuse treatment. *Journal of Substance Abuse Treatment, 27*(2), 161-167.
- Felitti, V., Anda, R., Nordenberg, D., Williamson, D., Spitz, A., Edwards, V., & Marks, J. (1998). The relationship of adult health status to childhood abuse & household dysfunction. *American Journal of Preventative Medicine, 14*(4), 245-258.
- First, M., Spitzer, R., Gibbon, M., & Williams, J. (1995). *Structured clinical interview for DSM-IV Axis I disorders (SCID-I)*. Washington, DC: American Psychiatric Association.
- Fleiss, J. (1981). *Statistical methods for rates and proportions*. New York: Wiley.
- Flynn, P., Joe, G., Broome, K., Simpson, D., & Brown, B. (2003). Recovery from opioid addiction in DATOS. *Journal of substance abuse treatment, 25*(3), 177-186
- Gallup Polls (2014). *State of American well-being: 2014 financial well-being rankings*. Retrieved from: http://info.healthways.com/hubfs/Well-Being_Index/2014_Data/Gallup-Healthways_State_of_American_Well-Being_2014_Financial_Rankings.pdf
- Gallup Polls (2015). *State of American well-being: 2015 state well-being rankings*. Retrieved from: http://info.healthways.com/hubfs/Well-Being_Index/2015_Data/Gallup-Healthways_State_of_American_Well-Being_2015_State_Rankings.pdf?t=1454003500629
- Glasgow, R. (2013). What does it mean to be pragmatic? Pragmatic methods, measures, and models to facilitate research translation. *Health Education & Behavior, 40*(3), 257-265.
- Goff, A., Rose, E., Rose, S., & Purves, D. (2007). Does PTSD occur in sentenced prison populations? A systematic literature review. *Criminal Behavior and Mental Health, 17*(3), 152-162.
- Gunderson, E., Russell, J., & Nail, R. (1973). A drug involvement scale for classification of drug abusers. *Journal of Comparative Psychology, 1*, 399-403.
- Hale, C. (1996). Fear of crime: A review of the literature. *International Review of Victimology, 4*, 79-150.
- Hasin, D., & Paykin, A. (1999). Alcohol dependence and abuse diagnoses: Concurrent validity in a nationally representative sample. *Alcoholism: Clinical and Experimental Research, 23*(1), 144-150.
- Hasin, D., Trautman, K., Miele, G., Samet, S., Smith, M., & Endicott, J. (1996). Psychiatric Research Interview for Substance and Mental Disorders (PRISM): Reliability for substance abusers. *American Journal of Psychiatry, 153*(9), 1195-1201.
- Havassy, B., Hall, S., & Wasserman, D. (1991). Social support and relapse: Commonalities among alcoholics, opiate users, and cigarette smokers. *Addictive Behaviors, 16*, 235-246.
- Hennessy, C., Moriarty, D., Zach, M., Scherr, P., & Brackbill, R. (1994). Measuring health-related quality of life for public health surveillance. *Public Health Reports, 109*(5), 665-672.

- Hess, C., Milli, J., Hayes, J., Hegewisch, A., Mayayeva, Y., Roman, S., Anderson, J., & Augeri, J. (2015). *The Status of Women in the States: 2015*. Institute for Women's Policy Research; Washington, DC.
- Hodgins, D., & El-Guebaly, N. (1992). More data on the Addiction Severity Index: Reliability and validity with the mentally ill substance abuser. *Journal of Nervous & Mental Disease*, 180(3), 197-201.
- Hogg, R., & Bertrand, D. (2004). What genes tell us about nicotine addiction. *Science*, 306(5698), 983-985.
- Horton, J., Compton, W., & Cottler, L. (2000). Reliability of substance use disorder diagnoses among African-Americans and Caucasians. *Drug and Alcohol Dependence*, 57(3), 203-209.
- Hunsley, J. (2015). Translating evidence-based assessment principles and components into clinical practice settings. *Cognitive and Behavioral Practice*, 22, 101-109.
- Hunsley, J., & Mash, E. (2007). Evidence-based assessment. *Annual Review of Clinical Psychology*, 3, 29-51.
- Iceland, J., & Bauman, K. (2004). Income poverty and material hardship: How strong is the association? *National Poverty Center Working Paper Series#04-17*. University of Michigan, Gerald R. Ford School of Public Policy.
- Ilies, R., Hauserman, N., Schwochau, S., & Stibal, J. (2003). Reported incidence rates of work-related sexual harassment in the United States: Using meta-analysis to explain reported rate disparities. *Personnel Psychology*, 56, 607-631.
- Jensen-Doss, A. (2015). Practical, evidence-based clinical decision making: Introduction to the special series. *Cognitive and Behavioral Practice*, 22, 1-4.
- Jensen-Doss, A., & Hawley, K. (2010). Understanding barriers to evidence-based assessment: Clinician attitudes toward standardized assessment tools. *Journal of Clinical Child & Adolescent Psychology*, 39(6), 885-896.
- Jensen-Doss, A., & Weisz, J. (2008). Diagnostic agreement predicts treatment process and outcomes in youth mental health clinics. *Journal of Consulting and Clinical Psychology*, 76(5), 711-722.
- Jensen-Doss, A., Youngstrom, E., Youngstrom, J., Feeny, N., & Findling, R. (2014). Predictors and moderators of agreement between clinical and research diagnoses for children and adolescents. *Journal of Consulting and Clinical Psychology*, 82(6), 1151-1162.
- Jewell, J., Handwerk, M., Almquist, J., & Lucas, C. (2004). Comparing the validity of clinician-generated diagnosis of conduct disorder to the Diagnostic Interview for Children. *Journal of Clinical Child and Adolescent Psychology*, 33(3), 536-546.
- Kearl, H. (2014). *Unsafe and harassed in public spaces: A national street harassment report*. Reston, VA: Stop Street Harassment. Retrieved from: <http://www.stopstreetharassment.org/wp-content/uploads/2012/08/National-Street-Harassment-Report-November-29-20151.pdf> (accessed on September, 28 2016).
- Kessler, R., Sonnega, A., Bromet, E., Hughes, M., & Nelson, C. (1995). Posttraumatic stress disorder in the National Comorbidity Survey. *Archives of General Psychiatry*, 52, 1048-1060.
- Kramer, T., Robbins, J., Phillips, S., Miller, T., & Burns, B. (2003). Detection and outcomes of substance use disorders in adolescents seeking mental health treatment. *Journal of the American Academy of Child & Adolescent Psychiatry*, 42(11), 1318-1326.
- Kroenke, K., Spitzer, R., & Williams, J. (2001). The PHQ-9: Validity of a brief depression severity measure. *Journal of General Internal Medicine*, 16, 606-613.

- Larsen, R., Diener, E., & Emmons, R. (1985). An evaluation of subjective well-being measures. *Social Indicators Research, 17*, 10-18.
- Logan, T. (2016). Personal safety concerns and responses. Unpublished data.
- Logan, T., Cole, J., Scrivner, A., & Spence, M. (2014). Adult Kentucky Treatment Outcome Study 2014 Annual Report. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.
- Logan, T. & Walker, R. (2010). Toward a deeper understanding of the harms caused by partner stalking. *Violence and Victims, 25*, 4, 440-455.
- Logan, T., Walker, R., Cole, J., & Leukefeld, C. (2002). Victimization and substance use among women: contributing factors, interventions, and implications. *Review of General Psychology, 6*(4), 325-397.
- Logan, T., Walker, R., Jordan, C., & Leukefeld, C. (2006). Women and victimization: Contributing factors, interventions, and implications. Washington, DC: American Psychological Association Press.
- Löwe, B., Decker, O., Müller, S., Brähler, E., Schellberg, D., Herzog, W., & Herzberg, P. (2008). Validation and standardization of the Generalized Anxiety Disorder Screener (GAD-7) in the general population. *Medical Care, 46*(3), 266-274.
- Mäkelä, K. (2004). Studies of the reliability and validity of the Addiction Severity Index. *Addiction, 99*, 398-410.
- Marquis, K. (1981). *Quality of prisoner self-reports: Arrest and conviction response errors*. Santa Monica: Rand Corporation.
- Maxfield, M., Weiler, B., & Widom, C. (2000). Comparing self-reports and official records of arrests. *Journal of Quantitative Criminology, 16*(1), 87-110.
- McEwen, B. (2000). The neurobiology of stress: From serendipity to clinical relevance. *Brain Research, 886*, 172-189.
- McEwen, B. (2004). Protection and damage from acute and chronic stress: Allostasis and allostatic overload and relevance to the pathophysiology of psychiatric disorders. *Annals of the New York Academy of Sciences, 1032*(1), 1-7.
- McLellan, A., Cacciola, J., Kushner, H., Peters, R., Smith, I., & Pettinati, H. (1992). The fifth edition of the Addiction Severity Index: Historical critique and normative data. *Journal of Substance Abuse Treatment, 9*, 199-213.
- McLellan, A., Luborsky, L., Cacciola, J., Griffith, B., Evans, R., Barr, H., & O'Brien, C. (1985). New data from the Addiction Severity Index: Reliability and validity in three centers. *Journal of Nervous and Mental Disease, 173*, 412-423.
- Mertens, J., Lu, Y., Parthasarathy, S., Moore, C., & Weisner, C. (2003). Medical and psychiatric conditions of alcohol and drug treatment patients in an HMO: Comparison with matched controls. *Archives of Internal Medicine, 163*, 2511-2517.
- Morgen, K., Astone-Twerell, J., Hernitche, T., Gunneson, L., & Santangelo, K. (2007). Health-related quality of life among substance abusers in residential drug abuse treatment. *Applied research in Quality of Life, 2*(4), 239-246.
- Murphy, A., Steele, M., Dube, S., Bate, J., Bonuck, K., Meissner, P., Goldman, H., & Steele, H. (2014). Adverse Childhood Experiences (ACEs) questionnaire and Adult Attachment Interview (AAI): Implications for parent child relationships. *Child Abuse & Neglect, 38*(2), 224-233.
- Mulvey, K., Atkinson, D., Avula, D., & Luckey, J. (2005). Using the internet to measure program performance. *American Journal of Evaluation, 26*(4), 587- 597.

- Najavits, L., Runkel, R., Neuner, C., Frank, A., Thase, M., Crits-Christoph, P., & Blaine, J. (2003). Rates and symptoms of PTSD among cocaine-dependent patients. *Journal of Studies on Alcohol*, 64, 601-606.
- Najavits, L., Sonn, J., Walsh, M., & Weiss, R. (2004). Domestic violence in women with PTSD and substance abuse. *Addictive Behavior*, 29, 707-715.
- Newschaffer, C. (1998). *Validation of Behavioral Risk Factor Surveillance System (BRFSS) HRQOL measures in a statewide sample*. Atlanta: U.S. Department of health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion. [http://www.cdc.gov/hrqol/measurement_properties/newschaffer.htm].
- Nunno, M. (2006). The effects of the ARC organizational intervention on caseworker turnover, climate, and culture in children's services systems: Commentary. *Child Abuse and Neglect*, 30(8), 849-854.
- Office for Victims of Crime (2015). Section 6: Statistical Overviews. In *2015 National Crime Victims' Rights Week (NCVRW) Resource Guide*. Washington, DC: Office of Justice Programs, US Department of Justice. Retrieved from: <http://ovc.ncjrs.gov/ncvrw2015/pdf/FullGuide.pdf> (accessed on September, 28 2016).
- Pavot, W., & Diener, E. (1993). Review of the Satisfaction with Life Scale. *Psychological Assessment*, 5(2), 164-172.
- Perreault, S. (2015). Criminal victimization in Canada, 2014. *Juristat*, 35(1), 1-43.
- Peters, R., Sherman, P., & Osher, F. (2008). Treatment in jails and prisons. In K.T. Mueser, & D.V. Jeste (Eds.), *Clinical Handbook of Schizophrenia*, 354-364. New York, NY: Guilford Press.
- Pogge, D., Wayland-Smith, D., Zaccario, M., Borgaro, S., Stokes, J., & Harvey, P. (2001). Diagnosis of manic episodes in adolescent inpatients: Structured diagnostic procedures compared to clinical chart diagnoses. *Psychiatry Research*, 101(1), 47-54.
- Price, M., Szafranski, D., van Stolk-Cooke, K., & Gros, D. (2016). Investigation of an abbreviated 4 and 8 item version of the PTSD Checklist 5. *Psychiatry Research*, 239, 124-130.
- Prochaska, J., & DiClemente, C. (1983). Stages and processes of self-change of smoking: Toward an integrative model of change. *Journal of Consulting and Clinical Psychology*, 51(3), 390.
- Regier, D., Farmer, M., Rae, D., Locke, B., Keith, S., & Judd, L. (1990). Comorbidity of mental disorders with alcohol and other drug abuse: Results from the Epidemiologic Catchment Area (ECA) Study. *Journal of the American Medical Association*, 264, 2511-2518.
- Rikoon, S., Cacciola, J., Carise, D., Alterman, A., & McLellan, A. (2006). Predicting DSM-IV dependence diagnoses from Addiction Severity Index composite scores. *Journal of Substance Abuse Treatment*, 31, 17-24.
- Rudd, R., Aleshire, N., Zibbell, J., & Gladden, R. (2016). Increases in drug and opioid overdose deaths—United States, 2000-2014. *Morbidity and Mortality Weekly Report*, 64(50), 1378-1382.
- Sadeh, N., & McNiel, D. (2015). Posttraumatic stress disorder increases risk of criminal recidivism among justice-involved persons with mental disorders. *Criminal Justice and Behavior*, 42(6), 573-586.
- Sansone, R., & Sansone, L. (2010). Road rage: What's driving it? *Psychiatry (Edgemont)*, 7(7), 14-18.
- Scott, K., & Lewis, C. (2015). Using measurement-based care to enhance any treatment. *Cognitive and Behavioral Practice*, 22, 49-59.
- Selzer, M. (1971). The Michigan Alcoholism Screening Test: The quest for a new diagnostic instrument. *American Journal of Psychiatry*, 127, 1653-1658.

- Shane, P., Diamond, G., Mensinger, J., Shera, D., & Wintersteen, M. (2006). Impact of victimization on substance abuse treatment outcomes for adolescents in outpatient and residential substance abuse treatment. *The American Journal on Addictions, 15*(1), 34-42.
- Shannon, L., Havens, J., Mateyoke-Scrivner, A., & Walker, R. (2009). Contextual differences in substance use for rural Appalachian treatment-seeking women. *The American Journal of Drug and Alcohol Abuse, 35*(2), 59-62.
- She, P., & Livermore, G. (2007). Material hardship, poverty, and disability among working-age adults. *Social Science Quarterly, 88*(4), 970-989.
- Sheu, R., Lussier, D., Rosenblum, A., Fong, C., Portenoy, J., Joseph, H., & Portenoy, R. (2008). Prevalence and characteristics of chronic pain in patients admitted to an outpatient drug and alcohol treatment program. *Pain Medicine, 9*11-917.
- Singh, T., Arrazola, R., Corey, C., Husten, C., Neff, L., Homa, D., & King, B. (2016). Tobacco use among middle and high school students—United States, 2011-2015. *MMWR Morbidity and Mortality Weekly Report, 65*, 361-367.
- Sinha, R. (2008). Chronic stress, drug use, and vulnerability to addiction. *New York Academy of Sciences, 1141*, 105-130.
- Slavova, S., Bunn, T., & Gao, W. (2015). *Drug overdose hospitalizations in Kentucky, 2000-2013*. Lexington, KY: Kentucky Injury Prevention and Research Center.
- Smart, R., Mann, R., & Stoduto, G. (2003). The prevalence of road rage: Estimates from Ontario. *Canadian Journal of Public Health, 94*(4), 247-250.
- Social Security Administration (2011). *Annual statistical report on the Social Security Disability Insurance Program, 2011*. Retrieved from: https://www.ssa.gov/policy/docs/statcomps/di_asr/2011/sect01.html
- Spitzer, R., Kroenke, K., Williams, J., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: The GAD-7. *Archives of Internal Medicine, 166*, 1092-1097.
- Stevenson, E., Cole, J., Walker, R., and Logan, T. (2014). Association of chronic non-cancer pain with substance abuse treatment outcomes among a Community Mental Health Center sample. *Addictive Disorders and Their Treatment, 13*(1), 30-37. doi: 10.1097/ADT.0b013e31827b0cd9.
- Stockdale, M., Logan, T., Sliter, K., & Berry, S. (2014). Interpersonal Violence Victimization and Sexual Harassment: A Prospective Study of Revictimization. *Sex Roles, 71*, 55-70.
- Stöffelmayr, B., Mavis, B., & Kasim, R. (1994). The longitudinal stability for the Addiction Severity Index. *Journal of Substance Abuse Treatment, 11*(4), 373-378.
- Substance Abuse and Mental Health Services Administration (SAMHSA). (2015). *2013-2014 National Survey on Drug Use and Health: Model-based prevalence estimates (50 states and the District of Columbia)*. Rockville, MD: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality.
- Tenney, N., Schotte, C., Denys, D., van Megen, H., & Westenberg, H. (2003). Assessment of DSM-IV personality disorders in obsessive-compulsive disorder: Comparison of clinical diagnosis, self-report questionnaire, and semi-structured interview. *Journal of Personality Disorders, 17*(6), 550-561.
- Truman, J., & Langton, L. (2015). *Criminal victimization, 2014*. (NCJ 248973). Washington, DC: U.S. Department of Justice, Office of Justice Statistics, Bureau of Justice Statistics.

- United Health Foundation (2015). *America's health rankings: Annual report*. Retrieved from http://cdnfiles.americashealthrankings.org/SiteFiles/Reports/2015AHR_Annual-v1.pdf (accessed September 7, 2016).
- U.S. Department of Labor (2010). *2010 Standard Occupational Classification Major Groups*. Retrieved from http://www.bls.gov/soc/major_groups.htm
- Vitaliano, P., Russo, J., Young, H., Becker, J., & Maiuro, R. (1991). The screen for caregiver burden. *The Gerontologist*, *31*, 76-83.
- Wahler, E. (2012). *The relationship of social stress, economic hardship, and psychological distress to addiction severity among Kentucky substance abuse treatment participants* (Doctoral dissertation). Retrieved from http://uknowledge.uky.edu/csw_etds/1
- Wahler, E. (2015). Social disadvantage and economic hardship as predictors of follow-up addiction severity after substance abuse treatment: Does referral to treatment by the criminal justice system matter? *Alcoholism Treatment Quarterly*, *33*(1), 6-27.
- Wahler, E., & Otis, M. (2014). Social disadvantage, economic hardship, and psychological distress as predictors of sustained abstinence from substance use after treatment. *Substance Use and Misuse*, *49*(13), 1820-1832.
- Walker, R., Cole, J., & Logan, T. (2008). Identifying client-level indicators of recovery among DUI, criminal justice, and non-criminal justice treatment referrals. Special issue - *Substance Use and Misuse*, *43*, 1785-1801.
- Walton, M., Blow, F., Bingham, C., & Chermack, S. (2003). Individual and social/environmental predictors of alcohol and drug use 2 years following substance abuse treatment. *Addictive Behaviors*, *28*(4), 627-642.
- Weathers, F., Litz, B., Keante, T., Palmieri, P., Marx, B., & Schnurr, P. (2013). *The PTSD Checklist for DSM-5 (PCL-5)*. Boston, MA: National Center for PTSD.
- Wertz, J., Cleaveland, B., & Stephens, R. (1995). Problems in the application of the Addiction Severity Index (ASI) in rural substance abuse services. *Journal of Substance Abuse*, *7*(2), 175-188.
- Wolff, N., & Shi, J. (2012). Childhood and adult trauma experiences of incarcerated persons and their relationship to adult behavioral health problems and treatment. *International Journal of Environmental Research and Public Health*, *9*(5), 1908-1926.
- World Health Organization. (2014). *Global status report on alcohol and health* (2014 ed.). Geneva, Switzerland: World Health Organization Press.
- Wortmann, J., Alexander, J., Weathers, F., Resick, P., Dondanville, K., Hall-Clark, B., & Litz, B. (in press). Psychometric analysis of the PTSD Checklist-5 (PCL-5) among treatment-seeking military service members. *Psychological Assessment*.
- Zanis, D., McLellan, A., Randall, M. (1994). Can you trust patient self-reports of drug use during treatment? *Drug & Alcohol Dependence*, *35*, 127-132.
- Zanis, D., McLellan, T., & Corse, S. (1997). Is the Addiction Severity Index a reliable and valid assessment instrument among clients with severe and persistent mental illness and substance abuse disorders? *Community Mental Health Journal*, *33*(3), 213-227.

Appendix B: Reports Generated Using KTOS Data

In addition to the annual report submitted to the state, regional reports and other ad hoc data reports are generated upon request.

Annual Reports

Cole, J., Logan, TK, Miller, J., & Scrivner, A. (2016). *Adult Kentucky Treatment Outcome Study 2016 Annual Report*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Cole, J., Logan, TK, Scrivner, A. & Miller, J. (2015). *Adult Kentucky Treatment Outcome Study 2015 Annual Report*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Logan, TK, Cole, J., Scrivner, A. & Spence, M. (2014). *Adult Kentucky Treatment Outcome Study 2014 Annual Report*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Cole, J., Logan, TK, Scrivner, A., Stevenson, E. (2013). *Findings from the Kentucky Treatment Outcome Study 2013 Report*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Cole, J., Walker, R., Stevenson, E., Mateyoke-Scrivner, A., Logan, TK. (2012). *Kentucky Substance Abuse Treatment Outcome Study 2012 Annual Follow-up Report*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Mateyoke-Scrivner, A., Cole, J., Logan, TK, Stevenson, E., & Jackson, T. (2011). *Kentucky Substance Abuse Treatment Outcome Study 2011 Annual Follow-up Report*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Mateyoke-Scrivner, A., Cole, J., Logan, TK, Stevenson, E., & Jackson, T. (2010). *Kentucky Substance Abuse Treatment Outcome Study 2010 Annual Follow-up Report*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Mateyoke-Scrivner, A., Cole, J., Logan, TK, Stevenson, E., & Jackson, T. (2009). *Treatment Effectiveness in Kentucky: Kentucky Treatment Outcome Study 2009 Annual Follow-up Report*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Mateyoke-Scrivner, A., Cole, J., Logan, TK, Stevenson, E., Leukefeld, C., & Jackson, T. (2008). *Kentucky Substance Abuse Treatment Outcome Study 2008 Annual Follow-up Report*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Mateyoke-Scrivner, A., Cole, J., Logan, TK, Stevenson, E., Leukefeld, C., & Jackson, T. (2007). *Kentucky Substance Abuse Treatment Outcome Study 2007 Annual Follow-up Report*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Mateyoke-Scriver, A., Cole, J., Logan, TK, Stevenson, E., & Leukefeld, C. (2006). *Kentucky Substance Abuse Treatment Outcome Study 2006 Annual Follow-up Report*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Mateyoke-Scriver, A., Bradshaw, G., Logan, TK, Stevenson, E., Leukefeld, C., & Goltz, M. (2005). *Kentucky Substance Abuse Treatment Outcome Study FY 2003 Follow-up Findings*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Logan, TK, Bradshaw, G., Leukefeld, C., Goltz, M., & Stevenson, E. (2004). *Kentucky Substance Abuse Treatment Outcome Study FY 2002 Follow-up Findings*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Morgan, K., & Leukefeld, C. (2003). *Kentucky Substance Abuse Treatment Outcome Study 2000 Follow-up Findings*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Morgan, K., & Leukefeld, C. (2002). *Kentucky Substance Abuse Treatment Outcome Study 1999 Follow-up Findings*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Morgan, K., & Leukefeld, C. (2001). *Kentucky Substance Abuse Treatment Outcome Study 1998 Follow-up Findings*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Morgan, K., & Leukefeld, C. (1999). *Kentucky Substance Abuse Treatment Outcome Study Follow-up Findings October 1996 Through December 1997*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Select Regional and Other Ad Hoc Reports

Cole, J., Scriver, A., & Miller, J. (2016) *Kentucky Treatment Outcome Study RiverValley Behavioral Health Clients at Treatment Intake July 1, 2010 – June 30, 2014*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Comparison of KTOS Clients' Self-Reported Substance Use in the 12 Months Before Entering Treatment: Bluegrass Vs. Other Regions. (2016). Center on Drug and Alcohol Research, University of Kentucky.

Kentucky Treatment Outcome Study Intake Data for FY 2014 AOC Data Request: Hardin, McCracken, and Perry Counties. (2016). Center on Drug and Alcohol Research, University of Kentucky.

Kentucky Treatment Outcome Study Intake Data for FY 2014 AOC Data Request: Allen, Harrison, Nicholas, Robertson, and Simpson Counties. (2016). Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Miller, J., Cole, J., & Scrivner, A. (2016). *Kentucky Adult Treatment Outcome Study Program Report: Adanta*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Miller, J., Scrivner, A. & Cole, J. (2016). *Kentucky Adult Treatment Outcome Study Program Report: Four Rivers*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Miller, J., Scrivner, A. & Cole, J. (2016). *Kentucky Adult Treatment Outcome Study Program Report: Genesis*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Cole, J., & Miller, J. (2016). *Kentucky Adult Treatment Outcome Study Program Report: Comprehend, Inc.* Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Miller, J., & Cole, J. (2016). *Kentucky Adult Treatment Outcome Study Program Report: Bluegrass*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Miller, J., & Cole, J. (2016). *Kentucky Adult Treatment Outcome Study Program Report: Kentucky River*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Miller, J., & Cole, J. (2016). *Kentucky Adult Treatment Outcome Study Program Report: Pathways*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Miller, J., & Cole, J. (2016). *Kentucky Adult Treatment Outcome Study Program Report: Seven Counties*. Center on Drug and Alcohol Research, University of Kentucky.

Cole, J., & Miller, J. (2015) *Kentucky Treatment Outcome Study RiverValley Behavioral Health Clients at Treatment Intake July 1, 2010 – June 30, 2013*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Kentucky Treatment Outcome Study Intake Data for FY 2013 Christian County Residents. (2015). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Kentucky Treatment Outcome Study Intake Data for FY 2013 Floyd County Residents. (2015). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Cole, J. (2014). *Kentucky Treatment Outcome Study: Brief Report for River Valley Behavioral Health*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research

Cole, J. (2014). *Increasing Heroin Use in the Seven Counties Region*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Cole, J. (2014). *Substance Use Among Daviess County Clients Entering Publicly-Funded Substance Abuse Treatment*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Logan, T., Scrivner, A., Spence, M., & Cole, J. (2014). *Kentucky Adult Treatment Outcome Study Program Report: The Adanta Group*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Spence, M., & Cole, J. (2014). *Kentucky Adult Treatment Outcome Study Program Report: Bluegrass*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Cole, J., Spence, M., & Scrivner, A. (2014). *Kentucky Adult Treatment Outcome Study Program Report Draft: Communicare*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Spence, M., & Cole, J. (2014). *Kentucky Adult Treatment Outcome Study Program Report: Comprehend, Inc.* Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Spence, M., & Cole, J. (2014). *Kentucky Adult Treatment Outcome Study Program Report Draft: Cumberland River Comprehensive Care*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Spence, M., & Cole, J. (2014). *Kentucky Adult Treatment Outcome Study Program Report: Four Rivers Behavioral Health*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Spence, M., & Cole, J. (2014). *Kentucky Adult Treatment Outcome Study Program Report: Kentucky River Community Care*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Spence, M., & Cole, J. (2014). *Kentucky Adult Treatment Outcome Study Program Report: LifeSkills*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Spence, M., & Cole, J. (2014). *Kentucky Adult Treatment Outcome Study Program Report: Mountain Comprehensive Care*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Spence, M., & Cole, J. (2014). *Kentucky Adult Treatment Outcome Study Program Report: NorthKey Community Care*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Spence, M., & Cole, J. (2014). *Kentucky Adult Treatment Outcome Study Program Report: Pathways, Inc.* Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Spence, M., & Cole, J. (2014). *Kentucky Adult Treatment Outcome Study Program Report: Pennyroyal Center*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Spence, M., & Cole, J. (2014). *Kentucky Adult Treatment Outcome Study Program Report: RiverValley Behavioral Health*. Center on Drug and Alcohol Research, University of Kentucky.

Logan, T., Scrivner, A., Spence, M., & Cole, J. (2014). *Kentucky Adult Treatment Outcome Study Program Report: Seven Counties Services*. Center on Drug and Alcohol Research, University of Kentucky.

Kentucky Treatment Outcome Study Preliminary Profile for Individuals in Treatment at Seven Counties Services: Substance Use, Employment, and Criminal Justice System Involvement. (2012). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Lifekills Regional Report: Kentucky Treatment Outcome Study FY2010. (2012). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY2011 Intake Information Comparing RiverValley and Lifeskills. (2011). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Mateyoke-Scrivner, A., & Parrish, D. (2010). *Cumberland River Comprehensive Care Special Report on Substance Abuse Treatment Outcomes*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Mateyoke-Scrivner, A., & Parrish, D. (2010). *Pathways Special Report on Substance Abuse Treatment Outcomes*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Adult KTOS Draft Analysis for Mercer County Site Number 901038. (2009). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY05-FY08 Intake and Follow-up Comparison for Pathways. (2009). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY2009 Intake Comparison of Head Injury for Communicare Clients. (2009). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY2009 Intake: Kentucky River vs. Seven Counties and Western Regions. (2009). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

KTOS FY08 Intake- Kentucky River Compared to the Rest of the State. (2009). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

KTOS FY08 Intake- Seven Counties Compared to the Rest of the State. (2009). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Mateyoke-Scrivner, A., Stevenson, E., Walker, R. (2009). *Special Brief Report: RiverValley Stimulant Use Trends Using Kentucky Treatment Outcome Study Data Collected Between SFY2005 and SFY2008*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Mateyoke-Scriver, A., Cole, J., Logan, TK, Stevenson, E. (2008). *Special Report for Bluegrass Regional Mental Health-Mental Retardation Board, Inc.: Substance Abuse Treatment Outcomes*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Mateyoke-Scriver, A., Cole, J., Logan, TK, Stevenson, E. (2008). *Special Report for RiverValley Behavioral Health: Substance Abuse Treatment Outcomes*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Kentucky Substance Abuse Treatment Outcome Study Kentucky River Profile: Changes in Substance Use, Employment, and Criminal Justice Involvement Fiscal Years 2003-2005. (2007). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Kentucky Substance Abuse Treatment Outcome Study Mountain Region Profile: Changes in Substance Use, Employment, and Criminal Justice Involvement Fiscal Years 2003-2005. (2007). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

KTOS Special Report Seven Counties, Inc. Profile of Treatment Outcomes: Substance Use, Employment, and Criminal Justice Involvement Fiscal Years 2003-2005. (2007). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Regional Reports: Change from Intake to Follow-up. (2006). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY 2002 and FY 2003 Shepard's House Intake Data. (2005). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY 2004 Preliminary Analysis: Adanta Group. (2005). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY 2004 Preliminary Analysis: Bluegrass. (2005). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY 2004 Preliminary Analysis: Communicare. (2005). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY 2004 Preliminary Analysis: Comprehend. (2005). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY 2004 Preliminary Analysis: Cumberland River. (2005). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY 2004 Preliminary Analysis: Four Rivers. (2005). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY 2004 Preliminary Analysis: Kentucky River. (2005). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY 2004 Preliminary Analysis: LifeSkills. (2005). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY 2004 Preliminary Analysis: Mountain. (2005). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY 2004 Preliminary Analysis: NorthKey. (2005). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY 2004 Preliminary Analysis: Pathways. (2005). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY 2004 Preliminary Analysis: Pennyroyal. (2005). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY 2004 Preliminary Analysis: Seven Counties. (2005). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

FY 2004 Preliminary Analysis: Volta House. (2005). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Stevenson, E., Goltz, M. & Cole, C. (2005). *Special Substance Abuse Report: Adanta Region.* Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Stevenson, E., Goltz, M. & Cole, C. (2005). *Special Substance Abuse Report: Bluegrass Region.* Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Stevenson, E., Goltz, M. & Cole, C. (2005). *Special Substance Abuse Report: Communicare Region.* Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Stevenson, E., Goltz, M. & Cole, C. (2005). *Special Substance Abuse Report: Comprehend Region.* Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Stevenson, E., Goltz, M. & Cole, C. (2005). *Special Substance Abuse Report: Cumberland River Region.* Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Stevenson, E., Goltz, M. & Cole, C. (2005). *Special Substance Abuse Report: Four Rivers Region.* Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Stevenson, E., Goltz, M. & Cole, C. (2005). *Special Substance Abuse Report: Kentucky River Region.* Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Stevenson, E., Goltz, M. & Cole, C. (2005). *Special Substance Abuse Report: LifeSkills Region*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Stevenson, E., Goltz, M. & Cole, C. (2005). *Special Substance Abuse Report: Mountain Region*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Stevenson, E., Goltz, M. & Cole, C. (2005). *Special Substance Abuse Report: NorthKey Region*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Stevenson, E., Goltz, M. & Cole, C. (2005). *Special Substance Abuse Report: Pathways Region*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Stevenson, E., Goltz, M. & Cole, C. (2005). *Special Substance Abuse Report: Pennyroyal Region*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Stevenson, E., Goltz, M. & Cole, C. (2005). *Special Substance Abuse Report: River Valley Region*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R., Stevenson, E., Goltz, M. & Cole, C. (2005). *Special Substance Abuse Report: Seven Counties Region*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Kentucky Substance Abuse Treatment Outcome Study Preliminary Findings for Region 1: Four Rivers FY 2002 Report. (2004). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Special Substance Abuse Report: Adanta Group Region. (2004). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Special Substance Abuse Report: Bluegrass Region. (2004). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Special Substance Abuse Report: Four Rivers Region. (2004). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Report for Shepard's House. (2003). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Kentucky Substance Abuse Treatment Outcome Study Preliminary Findings for Region 3: River Valley 2000 Report. (2002). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Kentucky Substance Abuse Treatment Outcome Study Preliminary Findings for Region 4: Lifeskills 2000 Report. (2002). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Kentucky Substance Abuse Treatment Outcome Study Preliminary Findings for Region 14: The Adanta Group 2000 Report. (2002). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Kentucky Substance Abuse Treatment Outcome Study Preliminary Findings for Region 15: Bluegrass Regional Mental Health-Mental Retardation Board, Inc. 2000 Report. (2002). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Cumberland River Regional Mental Health/Mental Retardation Board, Inc. Substance Abuse Outcomes Process. (2000). Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Walker, R. (2000). *Jefferson County 1999 Substance Abuse Treatment Data: Women in Treatment.* Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Translational reports

Logan, TK, Cole, J., Miller, J. & Scrivner, A. (2016). *Adult Kentucky Treatment Outcome Study 2016 Findings at a Glance.* Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Logan, TK, Cole, J., Miller, J. & Scrivner, A. (2016). *Adult Kentucky Treatment Outcome Study 2016 Fact Sheet.* Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Cole, J., Logan, TK, Scrivner, A. & Miller, J. (2015). *Adult Kentucky Treatment Outcome Study 2015 Findings at a Glance.* Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Logan, TK, Cole, J., Scrivner, A. (2015). *Adult Kentucky Treatment Outcome Study 2015 Fact Sheet.* Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Logan, TK, Cole, J., Scrivner, A., and Spence, M. (2014). *Adult Kentucky Treatment Outcome Study 2014 Findings at a Glance.* Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Logan, TK, Cole, J., Scrivner, A., & Spence, M. (2014). *Adult Kentucky Treatment Outcome Study 2014 Fact Sheet.* Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Logan, TK, Cole, J., Scrivner, A. & Stevenson, E. (2013). *Kentucky Treatment Outcome Study 2013 Report: Findings at a Glance.* Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Logan, TK, Cole, J., Scrivner, A. & Stevenson, E. (2013). *2013 Adult Kentucky Treatment Outcome Study Fact Sheet.* Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Logan, TK, Scrivner, A., Parish, D., Stevenson, E., & Cole, J. (2012). *KTOS 2012 Outcomes Report: Findings at a Glance.* Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Cole, J., Logan, TK, Mateyoke, A., & Stevenson, E. (2011). *Kentucky Treatment Outcome Study 2011*. Lexington, KY: University of Kentucky, Center on Drug & Alcohol Research.

Cole, J., Stevenson, E., Mateyoke-Scriver, A., Logan, T., & Walker, R. (2011). Tobacco Use among Individuals in Substance Abuse Treatment in Kentucky. *KTOS In-Focus, Winter 2010*, 1-4. Center on Drug & Alcohol Research, University of Kentucky.

Scriver, A., Walker, R., Stevenson, E., Cole, J. & Logan, TK. (2011). Regional differences in substance use. *KY Treatment In-Focus*, 3(5), 1-4. Center on Drug & Alcohol Research, University of Kentucky.

Mateyoke-Scriver, A., Walker, R., Logan, TK, Cole, J., Stevenson, E. (2010). Risk factors for suicidality among substance abuse treatment clients. *KTOS In-Focus*, 3(1), 1-4. Center on Drug & Alcohol Research, University of Kentucky.

Walker, R., Mateyoke-Scriver, A., Cole, J., Logan, TK, Stevenson, E., & Jackson, T. (2009). *Treatment Effectiveness in Kentucky: 2009 Follow-up Findings Summary*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Cole, J., Walker, R., Logan, TK, Stevenson, E., & Mateyoke-Scriver, A. (2009). Following substance abuse treatment clients over time: Tips from the front lines. *KTOS In-Focus*, 2(1), 1-6. Center on Drug & Alcohol Research, University of Kentucky.

Cole, J., Walker, R., Stevenson, E., Logan, TK, & Mateyoke-Scriver, A. (2009). Lifetime victimization and mental health among women in state-funded substance abuse treatment. *KTOS In-Focus*, 2(4), 1-4. Center on Drug & Alcohol Research, University of Kentucky.

Mateyoke-Scriver, A., Walker, R., Cole, J., Stevenson, E., Logan, TK, & Shannon, L. (2009). Examining prescription opiate abuse in Kentucky: A look at an emerging trend. *KTOS In-Focus*, 2(3), 1-4. Center on Drug & Alcohol Research, University of Kentucky.

Stevenson, E., Shannon, L., Walker, R., Mateyoke-Scriver, A., Logan, TK., & Cole, J. (2008). Kentucky women in substance abuse treatment: Rural Appalachia. *KTOS In-Focus*, 1 (2), 1-4. Center on Drug & Alcohol Research, University of Kentucky.

Stevenson, E., Walker, R., Cole, J., Logan, TK, Mateyoke-Scriver, A., & Corrigan, J. (August, 2008). Substance use treatment clients with traumatic brain injury. *KTOS In-Focus*, 1 (1), 1-4. Center on Drug & Alcohol Research, University of Kentucky.

Walker, R., Mateyoke-Scriver, A., Cole, J., Logan, TK, Stevenson, E., Leukefeld, C., & Jackson, T. (2007). *Kentucky Substance Abuse Treatment Outcome Study 2007 Follow-up Summary Report*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Mateyoke-Scriver, A., Cole, J., Logan, TK, Stevenson, E., & Leukefeld, C. (2006). *Kentucky Substance Abuse Treatment Outcome Study 2006 Annual Follow-up Summary Report*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Mateyoke-Scriver, A., Bradshaw, G., Logan, TK, Stevenson, E., Leukefeld, C., & Goltz, M. (2005). *Kentucky Substance Abuse Treatment Outcome Study Summary Report FY 2003*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Walker, R., Logan, TK, Bradshaw, G., Leukefeld, C., Goltz, M., & Stevenson, E. (2004). *Kentucky Substance Abuse Treatment Outcome Study FY 2002 Summary of Statewide Follow-Up Findings*. Lexington, KY: University of Kentucky, Center on Drug and Alcohol Research.

Appendix C: KTOS Publications

There are eight publications from KTOS published in peer reviewed journals and one additional publication under review. A dissertation has also been completed using the KTOS data.

1. Victor, G., Walker, R., Cole, J., & Logan, T. (2017). Opioid analgesics and heroin: Examining drug misuse trends among a sample of drug treatment clients in Kentucky. *In Press Journal of International Drug Policy*.
2. Wahler, E. (2015). Social disadvantage and economic hardship as predictors of follow-up addiction severity after substance abuse treatment: Does referral to treatment by the criminal justice system matter? *Alcoholism Treatment Quarterly*, 33(1), 6-27.
3. Wahler, E. & Otis, M. (2014). Social disadvantage, economic hardship, and psychological distress as predictors of sustained abstinence from substance use after treatment. *Substance Use and Misuse*, 49(13), 1820-1832.
4. Cole, J., Logan, T., & Walker, R. (2011). Social exclusion, self-regulation, and stress among substance abuse treatment clients. *Drug and Alcohol Dependence*, 113, 13-20.
5. Shannon, L., Havens, J., Mateyoke-Scriver, A., & Walker, R. (2009). Contextual differences in substance use for rural Appalachian treatment-seeking women. *The American Journal of Drug and Alcohol Abuse*, 35(2), 59-62.
6. Walker, R., Cole, J., & Logan, T. (2008). Identifying client-level indicators of recovery among DUI, criminal justice, and non-criminal justice treatment referrals. Special issue - *Substance Use and Misuse*, 43, 1785-1801.
7. Walker, R., Cole, J., Logan, T. (2007). Screening substance abuse treatment clients for traumatic brain injury: Prevalence and characteristics. *Journal of Head Trauma Rehabilitation*, 22, 360-368.
8. Walker, R., Hiller, M., Staton-Tindall, M., & Leukefeld, C. (2003). Head injury among drug abusers: An indicator of co-occurring problems. *Journal of Psychoactive Drugs*, 35(3).
9. Walker, R., Staton, M., & Leukefeld, C. (2001). History of head injury among substance users: Preliminary findings. *Substance Use and Misuse*, 36(6-7), 757-768.
10. Wahler, E. A. (2012). *The relationship of social stress, economic hardship, and psychological distress to addiction severity among Kentucky substance abuse treatment participants* (Doctoral dissertation). Retrieved from http://uknowledge.uky.edu/csw_etds/1